

BL

## (12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property Organization  
International Bureau(43) International Publication Date  
12 June 2003 (12.06.2003)

PCT

(10) International Publication Number  
WO 03/048130 A2(51) International Patent Classification<sup>7</sup>: C07D 233/00

(21) International Application Number: PCT/US02/36128

(22) International Filing Date:  
26 November 2002 (26.11.2002)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:  
60/334,453 30 November 2001 (30.11.2001) US(71) Applicant (for all designated States except US): **ELI LILLY AND COMPANY** [US/US]; Lilly Corporate Center, Indianapolis, IN 46285 (US).

(72) Inventors; and

(75) Inventors/Applicants (for US only): **GIBSON, Tracey, Ann** [US/US]; 4416 Par Drive, Indianapolis, IN 46268 (US). **JOHNSTON, Richard, Duane** [US/US]; 3712 Forest Lane, Greenfield, IN 46140 (US). **MANTLO, Nathan, Bryan** [US/US]; 7325 East 800 North, Brownsburg, IN 46112 (US). **THOMPSON, Richard, Craig** [US/US]; 763 North 900 West, Frankfort, IN 46041 (US). **WANG, Xiaodong** [CN/US]; 14066 Honey Tree Drive, Carmel, IN 46032 (US). **WINNEROSKI, Leonard, Larry, Junior** [US/US]; 587 Thoroughbred Lane, Greenwood, IN 46142 (US). **XU, Yanping** [CN/US]; 13832 Barnett Place, Fishers, IN 46038 (US).(74) Agents: **VORNDRAN-JONES, MaCharri et al.**; Eli Lilly and Company, P. O. Box 6288, Indianapolis, IN 46206-6288 (US).

(81) Designated States (national): AE, AG, AL, AM, AT (utility model), AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU, CZ (utility model), CZ, DE (utility model), DE, DK (utility model), DK, DM, DZ, EC, EE (utility model), EE, ES, FI (utility model), FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NO, NZ, OM, PH, PL, PT, RO, RU, SC, SD, SE, SG, SI, SK (utility model), SK, SL, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, YU, ZA, ZM, ZW.


(84) Designated States (regional): ARIPO patent (GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, SK, TR), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).

## Declaration under Rule 4.17:

— as to applicant's entitlement to apply for and be granted a patent (Rule 4.17(ii)) for the following designations AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NO, NZ, OM, PH, PL, PT, RO, RU, SC, SD, SE, SG, SI, SK, SL, TJ, TM, TN, TR, TT, TZ, UA, UG, UZ, VC, VN, YU, ZA, ZM, ZW, ARIPO patent (GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, SK, TR), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG)

[Continued on next page]

(54) Title: PEROXISOME PROLIFERATOR ACTIVATED RECEPTOR AGONISTS



C<sub>1</sub>-C<sub>4</sub> alkyl amide, OR10 and C<sub>3</sub>-C<sub>6</sub> cycloalkyl; (c) W is O or S; (d) X is an optionally substituted C<sub>1</sub>-C<sub>5</sub> alkylene linker wherein one carbon atom of the linker may optionally be replaced with O, NH, S, and optionally two carbons together may form a double bond; (e) Y is selected from the group consisting of C, O, S, NH and a single bond; and (f) E is selected from the group consisting of C(R<sub>3</sub>) (R<sub>4</sub>) A, A, and a substituted or unsubstituted group selected from the group consisting of (CH<sub>2</sub>)<sub>n</sub> COOR<sub>19</sub>.

(57) Abstract: The present invention is directed to compounds represented by the following structural Formula (I), (a) R<sub>1</sub> is selected from the group consisting of hydrogen, substituted or unsubstituted group selected from C<sub>1</sub>-C<sub>8</sub> alkyl, aryl-C<sub>0.4</sub>-alkyl, heteroaryl-C<sub>0.4</sub>-alkyl, C<sub>3</sub>-C<sub>6</sub> cycloalkyl, aryl-C<sub>0.2</sub>-alkyl, and -CH<sub>2</sub>-C(O)-R<sub>17</sub>-R<sub>18</sub>, wherein R<sub>17</sub> is O or NH and R<sub>18</sub> is optionally substituted benzyl; (b) R<sub>2</sub> is selected from the group consisting of C<sub>1</sub>-C<sub>6</sub> alkyl, C<sub>1</sub>-C<sub>6</sub> alkenyl, aryl-C<sub>0.4</sub>-alkyl, heteroaryl-C<sub>0.4</sub>-alkyl, C<sub>1</sub>-C<sub>4</sub> alkyl sulfonamide,

WO 03/048130 A2

38

WO 03/048130 A2



**Published:**

— without international search report and to be republished  
upon receipt of that report

*For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.*

**PEROXISOME PROLIFERATOR ACTIVATED RECEPTOR AGONISTS****BACKGROUND OF THE INVENTION**

Peroxisome Proliferator Activated Receptors (PPARs) are  
5 members of the nuclear hormone receptor super family, which  
are ligand-activated transcription factors regulating gene  
expression. Various subtypes of PPARs have been discovered.  
These include, for example, PPAR $\alpha$ , PPAR $\gamma$  and PPAR $\delta$ .

PPAR $\alpha$ , PPAR $\gamma$  and PPAR $\delta$  receptors have been  
10 implicated in diabetes mellitus, cardiovascular disease,  
obesity, Syndrome X and gastrointestinal disease, such as,  
inflammatory bowel disease. Syndrome X is the combination  
of symptoms which include hyperinsulemia combined with  
hypertension, elevated body weight, elevated triglycerides  
15 and elevated LDL.


Current PPAR agonist treatment for Syndrome X relates  
to the use of thiazolidinediones (TZDs) or other insulin  
sensitivity enhancers (ISEs). TZDs are a class of PPAR  
gamma agonists which have been shown to increase the  
20 sensitivity of insulin sensitive cells. Increasing insulin  
sensitivity rather than the amount of insulin in the blood  
reduces the likelihood of hypoglycemic coma. However, TZDs  
and ISEs typically have little effect in preventing the  
cardiovascular part of Syndrome X in that their  
25 administration usually does not result in the lowering of  
triglycerides and LDL-cholesterol while raising HDL-  
cholesterol. Furthermore, clinically significant side  
effects are commonly associated with treatment with TZDs.  
Therefore, a need exists for new pharmaceutical agents which  
30 affect treat or prevent cardiovascular disease, particularly  
that associated with Syndrome X, while preventing or  
minimizing weight gain, and more preferably while improving

insulin sensitivity. The present invention provides novel compounds exhibiting desired pharmacological activity.

#### SUMMARY OF THE INVENTION

The present invention is directed to compounds represented by the following structural formula:

Formula I



and pharmaceutically acceptable salts, solvates and hydrates thereof, wherein:

- (a) R1 is selected from the group consisting of hydrogen, substituted or unsubstituted group selected from C<sub>1</sub>-C<sub>8</sub> alkyl, aryl-C<sub>0-4</sub>-alkyl, heteroaryl-C<sub>0-4</sub>-alkyl, C<sub>3</sub>-C<sub>6</sub> cycloalkylaryl-C<sub>0-2</sub>-alkyl, and -CH<sub>2</sub>-C(O)-R17-R18, wherein R17 is O or NH and R18 is optionally substituted benzyl;
- (b) R2 is H or a substituted or unsubstituted group selected from the group consisting of C<sub>1</sub>-C<sub>6</sub> alkyl, C<sub>1</sub>-C<sub>6</sub> alkenyl, aryl-C<sub>0-4</sub>-alkyl, heteroaryl-C<sub>0-4</sub>-alkyl, C<sub>1</sub>-C<sub>4</sub> alkyl sulfonamide, C<sub>1</sub>-C<sub>4</sub> alkyl amide, OR10 and C<sub>3</sub>-C<sub>6</sub> cycloalkyl;
- (c) W is O or S;
- (d) X is an optionally substituted C<sub>1</sub>-C<sub>5</sub> alkylene linker wherein one carbon atom of the linker may optionally be replaced with O, NH, S, and



optionally two carbons together may form a double bond;

(e) Y is selected from the group consisting of C, O, S, NH and a single bond; and

5 (f) E is selected from the group consisting of C(R3)(R4)A, A, and a substituted or unsubstituted group selected from the group consisting of (CH<sub>2</sub>)<sub>n</sub>COOR<sub>19</sub>; and wherein

(i) n is 0, 1, 2 or 3;

10 (ii) A is an functional group selected from the group consisting of carboxyl, C<sub>1</sub>-C<sub>3</sub>alkylnitrile, carboxamide, substituted or unsubstituted sulfonamide, substituted or unsubstituted acylsulfonamide substituted or  
15 unsubstituted tetrazole, and substituted or unsubstituted isoxazole;

(iii) R<sub>3</sub> is selected from the group consisting of H, C<sub>1</sub>-C<sub>5</sub> alkyl, arylC<sub>0</sub>-C<sub>2</sub>alkoxy, and C<sub>1</sub>-C<sub>5</sub> alkoxy, and

20 (iv) R<sub>4</sub> is selected from the group consisting of H, halo, and a substituted or unsubstituted group selected from C<sub>1</sub>-C<sub>5</sub> alkyl, C<sub>1</sub>-C<sub>5</sub> alkoxy, C<sub>3</sub>-C<sub>6</sub> cycloalkyl, aryl C<sub>0</sub>-C<sub>4</sub> alkyl, arylC<sub>0</sub>-C<sub>2</sub>alkoxy and phenyl; or R<sub>3</sub> and R<sub>4</sub> are  
25 combined to form a C<sub>3</sub>-C<sub>8</sub> cycloalkyl;

(v) R<sub>19</sub> is selected from the group consisting of hydrogen, optionally substituted arylmethyl and optionally substituted C<sub>1</sub>-C<sub>4</sub>alkyl;


30 (g) R<sub>8</sub> is selected from the group consisting of hydrogen, C<sub>1</sub>-C<sub>4</sub> alkyl, C<sub>1</sub>-C<sub>4</sub> alkylenyl, and halo;

(h) R<sub>9</sub> is selected from the group consisting of hydrogen, C<sub>1</sub>-C<sub>4</sub> alkyl, C<sub>1</sub>-C<sub>4</sub> alkylenyl, halo,

substituted or unsubstituted aryl, substituted or unsubstituted aryl-C<sub>1</sub>-C<sub>4</sub> alkyl, substituted or unsubstituted heteroaryl, C<sub>1</sub>-C<sub>6</sub> alkenyl, and OR<sub>10</sub>;

- (i) R<sub>10</sub> is independently selected from the group consisting of hydrogen and C<sub>1</sub>-C<sub>4</sub> alkyl;
- (j) R<sub>21</sub> is selected from the group consisting of hydrogen, =O, and a substituted or unsubstituted group selected from the group consisting of C<sub>1</sub>-C<sub>6</sub> alkyl, aryl, C<sub>1</sub>-C<sub>4</sub> alkylaryl, and heteroaryl;
- (k) R<sub>22</sub> is selected from the group consisting of hydrogen, and a substituted or unsubstituted group selected from the group consisting of C<sub>1</sub>-C<sub>6</sub> alkyl, aryl, C<sub>1</sub>-C<sub>4</sub> alkylaryl, and heteroaryl; and
- (l) ---- represents an optional double bond.

A compound of Formula I'



and pharmaceutically acceptable salts, solvates and hydrates thereof, wherein:


- (a) R<sub>1</sub> is selected from the group consisting of hydrogen, substituted or unsubstituted group selected from C<sub>1</sub>-C<sub>8</sub> alkyl, aryl-C<sub>0-4</sub>-alkyl, heteroaryl-C<sub>0-4</sub>-alkyl, C<sub>3</sub>-C<sub>6</sub> cycloalkylaryl-C<sub>0-2</sub>-alkyl, and -CH<sub>2</sub>-C(O)-R<sub>17</sub>-R<sub>18</sub>, wherein R<sub>17</sub> is O or NH and R<sub>18</sub> is optionally substituted benzyl;
- (b) R<sub>2</sub> is H or a substituted or unsubstituted group

selected from the group consisting of C<sub>1</sub>-C<sub>6</sub> alkyl, C<sub>1</sub>-C<sub>6</sub> alkenyl, aryl-C<sub>0-4</sub>-alkyl, heteroaryl-C<sub>0-4</sub>-alkyl, C<sub>1</sub>-C<sub>4</sub> alkyl sulfonamide, C<sub>1</sub>-C<sub>4</sub> alkyl amide, OR10 and C<sub>3</sub>-C<sub>6</sub> cycloalkyl;


- 5 (c) W is O or S;
- (d) X is an optionally substituted C<sub>1</sub>-C<sub>5</sub> alkylene linker wherein one carbon atom of the linker may be replaced with O, NH or S;
- (e) Y is C, O, S, NH or a single bond; and
- 10 (f) E is selected from the group consisting of C(R3)(R4)A, A, substituted or unsubstituted selected from the group consisting of (CH<sub>2</sub>)<sub>n</sub> COOR19; and wherein
- (i) n is 0, 1, 2 or 3,
- 15 (ii) A is an functional group selected from the group consisting of carboxyl, C<sub>1</sub>-C<sub>3</sub>alkylnitrile, carboxamide, substituted or unsubstituted sulfonamide, substituted or unsubstituted acylsulfonamide and substituted
- 20 or unsubstituted tetrazole;
- (iii) R3 is H, saturated or unsaturated C<sub>1</sub>-C<sub>5</sub> alkyl, arylC<sub>0-2</sub>alkoxy, C<sub>1</sub>-C<sub>5</sub> alkoxy, and
- (iv) R4 is H, halo, a substituted or unsubstituted group selected from C<sub>1</sub>-C<sub>5</sub> alkyl, C<sub>1</sub>-C<sub>5</sub>
- 25 alkoxy, C<sub>3</sub>-C<sub>6</sub> cycloalkyl, aryl C<sub>0-4</sub> alkyl, arylC<sub>0-2</sub>alkoxy and phenyl, or R3 and R4 are combined to form a C<sub>3</sub>-C<sub>8</sub> cycloalkyl;
- (v) R19 is selected from the group consisting of hydrogen, optionally substituted arylmethyl
- 30 and optionally substituted C<sub>1</sub>-C<sub>4</sub>alkyl;

- (g) R8 is selected from the group consisting of hydrogen, C<sub>1</sub>-C<sub>4</sub> alkyl, C<sub>1</sub>-C<sub>4</sub> alkylenyl, and halo;
- (h) R9 is selected from the group consisting of hydrogen, C<sub>1</sub>-C<sub>4</sub> alkyl, C<sub>1</sub>-C<sub>4</sub> alkylenyl, halo, substituted or unsubstituted aryl, substituted or unsubstituted aryl-C<sub>1</sub>-C<sub>4</sub> alkyl, substituted or unsubstituted heteroaryl, C<sub>1</sub>-C<sub>6</sub> alkenyl, and OR10;
- (i) R10 is independently selected from the group consisting of hydrogen and C<sub>1</sub>-C<sub>4</sub> alkyl;
- (j) R21 is selected from the group consisting of hydrogen, =O, and a substituted or unsubstituted group selected from the group consisting of C<sub>1</sub>-C<sub>6</sub> alkyl, aryl, C<sub>1</sub>-C<sub>4</sub> alkylaryl, and heteroaryl;
- (k) R22 is selected from the group consisting of hydrogen, and a substituted or unsubstituted group selected from the group consisting of C<sub>1</sub>-C<sub>6</sub> alkyl, aryl, C<sub>1</sub>-C<sub>4</sub> alkylaryl, and heteroaryl; and
- (l) ---- represents an optional double bond.


An additional embodiment is a compound and pharmaceutically acceptable salts, solvates and hydrates of Structural Formula II:



An additional embodiment is a compound and pharmaceutically acceptable salts, solvates and hydrates of Structural Formula III:



Another embodiment of this invention is a compound of Formula IV represented by the following structure:



5 and salts, solvates and hydrates thereof.

One preferred embodiment of this invention is a compound selected from the group consisting of 2-Methyl-3-(4-{2-[3-methyl-2-oxo-1-(4-trifluoromethyl-benzyl)-imidazolidin-4-yl]-ethoxy}-phenyl)-2-phenoxy-propionic acid  
 10 and 2-methyl-3-(4-{3-[3-methyl-2-oxo-1-(4-trifluoromethyl-benzyl)-imidazolidin-4-yl]-propyl}-phenyl)-2-phenoxy-propionic acid.

In another feature of this invention, a compound claimed herein is radiolabeled.

15 In one embodiment, the present invention also relates to pharmaceutical compositions which comprising at least one compound of the present invention, or a pharmaceutically acceptable salt, solvate, hydrate or prodrug thereof, and a pharmaceutically acceptable carrier.

20 In another embodiment, the present invention relates to a method of modulating one or more PPAR receptor(s) by contacting the receptor with at least one compound

represented by Structural Formula I, and pharmaceutically acceptable salts, solvates and hydrates thereof.

In another embodiment, the present invention relates to a method of selectively modulating a PPAR receptor by  
5 contacting the receptor with at least one compound represented by Structural Formula I, and pharmaceutically acceptable salts, solvates and hydrates thereof. Another embodiment of this invention is preferentially modulating one PPAR receptor and additionally modulating a second,  
10 different, PPAR receptor to provide desired dual agonism.

In a further embodiment, the present invention relates to a method of making a compound represented by Structural Formula I.

The compounds of the present invention and  
15 pharmaceutically acceptable salts, solvates and hydrates thereof are believed to be effective in treating and preventing Syndrome X, Type II diabetes, hyperglycemia, hyperlipidemia, obesity, coagulopathy, hypertension, atherosclerosis, and other disorders related to Syndrome X  
20 and cardiovascular diseases. In addition, the compounds exhibit fewer side effects than compounds currently used to treat these conditions. Further, compounds of this invention can be useful for lowering fibrinogen, increasing HDL levels, treating renal disease, controlling desirable  
25 weight, treating demyelinating diseases, treating certain viral infections, and treating liver disease.

#### DETAILED DESCRIPTION OF THE INVENTION

The terms used to describe the instant invention have  
30 the following meanings herein.

As used herein, alkyl groups include straight chained or branched hydrocarbons, which are completely saturated.

Said branched hydrocarbon may be primary, secondary, tertiary, or quaternary, as appropriate.


As used herein, alkylene linker is a C<sub>1</sub>-C<sub>5</sub> straight or branched chain hydrocarbon group. However, the term  
5 "alkylene linker wherein one carbon of the linker may optionally be replaced with O, NH, S, and optionally two carbons together may form a double bond" refers to an alkylene linking having an O, NH or S in the link. Alkylene linkers having two carbons together forming a double bond  
10 means for example, -CH-CHCH<sub>2</sub>-, -CH<sub>2</sub>CHCH-, -CH<sub>2</sub>CH<sub>2</sub>CHCH-, and the like. For example, but not limited to, -CH<sub>2</sub>CH<sub>2</sub>O-, -CH<sub>2</sub>CH<sub>2</sub>NH-, CH<sub>2</sub>CH<sub>2</sub>S-, -CH<sub>2</sub>OCH<sub>2</sub>-, -OCH<sub>2</sub>CH<sub>2</sub>-, and the like.

Cycloalkyl groups, as used herein, include cyclic hydrocarbons, which are partially or completely saturated.

15 As used herein, aryl groups include carbocyclic aromatic ring systems (e.g. phenyl), fused polycyclic aromatic ring systems (e.g. naphthyl and anthracenyl) and aromatic ring systems fused to carbocyclic non-aromatic ring systems (e.g., 1,2,3,4-tetrahydronaphthyl and benzodioxyl).


20 Heterocyclic group, as used herein, is a ring system having at least one heteroatom such as nitrogen, sulfur or oxygen. Heterocyclic groups include benzofuranyl, benzothiazolyl, benzothienyl, isoquinolyl, isoxazolyl, morpholino, oxadiazolyl, pyridyl, pyrimidinyl, pyrrolyl,  
25 quinolyl, tetrahydropyranyl and thienyl.

Heteroaryl, as used herein is an aromatic ring system, fused polycyclic aromatic ring systems and aromatic ring systems fused to non-aromatic ring systems wherein at least one carbon atom is replaced with a heteroatom such as  
30 nitrogen, sulfur or oxygen. It may be preferred that heteroaryl contains from 1-3 heteroatoms. It may be preferred that heteroaryl contains 1-2 nitrogen atoms. One



heteroaryl group that may be desired is

Another heteroaryl group that may be desired is:



The term "arylalkyl", "arylmethyl", "arylC<sub>0</sub>-C<sub>2</sub>alkoxy" and "aryloxy" each represent a substituent in which the aryl group is linked to the parent molecule via the alkyl, methyl, and oxy, respectively. Additionally, when the phrase "arylC<sub>0</sub>-C<sub>2</sub>alkoxy" is C<sub>0</sub> alkoxy, this means that the aryl is linked to the parent molecule via an oxy group.

Examples of R<sub>1</sub>, R<sub>5</sub>, E, R<sub>19</sub> and R<sub>9</sub> suitable substituents when said R<sub>1</sub>, E, R<sub>5</sub>, R<sub>19</sub> or R<sub>9</sub> are at least one or more independently selected from the group consisting of C<sub>1</sub>-C<sub>8</sub>alkyl, C<sub>1</sub>-C<sub>4</sub> alkyl, aryl, arylmethyl, (CH<sub>2</sub>)<sub>n</sub>COOR<sub>19</sub>, C<sub>1</sub>-C<sub>6</sub> alkenyl, thio-C<sub>1</sub>-C<sub>4</sub>alkyl, thioaryl, C<sub>1</sub>-C<sub>4</sub>alkoxyaryl, C<sub>1</sub>-C<sub>4</sub>alkoxy C<sub>1</sub>-C<sub>4</sub>alkyl, aminoaryl, aminoC<sub>1</sub>-C<sub>4</sub>alkyl, aryl-C<sub>0</sub>-C<sub>4</sub>alkyl, heteroarylC<sub>0</sub>-C<sub>4</sub>alkyl, heterocyclic, -CH<sub>2</sub>-C(O)-R<sub>17</sub>-R<sub>18</sub>, (C<sub>3</sub>-C<sub>6</sub>)cycloalkylaryl-C<sub>0</sub>-C<sub>2</sub>-alkyl and cycloalkyl, then suitable substituted groups include, for example, C<sub>1</sub>-C<sub>5</sub> alkyl, C<sub>1</sub>-C<sub>5</sub> alkoxy, C<sub>0</sub>-C<sub>5</sub> haloalkyl, C<sub>1</sub>-C<sub>5</sub> trihaloalkyl, C<sub>1</sub>-C<sub>5</sub> trihaloalkoxy, C<sub>1</sub>-C<sub>5</sub> haloalkoxy, nitro, cyano, CHO, =O, hydroxyl, C<sub>1</sub>-C<sub>4</sub> alkanolic acid, phenyl, aryloxy, SO<sub>2</sub>R<sub>7</sub>, SR<sub>7</sub>, benzyloxy, alkylcarboxamido or COOH. R<sub>7</sub> is an alkyl or a haloalkyl. When R<sub>1</sub>, R<sub>5</sub>, E, R<sub>19</sub> or R<sub>9</sub> is substituted, it is preferred that there are from 1-3 substitutions on said R<sub>1</sub>, R<sub>5</sub>, E, R<sub>19</sub> or R<sub>9</sub> group. An especially preferred trihaloalkyl group is trifluoro C<sub>1</sub>-C<sub>5</sub> alkyl.

Examples of suitable substituents for an "optionally substituted C<sub>2</sub>-C<sub>5</sub> alkylene linker," include one or more



independently selected from the group consisting of C<sub>1</sub>-C<sub>6</sub>alkyl, oxo, substituted or unsubstituted arylC<sub>0</sub>-C<sub>3</sub>alkyl, C<sub>1</sub>-C<sub>3</sub>alkoxy, hydroxy, C<sub>3</sub>-C<sub>6</sub>cycloalkyl and halo. When the alkylene linker is substituted, it is preferred that there  
5 are from one to three independent substitutions.

Examples of suitable substituents for a substituted C<sub>1</sub>-C<sub>3</sub> alkylene, include one or more independently selected from C<sub>1</sub>-C<sub>6</sub>alkyl, oxo, aryl C<sub>0</sub>-C<sub>3</sub>alkyl, C<sub>1</sub>-C<sub>3</sub>alkoxy, hydroxy, and halo. When the alkylene is substituted it is preferred that  
10 there are from 1-3 independent substitutions.

Suitable substituents for substituted R<sub>2</sub> groups wherein R<sub>2</sub> is C<sub>1</sub>-C<sub>6</sub>alkyl, C<sub>1</sub>-C<sub>6</sub>alkenyl, arylC<sub>0</sub>-C<sub>4</sub>alkyl, arylC<sub>0</sub>-C<sub>4</sub>alkyl, C<sub>1</sub>-C<sub>4</sub> alkyl sulfonamide, C<sub>1</sub>-C<sub>4</sub> alkyl amide, OR<sub>10</sub>, or C<sub>3</sub>-C<sub>6</sub>cycloalkyl, include for example, one or more  
15 independently selected from the group consisting of OH, alkoxy, haloalkyl, amino, COOH, heteroaryl-O-, heteroaryl-C(O)-, alkyl-O-, alkyl-C(O)-, C<sub>3</sub>-C<sub>6</sub> cycloalkyl, aryl-O-, aryl-C(O)-, heteroaryl, aryl, heterocycloalkyl, heterocycloalkyl-O-, and heterocycloalkyl-C(O)-. When R<sub>2</sub> is  
20 substituted it is preferred that there are from 1-3 independent substitutions on the R<sub>2</sub> group.

Examples of suitable substituents for A groups, wherein the A is a sulfonamide, include one or more independently selected from C<sub>1</sub>-C<sub>4</sub> alkyl, C<sub>1</sub>-C<sub>4</sub> haloalkyl, substituted or  
25 unsubstituted heteroaryl, or substituted or unsubstituted aryl. When the A group is substituted, it is preferred that there are from 1-3 independent substitutions on the A group.

Examples of suitable substituents for A groups, wherein A is acylsulfonamide and tetrazole include, for example, one  
30 or more independently selected from C<sub>1</sub>-C<sub>4</sub> alkyl, C<sub>1</sub>-C<sub>4</sub> haloalkyl, substituted or unsubstituted heteroaryl, or substituted or unsubstituted aryl.

Suitable substituents for R4 wherein R4 is C<sub>1</sub>-C<sub>5</sub> alkyl, C<sub>1</sub>-C<sub>5</sub>alkoxy, C<sub>1</sub>-C<sub>6</sub>cycloalkyl, arylC<sub>0</sub>-C<sub>4</sub>alkyl, arylC<sub>0</sub>-C<sub>2</sub>alkoxy or phenyl, include, for example halo, phenyl, C<sub>1</sub>-C<sub>4</sub> alkoxy, hydroxy, and arylC<sub>0</sub>-C<sub>2</sub>alkoxy. When R4 is substituted, it is  
5 preferred that there are from 1-4 independently selected substitutions on the R4 group.

Preferably, for the compounds of the present invention, represented by Structural Formula I, and with their respective pharmaceutical compositions, W is an oxygen.

10 The compounds of Structural Formula I may contain one or more chiral centers, and exist in different optically active forms. When compounds of Structural Formula I contain one chiral center, the compounds exist in two enantiomeric forms and the present invention includes both  
15 enantiomers and mixtures of enantiomers, such as racemic mixtures. The enantiomers may be resolved by methods known to those skilled in the art, for example by formation of diastereoisomeric salts which may be separated, for example, by crystallization; formation of diastereoisomeric  
20 derivatives or complexes which may be separated, for example, by crystallization, gas-liquid or liquid chromatography; selective reaction of one enantiomer with an enantiomer-specific reagent, for example enzymatic esterification; or gas-liquid or liquid chromatography in a  
25 chiral environment, for example on a chiral support for example silica with a bound chiral ligand or in the presence of a chiral solvent. It will be appreciated that where the desired enantiomer is converted into another chemical entity by one of the separation procedures described above, a  
30 further step is required to liberate the desired enantiomeric form. Alternatively, specific enantiomers may be synthesized by asymmetric synthesis using optically

active reagents, substrates, catalysts or solvents, or by converting one enantiomer into the other by asymmetric transformation.

When a compound represented by Structural Formula I has  
5 more than one chiral substituent it may exist in diastereoisomeric forms. The diastereoisomeric pairs may be separated by methods known to those skilled in the art, for example chromatography or crystallization and the individual enantiomers within each pair may be separated as described  
10 above. The present invention includes each diastereoisomer of compounds of Structural Formula I and mixtures thereof.

Certain compounds of Structural Formula I may exist in different stable conformational forms which may be separable. Torsional asymmetry due to restricted rotation  
15 about an asymmetric single bond, for example because of steric hindrance or ring strain, may permit separation of different conformers. The present invention includes each conformational isomer of compounds of Structural Formula I and mixtures thereof.

20 Certain compounds of Structural Formula I may exist in zwitterionic form and the present invention includes each zwitterionic form of compounds of Structural Formula I and mixtures thereof.

Certain compounds of Structural Formula I and their  
25 salts may also exist in the form of solvates, for example hydrates, and the present invention includes each solvate and mixtures thereof.

"Pharmaceutically-acceptable salt" refers to salts of the compounds of the Structural Formula I which are  
30 substantially non-toxic to mammals. Typical pharmaceutically acceptable salts include those salts prepared by reaction of the compounds of the present

invention with a mineral or organic acid or an organic or inorganic base. Such salts are known as base addition salts, respectively. It should be recognized that the particular counterion forming a part of any salt of this invention is not of a critical nature, so long as the salt as a whole is pharmaceutically acceptable and as long as the counterion does not contribute undesired qualities to the salt as a whole.

By virtue of its acidic moiety, a compound of Structural Formula I forms salts with pharmaceutically acceptable bases. Some examples of base addition salts include metal salts such as aluminum; alkali metal salts such as lithium, sodium or potassium; and alkaline earth metal salts such as calcium and magnesium; and ammonium or substituted ammonium salts. Examples of substituted ammonium salts include, for instance, those with lower alkylamines such as trimethylamine, triethylamine; hydroxyalkylamines such as 2-hydroxyethylamine, bis-(2-hydroxyethyl)-amine or tri-(2-hydroxyethyl)-amine, cycloalkylamines such as bicyclohexylamine or dibenzylpiperidine, N-benzyl- $\beta$ -phenethylamine, dehydroabietylamine, N,N'-bisdehydro-abietylamine, glucamine, N-methylglucamine; bases of the pyridine type such as pyridine, collidine, quinine or quinoline; and salts of basic amino acids such as lysine and arginine.

Examples of inorganic bases include, without limitation, sodium hydroxide, potassium hydroxide, potassium carbonate, sodium carbonate, sodium bicarbonate, potassium bicarbonate, calcium hydroxide, calcium carbonate, and the like.

Compounds of Structural Formula I, which are substituted with a basic group, may exist as salts with

pharmaceutically acceptable acids. The present invention includes such salts. Examples of such salts include hydrochlorides, hydrobromides, sulfates, methanesulfonates, nitrates, maleates, acetates, citrates, fumarates, tartrates  
5 [e.g. (+)-tartrates, (-)-tartrates or mixtures thereof including racemic mixtures], succinates, benzoates and salts with amino acids such as glutamic acid.

These salts may be prepared by methods known to those skilled in the art.

10 Certain compounds of Structural Formula I and their salts may also exist in the form of solvates, for example hydrates, and the present invention includes each solvate and mixtures thereof.

In addition, it is generally not desirable to formulate  
15 pharmaceuticals containing substantial amounts of organic solvent (e.g., ethyl acetate) due to potential solvent toxicity to the recipient thereof and changes in potency of the pharmaceutical as a function of the solvent. In addition, from a manufacturing perspective, it is also  
20 generally less desirable to prepare non-crystalline materials whenever said preparation involves a collection of the final product via filtration. Such filtrations are often more difficult to perform when the material collected is non-crystalline. Moreover, it is also generally less  
25 desirable, from a manufacturing perspective, to formulate pharmaceuticals containing substantial amounts of water (hydrates) because the level of hydration will typically be some function of the relative humidity at which the pharmaceutical is produced and stored. In other words,  
30 potency variability is typically more problematic with a hydrate relative to its anhydrous form. The present invention provides a desired crystalline form.

Prodrugs are compounds of the present invention, which have chemically or metabolically cleavable groups and become by solvolysis or under physiological conditions the compounds of the invention which are pharmaceutically active  
5 in vivo. Prodrugs include acid derivatives well known to practitioners of the art, such as, for example, esters prepared by reaction of the parent acidic compound with a suitable alcohol, or amides prepared by reaction of the parent acid compound with a suitable amine. Simple  
10 aliphatic or aromatic esters derived from acidic groups pendent on the compounds of this invention are preferred prodrugs. In some cases it is desirable to prepare double ester type prodrugs such as (acyloxy) alkyl esters or ((alkoxycarbonyl)oxy)alkyl esters. Particularly preferred  
15 esters as prodrugs are methyl, ethyl, propyl, isopropyl, n-butyl, isobutyl, tert-butyl, morpholinoethyl, and N,N-diethylglycolamido.

Methyl ester prodrugs may be prepared by reaction of the acid form of a compound of Formula I in a medium such as  
20 methanol with an acid or base esterification catalyst (e.g., NaOH, H<sub>2</sub>SO<sub>4</sub>). Ethyl ester prodrugs are prepared in similar fashion using ethanol in place of methanol.

Morpholinylethyl ester prodrugs may be prepared by reaction of the sodium salt of a compound of Structural  
25 Formula I (in a medium such as dimethylformamide) 4-(2-chloroethyl)morphine hydrochloride (available from Aldrich Chemical Co., Milwaukee, Wisconsin USA, Item No. C4,220-3).

The term, "active ingredient" means the compounds generically described by Structural Formula I as well as the  
30 salts, solvates, and prodrugs of such compounds.

The term "pharmaceutically acceptable" means that the carrier, diluent, excipients and salt must be compatible

with the other ingredients of the composition, and not deleterious to the recipient thereof. Pharmaceutical compositions of the present invention are prepared by procedures known in the art using well-known and readily available ingredients.

"Preventing" refers to reducing the likelihood that the recipient will incur or develop any of the pathological conditions described herein.

"Treating" refers to mediating a disease or condition and preventing, or mitigating, its further progression or ameliorate the symptoms associated with the disease or condition.

"Pharmaceutically-effective amount" means that amount of a compound, or of its salt, solvate, hydrate or prodrug thereof, that will elicit the biological or medical response of a tissue, system, or mammal. Such an amount can be administered prophylactically to a patient thought to be susceptible to development of a disease or condition. Such amount when administered prophylactically to a patient can also be effective to prevent or lessen the severity of the PPAR mediated condition. Such an amount is intended to include an amount which is sufficient to modulate a PPAR receptor or to prevent or mediate a disease or condition. Conditions prevented or treated by PPAR receptors include, but are not limited to, diabetes mellitus, cardiovascular disease, Syndrome X, obesity and gastrointestinal disease.

A "mammal" is an individual animal that is a member of the taxonomic class Mammalia. The class Mammalia includes humans, monkeys, chimpanzees, gorillas, cattle, swine, horses, sheep, dogs, cats, mice, and rats.

Administration to a human is most preferred. The compounds and compositions of the present invention can be

useful for the treatment and/or prophylaxis of cardiovascular disease, for raising serum HDL cholesterol levels, for lowering serum triglyceride levels and for lower serum LDL cholesterol levels. Elevated triglyceride and LDL  
5 levels, and low HDL levels, are considered risk factors for the development of heart disease, stroke, and circulatory system disorders and diseases.

The compounds and compositions of the present invention can also be useful for treating and/or preventing obesity.

10 Further, these compounds and compositions can be useful for the treatment and/or prophylaxis of non-insulin dependent diabetes mellitus (NIDDM) with reduced or no body weight gains by the patients. Furthermore, the compounds and compositions of the present invention can be useful to treat  
15 or prevent acute or transient disorders in insulin sensitivity, such as sometimes occur following surgery, trauma, myocardial infarction, and the like. The physician of ordinary skill can identify humans who will benefit from administration of the compounds and compositions of the  
20 present invention.

The present invention further provides a method for the treatment and/or prophylaxis of hyperglycemia in a human or non-human mammal which comprises administering an effective, non-toxic amount of a compound of the general formula (I),  
25 or a tautomeric form thereof and/or a pharmaceutically acceptable salt thereof and/or a pharmaceutically acceptable solvate thereof to a hyperglycemic human or non-human mammal in need thereof.

They are useful as therapeutic substances in preventing  
30 or treating Syndrome X, diabetes mellitus and related endocrine and cardiovascular disorders and diseases in human or non-human animals.



The invention also relates to the use of a compound of Formula I as described above, for the manufacture of a medicament for treating a PPAR receptor mediated condition.

A therapeutically effective amount of a compound of Structural Formula I can be used for the preparation of a medicament useful for treating Syndrome X, diabetes, treating obesity, lowering triglyceride levels, lowering serum LDL levels, raising the plasma level of high density lipoprotein, and for treating, preventing or reducing the risk of developing atherosclerosis, and for preventing or reducing the risk of having a first or subsequent atherosclerotic disease event in mammals, particularly in humans. In general, a therapeutically effective amount of a compound of the present invention typically reduces serum triglyceride levels of a patient by about 20% or more, and increases serum HDL levels in a patient. Preferably, HDL levels will be increased by about 30% or more. In addition, a therapeutically effective amount of a compound, used to prevent or treat NIDDM, typically reduces serum glucose levels, or more specifically HbA1c, of a patient by about 0.7% or more.

Advantageously, compositions containing the compound of Structural Formula I and/or the salts thereof may be provided in dosage unit form. Preferably each dosage unit contain from about 1 to about 500 mg of active ingredient. Although it will, of course, readily be understood that the amount of the compound or compounds of Structural Formula I actually to be administered will be determined by a physician, in the light of all the relevant circumstances.

When used herein Syndrome X includes pre-diabetic insulin resistance syndrome and the resulting complications thereof, insulin resistance, non-insulin dependent diabetes,

dyslipidemia, hyperglycemia obesity, coagulopathy, hypertension and other complications associated with diabetes. The methods and treatments mentioned herein include the above and encompass the treatment and/or prophylaxis of any one of or any combination of the following: pre-diabetic insulin resistance syndrome, the resulting complications thereof, insulin resistance, Type II or non-insulin dependent diabetes, dyslipidemia, hyperglycemia, obesity and the complications associated with diabetes including cardiovascular disease, especially atherosclerosis.

The compounds of the present invention can be useful for treating PPAR mediated conditions and for use as research tools. Certain compounds and conditions within the scope of this invention are preferred. The following conditions, invention embodiments, and compound characteristics listed in tabular form may be independently combined to produce a variety of preferred compounds and treatment conditions. The following list of embodiments of this invention is not intended to limit the scope of this invention in any way.

Some preferred characteristics of compounds of Formula I are:

- (a) R3 is methyl;
- (b) R3 is arylC<sub>0</sub>-C<sub>2</sub>alkoxy;
- (c) R4 is arylC<sub>0</sub>-C<sub>2</sub>alkoxy;
- (d) R4 is arylalkyl;
- (e) R3 and R4 are each C<sub>1</sub>-C<sub>6</sub> alkyl;
- (f) A is carboxyl;
- (g) W is O;
- (h) W is S;
- (i) X is -CH<sub>2</sub>CH<sub>2</sub>CH<sub>2</sub>-;

- (j) X is  $-\text{CH}_2\text{CH}_2\text{O}-$ ;
- (k) R9 is methyl;
- (l) R9 is benzyl;
- (m) R9 is heteroarylalkyl;
- 5 (n) R8 is hydrogen;
- (o) R10 is methyl
- (p) Y is  $\text{CH}_2$ ;
- (q) Y is O;
- (r) R21 is  $=\text{O}$ ;
- 10 (s) R21 is H;
- (t) R22 is H;
- (u) R1 is arylalkyl;
- (v) R1 is substituted arylalkyl;
- (w) R2 is methyl;
- 15 (x) R2 is H;
- (y) E is  $\text{C}(\text{R}_3)(\text{R}_4)\text{A}$ ;
- (z) A is tetrazole;
- (aa) A is acylsulfonamide;
- (bb) R21 is arylalkyl;
- 20 (cc) R1 arylalkyl is substituted by  $\text{CF}_3$ ;
- (dd) Aryl is a phenyl group;
- (ee) Heteroaryl group contains a N;
- (ff) "----" represents a double bond, as shown  
by Formula I herein;
- 25 (gg) A compound of Formula I that selectively  
modulates an alpha receptor;
- (hh) A compound of Formula I that is a PPAR  
coagonist that modulates an alpha  
receptor and a gamma receptor; and
- 30 (ii) A compound of Formula I for use in the  
treatment of cardiovascular disease.

(jj) A compound of Formula I for use in the treatment of Type II diabetes and/or Syndrome X.

5       The compositions are formulated and administered in the same general manner as detailed herein. The compounds of the instant invention may be used effectively alone or in combination with one or more additional active agents depending on the desired target therapy. Combination  
10   therapy includes administration of a single pharmaceutical dosage composition which contains a compound of Structural Formula I and one or more additional active agents, as well as administration of a compound of Structural Formula I and each active agent in its own separate pharmaceutical dosage  
15   formulation. For example, a compound of Structural Formula I or thereof and an insulin secretagogue such as biguanides, thiazolidinediones, sulfonylureas, insulin, or  $\alpha$ -glucosidase inhibitors can be administered to the patient together in a single oral dosage composition such as a tablet or capsule,  
20   or each agent administered in separate oral dosage formulations. Where separate dosage formulations are used, a compound of Structural Formula I and one or more additional active agents can be administered at essentially the same time, i.e., concurrently, or at separately  
25   staggered times, i.e., sequentially; combination therapy is understood to include all these regimens.

      An example of combination treatment or prevention of atherosclerosis may be wherein a compound of Structural Formula I or salts thereof is administered in combination  
30   with one or more of the following active agents:  
antihyperlipidemic agents; plasma HDL-raising agents;  
antihypercholesterolemic agents, fibrates, vitamins,

aspirin, and the like. As noted above, the compounds of Structural Formula I can be administered in combination with more than one additional active agent.

Another example of combination therapy can be seen in treating diabetes and related disorders wherein the compounds of Structural Formula I, salts thereof can be effectively used in combination with, for example, sulfonylureas, biguanides, thiazolidinediones,  $\alpha$ -glucosidase inhibitors, other insulin secretagogues, insulin as well as the active agents discussed above for treating atherosclerosis.

The compounds of the present invention, and the pharmaceutically acceptable salts, solvates and hydrates thereof, have valuable pharmacological properties and can be used in pharmaceutical compositions containing a therapeutically effective amount of a compound of the present invention, or pharmaceutically acceptable salts, esters or prodrugs thereof, in combination with one or more pharmaceutically acceptable excipients. Excipients are inert substances such as, without limitation carriers, diluents, fillers, flavoring agents, sweeteners, lubricants, solubilizers, suspending agents, wetting agents, binders, disintegrating agents, encapsulating material and other conventional adjuvants. Proper formulation is dependent upon the route of administration chosen. Pharmaceutical compositions typically contain from about 1 to about 99 weight percent of the active ingredient which is a compound of the present invention.

Preferably, the pharmaceutical formulation is in unit dosage form. A "unit dosage form" is a physically discrete unit containing a unit dose, suitable for administration in human subjects or other mammals. For example, a unit dosage

form can be a capsule or tablet, or a number of capsules or tablets. A "unit dose" is a predetermined quantity of the active compound of the present invention, calculated to produce the desired therapeutic effect, in association with one or more pharmaceutically-acceptable excipients. The quantity of active ingredient in a unit dose may be varied or adjusted from about 0.1 to about 1000 milligrams or more according to the particular treatment involved.

The dosage regimen utilizing the compounds of the present invention is selected by one of ordinary skill in the medical or veterinary arts, in view of a variety of factors, including, without limitation, the species, age, weight, sex, and medical condition of the recipient, the severity of the condition to be treated, the route of administration, the level of metabolic and excretory function of the recipient, the dosage form employed, the particular compound and salt thereof employed, and the like.

Preferably, the compounds of the present invention are administered in a single daily dose, or the total daily dose may be administered in divided doses, two, three, or more times per day. Where delivery is via transdermal forms, of course, administration is continuous.

Suitable routes of administration of pharmaceutical compositions of the present invention include, for example, oral, eyedrop, rectal, transmucosal, topical, or intestinal administration; parenteral delivery (bolus or infusion), including intramuscular, subcutaneous, intramedullary injections, as well as intrathecal, direct intraventricular, intravenous, intraperitoneal, intranasal, or intraocular injections. The compounds of the invention can also be administered in a targeted drug delivery system,

such as, for example, in a liposome coated with endothelial cell-specific antibody.

For oral administration, the compounds can be formulated readily by combining the active compounds with pharmaceutically acceptable carriers well known in the art. Such carriers enable the compounds of the invention to be formulated as tablets, pills, powders, sachets, granules, dragees, capsules, liquids, elixers, tinctures, gels, emulsions, syrups, slurries, suspensions and the like, for oral ingestion by a patient to be treated. Pharmaceutical preparations for oral use can be obtained by combining the active compound with a solid excipient, optionally grinding a resulting mixture, and processing the mixture of granules, after adding suitable auxiliaries, if desired, to obtain tablets or dragee cores.

For oral administration in the form of a tablet or capsule, the active ingredient may be combined with an oral, non-toxic, pharmaceutically-acceptable carrier, such as, without limitation, lactose, starch, sucrose, glucose, methyl cellulose, calcium carbonate, calcium phosphate, calcium sulfate, sodium carbonate, mannitol, sorbitol, and the like; together with, optionally, disintegrating agents, such as, without limitation, cross-linked polyvinyl pyrrolidone, maize, starch, methyl cellulose, agar, bentonite, xanthan gum, alginic acid, or a salt thereof such as sodium alginate, and the like; and, optionally, binding agents, for example, without limitation, gelatin, acacia, natural sugars, beta-lactose, corn sweeteners, natural and synthetic gums, acacia, tragacanth, sodium alginate, carboxymethyl-cellulose, polyethylene glycol, waxes, and the like; and, optionally, lubricating agents, for example, without limitation, magnesium stearate, sodium stearate,

stearic acid, sodium oleate, sodium benzoate, sodium acetate, sodium chloride, talc, and the like. When a dosage unit form is a capsule, it may contain, in addition to materials of the above type, a liquid carrier such as a fatty oil.

Solid form formulations include powders, tablets and capsules. A solid carrier can be one or more substance which may also act as flavoring agents, lubricants, solubilisers, suspending agents, binders, tablet disintegrating agents and encapsulating material.

In powders, the carrier is a finely divided solid which is in admixture with the finely divided active ingredient. In tablets, the active ingredient is mixed with a carrier having the necessary binding properties in suitable proportions and compacted in the shape and size desired.

Various other materials may be present as coatings or to modify the physical form of the dosage unit. For instance, tablets may be coated with shellac, sugar or both. A syrup or elixir may contain, in addition to the active ingredient, sucrose as a sweetening agent, methyl and propylparabens as preservatives, a dye and a flavoring such as cherry or orange flavor.

Sterile liquid formulations include suspensions, emulsions, syrups, and elixirs. The active ingredient can be dissolved or suspended in a pharmaceutically acceptable carrier, such as sterile water, sterile organic solvent, or a mixture of both sterile water and sterile organic solvent.

The active ingredient can also be dissolved in a suitable organic solvent, for example, aqueous propylene glycol. Other compositions can be made by dispersing the finely divided active ingredient in aqueous starch or sodium carboxymethyl cellulose solution or in a suitable oil.



Dragee cores are provided with suitable coatings. For this purpose, concentrated sugar solutions may be used, which may optionally contain gum arabic, talc, polyvinyl pyrrolidone, carbopol gel, polyethylene glycol, and/or  
5 titanium dioxide, lacquer solutions, and suitable organic solvents or solvent mixtures. Dyestuffs or pigments may be added to the tablets or dragee coatings for identification or to characterize different combinations of active compound doses.

10 Pharmaceutical preparations which can be used orally include push-fit capsules made of gelatin, as well as soft, sealed capsules made of gelatin and a plasticizer, such as glycerol or sorbitol. The push-fit capsules can contain the active ingredients in admixture with filler such as lactose,  
15 binders such as starches, and/or lubricants such as talc or magnesium stearate and, optionally, stabilizers. In soft capsules, the active compounds may be dissolved or suspended in suitable liquids, such as fatty oils, liquid paraffin, or liquid polyethylene glycols. In addition, stabilizers may  
20 be added.

All formulations for oral administration should be in dosages suitable for such administration. Particularly suitable compositions for oral administration are unit dosage forms such as tablets and capsules.

25 For parental administration the compounds of the present invention, or salts thereof, can be combined with sterile aqueous or organic media to form injectable solutions or suspensions. Formulations for injection may be presented in unit dosage form, such as in ampoules or in  
30 multi-dose containers, with an added preservative. The compositions may take such forms as suspensions, solutions or emulsions in oily or aqueous vehicles, and may contain

formulatory agents such as suspending, stabilizing and/or dispersing agents. The pharmaceutical forms suitable for injectable use include sterile aqueous solutions or dispersions and sterile powders for the extemporaneous preparation of sterile injectable solutions or dispersions. In all cases, the form must be sterile and must be fluid to the extent that each syringability exists. It must be stable under the conditions of manufacture and storage and must be preserved against any contamination. The carrier can be solvent or dispersion medium containing, for example, water, preferably in physiologically compatible buffers such as Hanks' solution, Ringer's solution, or physiological saline buffer, ethanol, polyol (e.g. glycerol, propylene glycol and liquid polyethylene glycol), propylene glycol and liquid polyethylene glycol), suitable mixtures thereof, and vegetable oils. Under ordinary conditions of storage and use, these preparations contain a preservative to prevent the growth of microorganisms.

The injectable solutions prepared in this manner can then be administered intravenously, intraperitoneally, subcutaneously, or intramuscularly, with intramuscular administration being preferred in humans.

For transmucosal administration, penetrants appropriate to the barrier to be permeated are used in the formulation. Such penetrants are generally known in the art. The active compounds can also be administered intranasally as, for example, liquid drops or spray.

For buccal administration, the compositions may take the form of tablets or lozenges formulated in a conventional manner.

For administration by inhalation, the compounds for use according to the present invention are conveniently

delivered in the form of a dry powder inhaler, or an aerosol spray presentation from pressurized packs or a nebuliser, with the use of a suitable propellant, e.g., dichlorodifluoromethane, trichlorofluoromethane, 5 dichlorotetrafluoroethane, carbon dioxide or other suitable gas. In the case of pressurized aerosol the dosage unit may be determined by providing a valve to deliver a metered amount. Capsules and cartridges of gelatin for use in an inhaler or insufflator may be formulated containing a powder 10 mix of the compound and a suitable powder base such as lactose or starch.

Pharmaceutical compositions of the present invention can be manufactured in a manner that is itself known, e.g., by means of conventional mixing, dissolving, granulating, 15 dragee-making, levigating, emulsifying, encapsulating, entrapping or lyophilizing processes.

In making the compositions of the present invention, the active ingredient will usually be admixed with a carrier, or diluted by a carrier, or enclosed within a 20 carrier which may be in the form of a capsule, sachet, paper or other container. When the carrier serves as a diluent, it may be a solid, lyophilized solid or paste, semi-solid, or liquid material which acts as a vehicle, or can be in the form of tablets, pills, powders, lozenges, 25 elixirs, suspensions, emulsions, solutions, syrups, aerosols (as a solid or in a liquid medium), or ointment, containing, for example, up to 10% by weight of the active compound. The compounds of the present invention are preferably formulated prior to administration.

30 The following pharmaceutical formulations 1 through 8 are illustrative only and are not intended to limit the scope of the invention in any way. "Active Ingredient",

refers to a compound according to Structural Formula I or salts thereof.

#### Formulation 1

- 5 Hard gelatin capsules are prepared using the following ingredients:

	Quantity (mg/capsule)
Active Ingredient	250
Starch, dried	200
Magnesium stearate	<u>10</u>
Total	460 mg

#### Formulation 2

- 10 A tablet is prepared using the ingredients below:

	Quantity (mg/tablet)
Active Ingredient	250
Cellulose, microcrystalline	400
Silicon dioxide, fumed	10
Stearic acid	<u>5</u>
Total	665 mg

The components are blended and compressed to form tablets each weighing 665 mg.

15

#### Formulation 3

An aerosol solution is prepared containing the following components:



are dried at 50°C and passed through a No. 18 mesh U.S. sieve. The sodium carboxymethyl starch, magnesium stearate and talc, previously passed through a No. 60 mesh U.S. sieve, are then added to the granules which, after mixing,  
5 are compressed on a tablet machine to yield tablets each weighing 150 mg.

#### Formulation 5

Capsules, each containing 80 mg of Active Ingredient, are  
10 made as follows:

Active Ingredient	80 mg
Starch	59 mg
Microcrystalline cellulose	59 mg
Magnesium stearate	<u>2 mg</u>
Total	200 mg


The Active Ingredient, cellulose, starch, and magnesium stearate are blended, passed through a No. 45 mesh U.S. sieve, and filled into hard gelatin capsules in 200 mg  
15 quantities.

In yet another embodiment of the compounds of the present invention, the compound is radiolabelled, such as with carbon-14, or tritiated. Said radiolabelled or  
20 tritiated compounds are useful as reference standards for in vitro assays to identify new PPAR $\alpha$  and or PPAR $\delta$  agonists.

#### SYNTHESIS

Compounds of the present invention have been formed as  
25 specifically described in the examples. Further, many compounds were prepared as more generally as shown in the


following schematic. Alternative synthesis methods may also be effective and known to the skilled artisan.



5


The process as shown by the following scheme may be used to make certain compounds of this invention:

## Hydantoin and Imidazolone Synthesis - I







## Hydantoin and Imidazolone Synthesis - II



## Hydantoin and Imidazolone Synthesis - III



## Hydantoin and Imidazolone Synthesis - IV




The reaction of intermediate A with an excess of trifluoroacetic acid at ambient temperature in methylene chloride overnight produces the fibrin acid (see Scheme 5 immediately above). The fibrin acid was washed until only traces of trifluoroacetic acid were present in the crude

product, and then the fibrates acid is used to prepare compounds claimed here.

Scheme 1


5 **General Example A:** 2-Ethoxy-2-methyl-3-(4-{2-[3-[methyl-2-




oxo-1-(4-trifluoromethyl-benzyl)-imidazolidin-4-yl]-ethoxy)-phenyl)propionic ethyl ester

To the solution of the  $\alpha$ -hydroxy ester (0.04 g, (0.08 mmol) in DMF, is added silver oxide (0.28 g, 1.2 mmol) and ethyl iodide (0.07 mL, 1.0 mmol). The mixture is stirred at room temperature for 16 h, and then stirred at 50 °C for another 16 h. Filtered, the filtrate is washed with water and extracted with EtOAc. The organic layer is evaporated, dried (MgSO<sub>4</sub>), concentrated and chromatographed (silica gel; hexane/EtOAc, 1:1 to 0:1). An oil is isolated as the ethyl ester (25 mg, 59%). MS (ESI) m/z 537 (M+H)<sup>+</sup>. The ester is then hydrolyzed with 5.0N NaOH in methanol at 60 °C for 4 h. Acidified with conc.HCl, extracted with EtOAc, the organic layer is dried (MgSO<sub>4</sub>) and concentrated to afford the title acid. MS (ESI) m/z 509 (M+H)<sup>+</sup>. Other members of this family could be prepared using the same method and the appropriate alkyl iodide.

Scheme 2




5

**General example B:**

To a mixture of 54 mg (0.13 mmol) of 2-Methyl-3-{4-[2-  
 10 (3-methyl-2-oxo-imidazolidin-4-yl)-ethoxy]-phenyl}-2-  
 phenoxy-propionic acid ethyl ester in 2 mL of dry DMF at 0  
 °C under an atmosphere of nitrogen, 10 mg of NaH (0.253  
 mmol) is added. The resulting solution is allowed to stand  
 at r.t. for 20 min. Then 0.26 mmol of the appropriate aryl  
 15 halide is added and resulting mixture is allowed to stand at

r.t. for 3h. The reaction mixture is diluted with ethyl acetate and 1N HCl. The organic layer is then washed with 1N HCl (2 x 10 mL), brine (2 x 10 mL), dried over Na<sub>2</sub>SO<sub>4</sub>, filtered and concentrated under vacuum. The resulting  
5 esters were purified by flash chromatography, and could be used as starting materials in subsequent coupling reactions, or hydrolyzed to final compounds: To hydrolyze, the ester is dissolved in 2 mL of MeOH and 0.3 mL of 5N NaOH, and the mixture is heated to 50 °C for 2 h. The organic solvent is  
10 then removed under vacuum and the residue is dissolved in CH<sub>2</sub>Cl<sub>2</sub> and 1N HCl. Aqueous layer is washed with CH<sub>2</sub>Cl<sub>2</sub> (2 x 10 mL). Combined organic layer is dried over Na<sub>2</sub>SO<sub>4</sub>, filtered and concentrated under vacuum. The crude material is purified by MS/LC or flash chromatography to give the  
15 final products.

**General example C: Suzuki couplings**



To the aryl bromide (or chloride) (0.07 mmol) in 1,4 dioxane  
20 (2 mL) in a N<sub>2</sub> purged sealed tube, is added aryl boronic acid (0.12 mmol), Cesium fluoride (0.18 mmol) and tetrakis(triphenylphosphine) Pd (0) (0.007 mmol). The reaction mixture is stirred at 110 °C for 2 h. Cooled, diluted with EtOAc and quenched with water. The organic  
25 layer is dried (MgSO<sub>4</sub>), filtered and concentrated. The residue is then hydrolyzed with 5.0 N NaOH in methanol at 60


°C for 2h. Acidified, extracted with EtOAc, the crude acid is chromatographed (silica gel; EtOAc/MeOH, 10:0 to 10:1).

#### General example D: Buchwald couplings


##### 5 General procedure:

The phenyl bromide (0.04 g, 0.067 mmol), phenol (12 mg, 0.13 mmol), Cs<sub>2</sub>CO<sub>3</sub> (33 mg, 0.1 mmol), (CuOTf)<sub>2</sub>•PhH (2mg), ethyl acetate (0.0033 mmol, 5.0 mol %) and toluene were added to a sealed tube which is purged with nitrogen. The mixture is  
10 heated to 100 °C until the phenyl bromide is consumed as determined by LC-MS. Cooled, diluted with ethyl acetate and washed with H<sub>2</sub>O. The organic layer is dried (MgSO<sub>4</sub>), concentrated to afford the crude product. The crude ester is  
15 then hydrolyzed with 5.0N NaOH in MeOH at 60 °C for 2 h. Acidified, extracted with ethyl acetate, the organic layer is dried (MgSO<sub>4</sub>), concentrated and chromatographed (silica gel; EtOAc/MeOH, 10:0 to 10:1) to afford the pure acid.


Scheme 4



2-Isopropoxy-3-(4-{2-[1-(4-methoxy-benzyl)-3-methyl-2-oxo-imidazolidin-4-yl]-ethoxy}-phenyl)-propionic acid methyl ester



- 5 3-(4-Hydroxy-phenyl)-2-isopropoxy-propionic acid methyl ester (1.7g, 7.3 mmol), Toluene-4-sulfonic acid 2-[1-(4-methoxy-benzyl)-3-methyl-2-oxo-imidazolidin-4-yl]-ethyl ester (3.4 g, 8.0 mmol), and  $\text{Cs}_2\text{CO}_3$  (3.6g, 11 mmol) were combined in 10 mL of DMF and heated at 55 oC for 18 h.
- 10 After cooling, the mixture is diluted with 100 mL of water and extracted with 3x20 mL of ethyl acetate. The organics were dried, concentrated, and the residue is purified by flash chromatography using 7/3 ethyl acetate/hexanes to give 3.0 g (86 %) of the title compound as an oil. NMR
- 15 **General example E:**  
 3-(4-[2-(1-Substitutedbenzyl-3-methyl-2-oxo-imidazolidin-4-yl)-ethoxy]-phenyl)-2-isopropoxy-propionic acid methyl ester




- Prepared as in **General example B** using 2-Isopropoxy-3-{4-[2-(3-methyl-2-oxo-imidazolidin-4-yl)-ethoxy]-phenyl}-propionic acid methyl ester and the appropriate benzyl halide. The resulting esters were purified by flash chromatography, and could be used as starting materials in
- 20

subsequent coupling reactions, or hydrolized to final compounds: To hydrolize, the ester is dissolved in 2 mL of MeOH and 0.3 mL of 5N NaOH, and the mixture is heated to 50 °C for 2 h. The organic solvent is then removed under vacuum and the residue is dissolved in CH<sub>2</sub>Cl<sub>2</sub> and 1N HCl. The aqueous layer is washed with CH<sub>2</sub>Cl<sub>2</sub> (2 x 10 mL). The organics were combined, dried over Na<sub>2</sub>SO<sub>4</sub>, filtered and concentrated under vacuum. The crude material is purified by MS/LC or flash chromatography to give the final products.

**General example F:**

**3-{4-[2-(1-Substituted-biphenyl-4-ylmethyl-3-methyl-2-oxo-imidazolidin-4-yl)-ethoxy]-phenyl}-2-isopropoxy-propionic acid**



Prepared as in **General example C** by coupling 3-(4-{2-[1-(4-bromo-benzyl)-3-methyl-2-oxo-imidazolidin-4-yl]-ethoxy}-phenyl)-2-isopropoxy-propionic acid methyl ester and the appropriate boronic acid, followed by hydrolysis to the carboxylic acid.

The Examples provided herein are illustrative of the invention claimed herein and are not intended to limit the scope of the claimed invention in any way.



## EXEMPLIFICATION

Instrumental Analysis


Infrared spectra were recorded on a Perkin Elmer 781 spectrometer. <sup>1</sup>H NMR spectra were recorded on a Varian 400  
5 MHz spectrometer at ambient temperature. Data are reported as follows: chemical shift in ppm from internal standard tetramethylsilane on the  $\delta$  scale, multiplicity (b = broad, s = singlet, d = doublet, t = triplet, q = quartet, qn = quintet and m = multiplet), integration, coupling constant  
10 (Hz) and assignment. <sup>13</sup>C NMR were recorded on a Varian 400 MHz spectrometer at ambient temperature. Chemical shifts are reported in ppm from tetramethylsilane on the  $\delta$  scale, with the solvent resonance employed as the internal standard (CDCl<sub>3</sub> at 77.0 ppm and DMSO-d<sub>6</sub> at 39.5 ppm). Combustion  
15 analyses were performed by Eli Lilly and Company Microanalytical Laboratory. High-resolution mass spectra were obtained on VG ZAB 3F or VG 70 SE spectrometers. Analytical thin layer chromatography was performed on EM Reagent 0.25 mm silica gel 60-F plates. Visualization was  
20 accomplished with UV light.

PreparationsPreparation 1

Preparation of toluene-4-sulfonic acid 2-[1-(4-methoxy-  
25 benzyl)-3-methyl-2-oxo-imidazolidin-4-yl]-ethyl ester

Step A

(2,5-Dioxo-imidazolidin-4-yl)-acetic acid methyl ester




30 5-Hydantoin acetic acid (20.57 g, 0.130 mol) in MeOH (210 mL) was treated with conc. H<sub>2</sub>SO<sub>4</sub> (7 mL) and heated to reflux


under N<sub>2</sub> for 2.5 h. The resultant clear solution was cooled and the solvent removed *in vacuo* to give an oil which was diluted with water (65 mL) and extracted 4 times with EtOAc. The organic layer was dried (MgSO<sub>4</sub>) and the solvent removed  
5 *in vacuo* to afford 19.81 g (88%) of (2,5-dioxo-imidazolidin-4-yl)-acetic acid methyl ester. MS (ES<sup>+</sup>) Calc'd for C<sub>6</sub>H<sub>9</sub>N<sub>2</sub>O<sub>4</sub> (M +1) 173. Found *m/z* 173 (100%). <sup>1</sup>H NMR.

Step B


10 **[1-(4-Methoxy-benzyl)-2,5-dioxo-imidazolidin-4-yl]-acetic acid methyl ester**



A solution of (2,5-dioxo-imidazolidin-4-yl)-acetic acid methyl ester (26.6 g, 0.155 mol) in DMF (500 mL) was treated  
15 with 4-methoxybenzyl chloride (26.6 g, 0.170 mol), MgSO<sub>4</sub> (18.6 g, 0.154 mol) and then 325 mesh K<sub>2</sub>CO<sub>3</sub> (42.71 g, 0.309 mol) at 0 °C. The resultant mixture was warmed to room temperature under N<sub>2</sub> and then heated at 45 °C for 4 h. The reaction mixture was filtered, and then aqueous 1N HCl (200  
20 mL) was added to the filtrate. The filtrate was extracted with EtOAc and the organic layer dried (MgSO<sub>4</sub>). The solvent was removed *in vacuo* to give 11.43 g crude product which was purified by flash chromatography using a gradient of 3:1 to 1:1 hexanes:acetone to give 9.81 g (22%) [1-(4-methoxy-  
25 benzyl)-2,5-dioxo-imidazolidin-4-yl]-acetic acid methyl ester. MS (ES<sup>+</sup>) Calc'd for C<sub>14</sub>H<sub>17</sub>N<sub>2</sub>O<sub>5</sub> (M +1) 293. Found *m/z* 293 (100%). <sup>1</sup>H NMR.

Step C**[1-(4-methoxy-benzyl)-3-methyl-2,5-dioxo-imidazolidin-4-yl]-acetic acid methyl ester**

- 5 A 0 °C solution of compound [1-(4-methoxy-benzyl)-2,5-dioxo-imidazolidin-4-yl]-acetic acid methyl ester (9.84 g, 33.7 mmol) in DMF (30 mL) was treated with sodium hydride (60% dispersion, 1.37 g, 34.3 mmol) and warmed to room temperature and stirred under N<sub>2</sub> for 20 minutes. The
- 10 resultant mixture was cooled to 0 °C and then treated with methyl iodide (6.16 g, 43.4 mmol) and then warmed to room temperature and stirred for 16 h. The reaction was quenched with aqueous 1 N HCl (60 mL) and then worked up extractively with EtOAc and water. The organic layer was dried (MgSO<sub>4</sub>)
- 15 and the solvent removed *in vacuo* to give crude product that was purified by flash chromatography using 3:1 hexanes:acetone to afford 7.46 g (85%) [1-(4-methoxy-benzyl)-3-methyl-2,5-dioxo-imidazolidin-4-yl]-acetic acid methyl ester. MS (ES<sup>+</sup>) Calc'd for C<sub>15</sub>H<sub>19</sub>N<sub>2</sub>O<sub>5</sub> (M +1) 307.
- 20 Found *m/z* 307 (100%). <sup>1</sup>H NMR.


Step D**4-(2-Hydroxy-ethyl)-1-(4-methoxy-benzyl)-3-methyl-imidazolidin-2-one**

- 25 A solution of [1-(4-methoxy-benzyl)-3-methyl-2,5-dioxo-imidazolidin-4-yl]-acetic acid methyl ester (7.45 g, 24.3 mmol) in methanol (100 mL) was treated with aqueous 5 N NaOH (49 mL) and heated to reflux 1 h. The reaction mixture was

cooled, the solvent removed *in vacuo*. The resultant residue was acidified with aqueous 1 N HCl (300 mL) and extracted with EtOAc and water. The organic layer was dried (MgSO<sub>4</sub>) and the solvent removed *in vacuo* to afford 7.73 g (100%) acid that was utilized without purification. A solution of crude acid (7.73 g, assume 24.3 mmol) in THF (100 mL) was treated dropwise with 1 M solution of borane-THF complex in THF (146 mL, 0.145 mol) and then stirred at room temperature under N<sub>2</sub> for 16 h. The reaction was quenched with methanol (100 mL) and stirred at room temperature for 1 h. The solvent was removed *in vacuo* to give crude product that was purified by flash chromatography using 2:1 hexanes:acetone to afford 4.76 g (74%) 4-(2-hydroxy-ethyl)-1-(4-methoxy-benzyl)-3-methyl-imidazolidin-2-one. MS (ES<sup>+</sup>) Calc'd for C<sub>14</sub>H<sub>21</sub>N<sub>2</sub>O<sub>3</sub> (M + 1) 265. Found *m/z* 265 (100%). <sup>1</sup>H NMR.

#### Step E

**Toluene-4-sulfonic acid 2-[1-(4-methoxy-benzyl)-3-methyl-2-oxo-imidazolidin-4-yl]-ethyl ester**



A solution of 4-(2-hydroxy-ethyl)-1-(4-methoxy-benzyl)-3-methyl-imidazolidin-2-one (4.75 g, 18.0 mmol), pyridine (4.98 g, 62.9 mmol) and 4-dimethyl amino pyridine (0.66 g, 5.40 mmol) in CH<sub>2</sub>Cl<sub>2</sub> (200 mL) was treated with *p*-toluenesulfonic anhydride (9.38 g, 28.7 mmol) and the reaction stirred at room temperature for under N<sub>2</sub> for 1.5 h. The reaction mixture was washed with aqueous 1 N HCl (140 mL), the organic layer was dried (MgSO<sub>4</sub>), and the solvent removed *in vacuo* to afford crude product that was purified


by flash chromatography using 3:1 hexanes:acetone to afford 6.93 g (92%) toluene-4-sulfonic acid 2-[1-(4-methoxy-benzyl)-3-methyl-2-oxo-imidazolidin-4-yl]-ethyl ester. MS (ES<sup>+</sup>) Calc'd for C<sub>21</sub>H<sub>27</sub>N<sub>2</sub>O<sub>5</sub>S (M + 1) 419. Found m/z 419 (100%). <sup>1</sup>H NMR.

### Exemplified Compounds

10

#### **Example 1**


**2-(4-{2-[1-(4-tert-Butyl-benzyl)-3-methyl-2-oxo-imidazolidin-4-yl]-ethoxy}-2-methyl-phenoxy)-2-methyl-propionic acid**



15

#### Step A

**2-(4-{2-[1-(4-Methoxy-benzyl)-3-methyl-2-oxo-imidazolidin-4-yl]-ethoxy}-2-methyl-phenoxy)-2-methyl-propionic acid ethyl ester**




20

To a mixture of toluene-4-sulfonic acid 2-[1-(4-methoxy-benzyl)-3-methyl-2-oxo-imidazolidin-4-yl]-ethyl ester

(0.100g, 0.239 mmole) and 2-(4-hydroxy-2-methyl-phenoxy)-2-methyl-propionic acid ethyl ester (0.063g, 0.262 mmole) in anhydrous DMF (2.0 mL), is added cesium carbonate (0.093g, 0.287 mmole). After heating at 65°C for about 16 hours, the  
5 reaction mixture is partitioned between ethyl acetate (5 ml) and 1.0 N HCl aq. (5 ml), the aqueous layer is extracted with more ethyl acetate (2X5 ml). The combined organic layer is washed with brine (3X5 ml). The organic layer is dried over Na<sub>2</sub>SO<sub>4</sub>, filtered, and concentrated *in vacuo*.  
10 Chromatography using silica gel (0-40% acetone / hexane) produces a colorless oil (0.107g, 91%). Mass [EI+] 485 (M+H)<sup>+</sup>.

#### Step B

15 2-Methyl-2-{2-methyl-4-[2-(3-methyl-2-oxo-imidazolidin-4-yl)-ethoxy]-phenoxy}-propionic acid ethyl ester



2-(4-{2-[1-(4-Methoxy-benzyl)-3-methyl-2-oxo-imidazolidin-4-yl]-ethoxy}-2-methyl-phenoxy)-2-methyl-propionic acid ethyl  
20 ester (0.105g, 0.217 mmole) is treated with triethylsilane (0.064g, 0.743 mmole) in TFA (8.0 mL) at room temperature for about 6 hours. After evaporating the solvents, the residue is purified using silica gel (0-50% acetone /  
25 hexane). Colorless oil (0.041g, 51%). Mass [EI+] 365 (M+H)<sup>+</sup>, 729 (2M+H)<sup>+</sup>.


Step C

2-(4-{2-[1-(4-tert-Butyl-benzyl)-3-methyl-2-oxo-  
imidazolidin-4-yl]-ethoxy}-2-methyl-phenoxy)-2-methyl-  
propionic acid

To a solution of 2-methyl-2-{2-methyl-4-[2-(3-methyl-2-oxo-  
imidazolidin-4-yl)-ethoxy]-phenoxy}-propionic acid ethyl  
ester (0.040g, 0.110 mmole) in DMF (2.0 mL), is added sodium  
hydride (60% in mineral oil, 0.0066g, 0.165 mmole) in one  
portion. The mixture is stirred for 15 minutes at room  
temperature, then was added 4-tert-butyl-benzyl bromide  
(0.030 mL, 0.165 mmole). After stirring at room temperature  
for 4 hours, the reaction mixture is partitioned between  
ethyl acetate (5 mL) and saturated  $\text{NH}_4\text{Cl}$  aq. (5 mL), the  
aqueous layer is extracted with more ethyl acetate (2X5 mL).  
The combined organic layer is washed with brine (3X5 mL),  
dried over  $\text{Na}_2\text{SO}_4$ , filtered, and concentrated in vacuo. The  
crude product was purified by silica gel (0-40% Acetone /  
Hexane). The ethyl ester obtained above is treated with a  
mixture of MeOH (2 mL)/5.0N NaOH (1 mL) at room temperature  
overnight, and then concentrated. The resulting residue is  
diluted with water (2 mL), cooled down to 0 °C, acidified to  
pH = 2 by adding concentrated HCl dropwise. The aqueous  
suspension is loaded on a Chem elut 1005 tube, eluted with  
DCM (50 mL). Evaporation of methylene chloride gives the  
titled compound as an colorless oil (0.022 g, 42%). Mass  
[EI+] 483 (M+H)<sup>+</sup>, [EI-] 481 (M-H)<sup>-</sup>.

**Example 2**

**2-(4-{2-[1-(4-tert-Butyl-benzyl)-3-methyl-2-oxo-imidazolidin-4-yl]-ethoxy}-3-propyl-phenoxy)-2-methyl-propionic acid**



5

Step A

2-(4-{2-[1-(4-Methoxy-benzyl)-3-methyl-2-oxo-imidazolidin-4-yl]-ethoxy}-3-propyl-phenoxy)-2-methyl-propionic acid ethyl ester

10

The titled compound is prepared, according to the procedure of Example 1, Step A, using toluene-4-sulfonic acid 2-[1-(4-methoxy-benzyl)-3-methyl-2-oxo-imidazolidin-4-yl]-ethyl ester (0.200g, 0.478 mmole) and 2-(4-hydroxy-3-propyl-phenoxy)-2-methyl-propionic acid ethyl ester (0.140g, 0.526 mmole) to produce a colorless oil (0.122g, 50%). Mass [EI+] 513 (M+H)<sup>+</sup>.

15

Step B

2-Methyl-2-(4-[2-(3-methyl-2-oxo-imidazolidin-4-yl)-ethoxy]-3-propyl-phenoxy)-propionic acid ethyl ester

20

The titled compound is prepared, according to the procedure of Example 1, Step B, using 2-(4-{2-[1-(4-methoxy-benzyl)-3-methyl-2-oxo-imidazolidin-4-yl]-ethoxy}-3-propyl-phenoxy)-2-methyl-propionic acid ethyl ester (0.120g, 0.234 mmole) to

25



produce a colorless oil (0.054g, 57%). Mass [EI+] 393 (M+H)<sup>+</sup>, 785 (2M+H)<sup>+</sup>.

### Step C


5        2-(4-{2-[1-(4-tert-Butyl-benzyl)-3-methyl-2-oxo-imidazolidin-4-yl]-ethoxy}-3-propyl-phenoxy)-2-methyl-propionic acid

The titled compound is prepared, according to the procedure  
10 of Example 1, Step B, using 2-methyl-2-{4-[2-(3-methyl-2-oxo-imidazolidin-4-yl)-ethoxy]-3-propyl-phenoxy}-propionic acid ethyl ester (0.054g, 0.138 mmole) to produce a colorless oil (0.051g, 72%). Mass [EI+] 511 (M+H)<sup>+</sup>, [EI-] 509 (M-H)<sup>-</sup>.

15

### Example 3


2-(4-{3-[1-(4-tert-Butyl-benzyl)-3-methyl-2-oxo-imidazolidin-4-yl]-propyl}-2-methyl-phenoxy)-2-methyl-propionic acid



20

### Step A


1-(4-tert-Butyl-benzyl)-4-[3-(3-iodo-4-methoxy-phenyl)-propyl]-3-methyl-imidazolidin-2-one



To a solution of 1-(4-tert-Butyl-benzyl)-4-[3-(4-methoxy-phenyl)-propyl]-3-methyl-imidazolidin-2-one (0.630g, 1.60 mmole) in ethanol (15 mL), is added iodine (0.810g, 3.20 mmole) followed by silver sulfate (0.998g, 3.20 mmole). The reaction mixture is stirred at room temperature overnight. The precipitate is removed through filtration, and the solution is concentrated *in vacuo*. The residue is purified by column chromatography (silica gel, gradient elution 0-20% acetone in hexane) to provide a white foamy solid (0.584g, 70%). Mass [EI+] 521 (M+H)<sup>+</sup>.

#### Step B

1-(4-tert-Butyl-benzyl)-4-[3-(4-methoxy-3-methyl-phenyl)-propyl]-3-methyl-imidazolidin-2-one




1-(4-tert-Butyl-benzyl)-4-[3-(3-iodo-4-methoxy-phenyl)-propyl]-3-methyl-imidazolidin-2-one (0.300g, 0.580 mmole), methyl boronic acid (0.069g, 1.16 mmole) and cesium carbonate (0.264g, 1.74 mmole) are mixed in dioxane (6.0 mL). After bubbling with nitrogen for 15 minutes, 1,1'-bis(diphenylphosphino)ferrocene palladium (II) chloride (0.060g, 0.015 mmole) is added. The reaction is heated at 80°C for 4 hours. The solvent is removed on rota-vapor, and

the crude product is purified by column chromatography (silica gel, gradient elution 0-20% acetone in hexane) to provide a yellow oil (0.153g, 65%). Mass [EI+] 409 (M+H)<sup>+</sup>, 817 (M+H)<sup>+</sup>.

5

Step C

1-(4-tert-Butyl-benzyl)-4-[3-(4-hydroxy-3-methyl-phenyl)-propyl]-3-methyl-imidazolidin-2-one




To a solution of 1-(4-tert-Butyl-benzyl)-4-[3-(4-methoxy-3-methyl-phenyl)-propyl]-3-methyl-imidazolidin-2-one (0.250g, 0.612 mmole) in DCM (2.0 mL) at -78 °C, is added dropwise the solution of BBr<sub>3</sub> (0.230 mL, 2.44 mmole) in methylene chloride (2.0 mL). The reaction is kept for 30 minutes at about -78 °C, then warmed up to 0 °C and stirred for an hour. It is then quenched by 1:1 MeOH/DCM (20 mL), stirred for another hour at 0 °C. The reaction mixture is partitioned between DCM (25 mL) and water (25 mL), the organic layer is separated, washed by brine (3x25 mL), dried over Na<sub>2</sub>SO<sub>4</sub>, filtered and concentrated *in vacuo*. The crude product is purified by column chromatography (silica gel, gradient elution 0-20% acetone in hexane) to provide a yellow oil (0.126g, 53%). Mass [EI+] 395 (M+H)<sup>+</sup>, 789 (M+H)<sup>+</sup>, [EI-] 393 (M-H)<sup>-</sup>.

25

Step D

2-(4-{3-[1-(4-tert-Butyl-benzyl)-3-methyl-2-oxo-imidazolidin-4-yl]-propyl}-2-methyl-phenoxy)-2-methyl-propionic acid ethyl ester



1- (4-tert-Butyl-benzyl) -4- [3- (4-hydroxy-3-methyl-phenyl) -  
propyl]-3-methyl-imidazolidin-2-one (0.060g, 0.152 mmol) is  
dissolved in DMF (1.5 mL), to it is added ethyl 2-  
bromoisobutyrate (0.141g, 0.760 mmol) followed by potassium  
5 carbonate (0.105g, 0.760 mmol). After heating at 50 °C  
overnight, the reaction mixture is diluted with ethyl  
acetate (2 mL), washed with water (2 mL), the separated  
organic layer is passed through a chem elut tube, and the  
tube is washed with more DCM (50 mL). Evaporation of  
10 solvent followed by chromatography on silica gel, (gradient  
elution 0-40% ethyl acetate in hexane) to provide an oil  
(0.068g, 88%). Mass [EI+] 510 (M+H)<sup>+</sup>.

#### Step E


15 2- (4- {3- [1- (4-tert-Butyl-benzyl) -3-methyl-2-oxo-  
imidazolidin-4-yl]-propyl} -2-methyl-phenoxy) -2-methyl-  
propionic acid

2- (4- {3- [1- (4-tert-Butyl-benzyl) -3-methyl-2-oxo-  
20 imidazolidin-4-yl]-propyl} -2-methyl-phenoxy) -2-methyl-  
propionic acid ethyl ester (0.065g, 0.128 mmole) is treated  
with 3:1 MeOH/5.0N NaOH (4 mL) at room temperature  
overnight, and then concentrated. The resulting residue is  
diluted with water (2 mL), cooled down to 0 °C, acidified to  
25 pH = 2 by adding concentrated HCl dropwise. The aqueous  
suspension is loaded on a Chem elut tube and eluted with DCM  
(50 mL). Evaporation of methylene chloride gives the titled

compound as an oil (0.060g, 98%). Mass [EI+] 481 (M+H)<sup>+</sup>,  
[EI-] 479 (M-H)<sup>-</sup>.


#### Example 4

5. 2-(2-Butyl-4-{3-[3-methyl-2-oxo-1-(4-trifluoromethyl-benzyl)-imidazolidin-4-yl]-propyl}-phenoxy)-2-methyl-propionic acid



#### Step A

10 2-Methyl-2-(4-{3-[3-methyl-2-oxo-1-(4-trifluoromethyl-benzyl)-imidazolidin-4-yl]-propyl}-phenoxy)-propionic acid ethyl ester




15 The titled compound is prepared, according to the procedure of Example 3, Step D, using 4-[3-(4-Hydroxy-phenyl)-propyl]-3-methyl-1-(4-trifluoromethyl-benzyl)-imidazolidin-2-one (0.445g, 1.13 mmole) to produce an oil (0.464g, 85%). Mass [EI+] 507 (M+H)<sup>+</sup>.

20

#### Step B


2-(2-Iodo-4-{3-[3-methyl-2-oxo-1-(4-trifluoromethyl-benzyl)-imidazolidin-4-yl]-propyl}-phenoxy)-2-methyl-propionic acid ethyl ester



The titled compound is prepared, according to the procedure of **Example 3, Step A**, using 2-Methyl-2-(4-(3-[3-methyl-2-oxo-1-(4-trifluoromethyl-benzyl)-imidazolidin-4-yl]-propyl)-phenoxy)-propionic acid ethyl ester (0.460g, 0.960 mmole) to  
 5 produce an oil (0.273g, 48%). Mass [EI+] 633 (M+H)<sup>+</sup>.

#### Step C

2-(2-Butyl-4-{3-[3-methyl-2-oxo-1-(4-trifluoromethyl-benzyl)-imidazolidin-4-yl]-propyl}-phenoxy)-2-methyl-propionic acid ethyl ester  
 10



The titled compound is prepared, according to the procedure of **Example 3, Step B**, using 2-(2-Iodo-4-{3-[3-methyl-2-oxo-1-(4-trifluoromethyl-benzyl)-imidazolidin-4-yl]-propyl}-phenoxy)-2-methyl-propionic acid ethyl ester (0.270g, 0.430  
 15 mmole) and n-butyl boronic acid (0.218g, 0.215 mmole) to produce an oil (0.115g, 48%). Mass [EI+] 563 (M+H)<sup>+</sup>.

Step D

2-(2-Butyl-4-{3-[3-methyl-2-oxo-1-(4-trifluoromethyl-benzyl)-imidazolidin-4-yl]-propyl}-phenoxy)-2-methyl-propionic acid

5

The titled compound is prepared, according to the procedure of **Example 3, Step E**, using 2-(2-Butyl-4-{3-[3-methyl-2-oxo-1-(4-trifluoromethyl-benzyl)-imidazolidin-4-yl]-propyl}-phenoxy)-2-methyl-propionic acid ethyl ester (0.110g, 0.196 mmole) to produce an oil (0.029g, 28%). Mass [EI+] 535 (M+H)<sup>+</sup>, [EI-] 533 (M-H)<sup>-</sup>.


10

The following Examples are prepared substantially as described herein above to yield the title compounds:

15

**Example 5**


**2-Methyl-3-(4-{2-[3-methyl-2-oxo-1-(3-phenoxy-benzyl)-imidazolidin-4-yl]-ethoxy}-phenyl)-2-phenoxy-propionic acid**



20 Yield(0.135g, 86%). Mass [EI+] 581 (M+H)<sup>+</sup>, [EI-] 579 (M-H)<sup>-</sup>

**Example 6**

**2-Methyl-3-(4-{2-[3-methyl-2-oxo-1-(3-phenoxy-benzyl)-imidazolidin-4-yl]-ethoxy}-phenyl)-2-phenoxy-propionic acid**



Yield(0.192g, 52%). Mass [EI+] 581 (M+H)<sup>+</sup>, [EI-] 579 (M-H)<sup>-</sup>


#### Example 7

2-Methyl-3-(4-{2-[3-methyl-2-oxo-1-(3-trifluoromethoxy-  
benzyl)-imidazolidin-4-yl]-ethoxy}-phenyl)-2-phenoxy-  
propionic acid

Yield(0.127g, 35%). Mass [EI+] 573 (M+H)<sup>+</sup>, [EI-] 571 (M-H)<sup>-</sup>

#### Example 8

2-Methyl-3-(4-{2-[3-methyl-2-oxo-1-(3-trifluoromethoxy-  
benzyl)-imidazolidin-4-yl]-ethoxy}-phenyl)-2-phenoxy-  
propionic acid




Yield (0.238g, 63%). Mass [EI+] 573 (M+H)<sup>+</sup>, [EI-] 571 (M-H)<sup>-</sup>


#### Example 9

2-(2-Fluoro-phenoxy)-3-(4-{2-[1-(4-methoxy-benzyl)-3-methyl-  
2-oxo-imidazolidin-4-yl]-ethoxy}-phenyl)-2-methyl-propionic  
acid



Step A

2-(2-Fluoro-phenoxy)-3-(4-hydroxy-phenyl)-2-methyl-propionic  
acid methyl ester




To a solution of LDA (1.5 M in cyclohexane, 33 mL, 49.5  
5 mmole) in THF (40 mL), at  $-78^{\circ}\text{C}$ , is added dropwise a  
solution of 2-(2-fluoro-phenoxy)-propionic acid ethyl ester  
(7.70g, 36.3 mmole) in THF (100 mL). After stirring for 30  
minutes at  $-78^{\circ}\text{C}$ , a solution of toluene-4-sulfonic acid 4-  
iodomethyl-phenyl ester (12.8g, 33.0 mmole) is injected to  
10 the reaction mixture. The reaction is stirred at  $-78^{\circ}\text{C}$  for  
one hour, then warmed up to room temperature and stirred  
overnight. The reaction is treated with 5.0N NaOH (50 mL)  
in presence of methanol (200 mL). After evaporating  
methanol, the aqueous solution is acidified with  
15 concentrated HCl to pH = 1 and is extracted with ethyl  
acetate (2x300 mL). The combined organic layers are dried,  
filtered and concentrated in vacuo. The residue is  
dissolved in MeOH (200 mL) and treated with concentrated  
sulfuric acid (2 mL) at  $80^{\circ}\text{C}$  overnight. The reaction  
20 mixture is partitioned between ethyl acetate (500 mL) and  
water (500 mL), the organic layer is washed by brine (3x500  
mL), dried, filtered and concentrated in vacuo. The  
compound is purified by chromatography (silica gel, gradient  
elution 0-20% ethyl acetate in hexane) to provide an oil

(6.02g, 60%), which is then resolved by chiral HPLC to generate the enantiomer as an oil (2.30g). Mass [EI+] 322 ( $M+NH_4$ )<sup>+</sup>, [EI-] 303 ( $M-H$ )<sup>-</sup>.

## 5

### Step B

2-(2-Fluoro-phenoxy)-3-(4-{2-[1-(4-methoxy-benzyl)-3-methyl-  
2-oxo-imidazolidin-4-yl]-ethoxy}-phenyl)-2-methyl-propionic  
acid methyl ester



The titled compound is prepared, according to the procedure of **Example 1, Step A**, using toluene-4-sulfonic acid 2-[1-(4-methoxy-benzyl)-3-methyl-2-oxo-imidazolidin-4-yl]-ethyl ester (0.888g, 2.12 mmole) and 2-(2-Fluoro-phenoxy)-3-(4-hydroxy-phenyl)-2-methyl-propionic acid methylester (0.645g, 2.12 mmole) to produce an oil (0.95g, 81%). Mass [EI+] 551 (M+H)<sup>+</sup>.

### Step C


2-(2-Fluoro-phenoxy)-3-(4-{2-[1-(4-methoxy-benzyl)-3-methyl-  
2-oxo-imidazolidin-4-yl]-ethoxy}-phenyl)-2-methyl-propionic  
acid

20 The titled compound is prepared, according to the procedure  
of **Example 3, Step E**, using 2-(2-Fluoro-phenoxy)-3-(4-{2-[1-  
(4-methoxy-benzyl)-3-methyl-2-oxo-imidazolidin-4-yl]-  
ethoxy}-phenyl)-2-methyl-propionic acid methyl ester  
(0.224g, 0.406 mmole) to produce a white foamy solid  
25 (0.199g, 91%). Mass [EI+] 537 (M+H)<sup>+</sup>, [EI-] 535 (M-H)<sup>-</sup>.

**Example 10**

**3-{4-[2-(1-Biphenyl-4-ylmethyl-3-methyl-2-oxo-imidazolidin-4-yl)-ethoxy]-phenyl}-2-(2-fluoro-phenoxy)-2-methyl-propionic acid**

5




The titled compound is prepared using substantially the processes described herein above to produce a white foamy solid (0.0858g, 40%). Mass [EI+] 583 (M+H)<sup>+</sup>, [EI-] 581 (M-H)<sup>-</sup>.

10

**Example 11**


**3-(4-{2-[1-(4-tert-Butyl-benzyl)-3-methyl-2-oxo-imidazolidin-4-yl]-ethoxy}-phenyl)-2-(2-fluoro-phenoxy)-2-methyl-propionic acid**



15 The titled compound is prepared, according to the procedures substantially as described hereinabove to produce a white foamy solid (0.101g, 77%). Mass [EI+] 563 (M+H)<sup>+</sup>, [EI-] 561 (M-H)<sup>-</sup>.

**Example 12**


20 **2-(2-Fluoro-phenoxy)-2-methyl-3-{4-[2-(3-methyl-1-naphthalen-2-ylmethyl-2-oxo-imidazolidin-4-yl)-ethoxy]-phenyl}-propionic acid**



The titled compound is prepared, substantially as described herein above to yield (0.0753g, 58%). Mass [EI+] 543


5 (M+H)<sup>+</sup>, [EI-] 541 (M-H)<sup>-</sup>.

### Example 13




10

#### Step A





A methylene chloride solution (300 ml) of 4-(4-methoxyphenyl)butanol (10.0g, 0.055mol) is cooled to 0 °C and stirred. Pyridinium chlorochromate (17.8g, 0.082 mol) is added slowly to the solution which is stirred, placed under a drying tube and warmed to room temperature overnight. The reaction mixture is filtered and the filtrate washed with aqueous brine then dried over sodium sulfate. Evaporation of the solvent gives the crude product as an oil. Purification by flash chromatography (hexanes:ethyl acetate) gives the desired aldehyde (6.8g).  
 20 C<sub>11</sub>H<sub>14</sub>O<sub>2</sub> (MW = 178.23); <sup>1</sup>H NMR

Step B

Potassium cyanide (10.3g, 0.158 mol) and ammonium carbonate (38.1g, 0.396 mol) are combined in methanol (100ml) and water (100ml). The product from Example 13, Step A (6.7g, 0.038 mol) is added and the reaction is stirred while heating to 45 °C overnight. The reaction is cooled and the precipitate filtered and washed with water. The obtained solid is dissolved in THF and dried with sodium sulfate.

Evaporation of the solvent results in a solid which is washed with ethyl acetate:hexanes (50:50) then air-dried to yield the desired hydantoin (16.6g).

$C_{13}H_{16}N_2O_3$  (MW = 248.3);  $^1H$  NMR


Example 14Step A**Preparation of:**

To a cooled (0°C) solution of boron tribromide (50g, 200mmol) in  $CH_2Cl_2$  (50 ml) is added a solution of methyl 4-(4-methoxyphenyl)butyrate (15.5g, 74.4mmol) in  $CH_2Cl_2$


- (100ml) dropwise over one hour. After stirring for an additional hour at 0°C, the reaction mixture is treated with 1:1 CH<sub>3</sub>OH:CH<sub>2</sub>Cl<sub>2</sub> (120 ml) with cooling and stirred overnight at ambient temperature. Concentration of the mixture gives
- 5 an oil which is partitioned between ethyl acetate (150ml) and water (150ml). The aqueous layer is extracted with ethyl acetate (2 x 50ml), and the combined organic extracts washed with water (50ml), brine (50ml), dried, then concentrated to give the desired phenol as an oil.
- 10 C<sub>11</sub>H<sub>14</sub>O<sub>3</sub> (MW = 194.23); MS: m/z (M<sup>+</sup> + 1) = 195

#### Step B


##### **Preparation of:**



- 15 The phenol from **Step A** (18.6g, 96 mmol) is dissolved in DMF (300ml) and treated with t-butyl 2-bromoisobutyrate (50ml, 288mmol), powdered K<sub>2</sub>CO<sub>3</sub> (53.0g, 384mmol) and MgSO<sub>4</sub> (1.6g, 96mmol), and the resulting mixture heated at 75°C overnight. After cooling to ambient temperature, the reaction mixture
- 20 is decanted into 1N aqueous HCl (300ml) and extracted with diethyl ether (3 x 150ml). The remaining solids from the decantation are washed several times with diethyl ether. The diethyl ether extracts and washes are combined and washed with 1N aqueous HCl (150ml), dried (Na<sub>2</sub>SO<sub>4</sub>), and
- 25 concentrated to a dark oil. Purification by flash chromatography (gradient elution, hexanes to 95:5 hexanes:ethyl acetate) gives the desired ether as an oil.
- C<sub>19</sub>H<sub>28</sub>O<sub>5</sub> (MW = 336.43); MS: m/z (M<sup>+</sup> + 1) = 337

Step C


The diester from **Step B** (10.7 g, 0.032 mol) is dissolved in dioxane (100 mL) then treated with an aqueous solution (50  
5 mL) of LiOH (1.5 g, 0.063 mol). The resulting mixture is stirred at room temperature for 2 hr at which time TLC indicates complete reaction. The solvent is concentrated to approximately 10 mL which is then diluted with H<sub>2</sub>O (200 mL) and washed with ether (1x100 mL). The aqueous extract is  
10 acidified with 5 N HCl (25 mL) and extracted with ethyl acetate (2x150 mL). The combined organic extracts are dried over Na<sub>2</sub>SO<sub>4</sub> and concentrated to give the desired carboxylic acid (10.6 g, >99%).  
C<sub>18</sub>H<sub>26</sub>O<sub>5</sub> (MW = 322.40); mass spectroscopy: (M+NH<sub>4</sub><sup>+</sup>) = 340.3,  
15 (M-H) = 321.3

Step D

A THF solution (100 mL) of the carboxylic acid from **Step C** (10 g, 0.031 mol) is cooled to 0 °C and treated with  
20 triethyl amine (5.6 mL, 0.040 mol) followed by ethyl chloroformate (3.8 mL, 0.040 mol). A thick white precipitate forms almost immediately. The reaction mixture is warmed to room temperature, and after stirring overnight, the mixture is filtered. The filtrate is cooled to 0 °C and


treated with sodium borohydride (3.5 g, 0.092 mol). A solution of methanol (20 mL) in THF (40 mL) is added dropwise over a 30-40 minute period then the resulting mixture is warmed to room temperature and stirred for 1  
5 hour. Again the solution is cooled to 0 °C and carefully quenched by the addition of 1 N HCl (100 mL). The mixture is diluted with water (400 mL) and extracted with CH<sub>2</sub>Cl<sub>2</sub> (2 x 300 mL). The combined organic extracts are dried over Na<sub>2</sub>SO<sub>4</sub>, filtered and concentrated. The crude oil is  
10 purified by flash chromatography (5:1 hexanes:ethyl acetate) to give the desired alcohol as a colorless oil (8.69 g, 88%).

Step E





15 The alcohol from **Step D** (8.7 g, 0.028 mol) is dissolved in CH<sub>2</sub>Cl<sub>2</sub> and cooled to 0 °C. Pyridinium chlorochromate (9.0 g, 0.042 mol) was added to the solution in portions after which the mixture is warmed to room temperature. After stirring overnight the reaction mixture is filtered using celite.  
20 The filtrate is concentrated and the resulting crude oil is purified by chromatography (10:1 hexanes:ethyl acetate) to give the desired aldehyde as a colorless oil (4.45 g, 52%).




Step F

Potassium cyanide (1.1 g, 7.2 mmol) and ammonium carbonate (1.7 g, 18.0 mmol) are combined in methanol (15 mL) and  
5 water (15 mL). The aldehyde from **Step E** (1.1 g, 3.6 mmol) is added and the reaction is stirred while heating to 50 °C. After 3 hr, additional ammonium carbonate is added (1.7 g, 18.0 mmol) and the resulting mixture is stirred overnight at 50 °C. The reaction is cooled to room temperature, diluted  
10 with water, then extracted with ethyl acetate (2X). The combined organic extracts are washed with brine, dried over Na<sub>2</sub>SO<sub>4</sub>, and concentrated. Purification by chromatography (gradient: 10:1 to 3:1 to 3:2 hexanes:ethyl acetate) gives the desired hydantoin (0.9 g, 66%)  
15 C<sub>20</sub>H<sub>28</sub>N<sub>2</sub>O<sub>5</sub> (MW = 376.46); mass spectroscopy: (MH<sup>+</sup>-t-butyl) = 321.2, (MH<sup>-</sup>) = 375.4

**Example 15**Step A

The hydantoin from **Example 14** (1.0g, 2.66 mmol) and 4-(tert-butyl)benzyl bromide (0.63ml, 2.93 mmol) are stirred together in DMF (20ml). Potassium carbonate (powdered, 1.5g, 10.64 mmol) and magnesium sulfate (0.5g, 4.00 mmol) are  
5 added and the mixture stirred at ambient temperature overnight. The reaction mixture is carefully added to 1 N hydrochloric acid (50ml) and the resulting solution is extracted twice with ethyl acetate. The organic layers are combined, washed with aqueous brine then dried over sodium  
10 sulfate. Evaporation of the solvent and subsequent purification by flash chromatography (hexanes:ethyl acetate) gives the desired product (1.05g).

$C_{31}H_{42}N_2O_5$  (MW = 522.7)




15

#### Step B


The hydantoin from **Example 15, Step A** (0.15g, 0.29mmol) is stirred in ethanol, sodium borohydride (0.11g, 2.90mmol) added slowly and the mixture was stirred at ambient temperature, under a drying tube, overnight. The reaction  
20 mixture was concentrated and the residue dissolved in ethyl acetate, washed with water and aqueous brine then dried. Evaporation of the solvent and subsequent purification by flash chromatography (ethyl hexanes:ethyl acetate) gives the desired product (0.071g).

25  $C_{31}H_{42}N_2O_4$  (MW = 506.7)


Step C

The product from **Example 15, Step B** (0.071g, 0.14mmol) is stirred in methylene chloride (4ml) with trifluoroacetic acid (0.1ml, 1.3mmol), under a drying tube, overnight. The solvent is evaporated and subsequent purification by flash chromatography (ethyl acetate:hexanes) gives the desired product (0.035g).

$C_{27}H_{34}N_2O_4$  (MW = 450.6); MS ( $M^+$ , 451.3)


**Example 16**

$C_{25}H_{30}N_2O_4$  (MW = 422.5); MS ( $M^+$ , 423.3)

Step A


Following the procedure of **Example 15, Step A**, the product was obtained using the hydantoin from **Example 14** (1.0g, 2.66 mmol) and 3,4-dimethylbenzyl chloride (Transworld, 0.42ml, 2.93mol). This reaction was heated to 50 °C to affect reaction completion, yielding the product (0.550g).

$C_{29}H_{38}N_2O_5$  (MW = 494.6)

Step B

Following the procedure of **Example 15, Step B**, the alkylated hydantoin from **Example 16, Step A** (0.15g, 0.3mmol) and  
5 sodium borohydride (0.11g, 3.00mmol) were used to yield the desired imidazolone (0.032g).


$C_{29}H_{38}N_2O_4$  (MW = 478.6)

Step C

10

Following the procedure of **Example 15, Step C**, the imidazolone from **Step B** (0.032g, 0.066mmol) and trifluoroacetic acid (0.1ml) were used to yield the desired product (0.026g).


15  $C_{25}H_{30}N_2O_4$  (MW = 422.5); MS ( $M^+$ , 423.3)

**Example 17**

The title compound is prepared using substantially the procedures described herein above. Yield: 26% for two steps.

$C_{24}H_{28}N_2O_5$  (MW = 424.50); mass spectroscopy:  $(MH^+) = 425.3$


5

**Example 18**

The title compound is prepared using substantially the procedures described herein above.

10



$C_{26}H_{27}N_3O_4$  (MW = 445.5); MS ( $M^+$ , 446.3)

**Example 19**


15

The title compound is prepared using substantially the procedures described herein above to yield the product.


$C_{24}H_{28}N_2O_4$  (MW = 408.50); mass spectroscopy:  $(MH^+) = 409.2$

**Example 20**Step A

5 The hydantoin from **Example 14** (0.500 g, 0.0013 mol) is dissolved in DMF and treated with 3, 5 difluoro bromobenzene (0.189 g, 0.0015 mol) and powdered  $K_2CO_3$  (0.718g, 0.0052 mol). The resulting mixture is stirred at room temperature  
10 overnight. The reaction mixture is poured into 1N HCl and combined with ethyl acetate. The organic layer is extracted and washed with brine then dried and concentrated. Purification of the crude material by flash chromatography (4:1 hexanes: ethyl acetate, 1:1 hexanes: ethyl acetate)  
15 yields the desired hydantoin as a clear oil (0.420 g, 64 %).  $C_{27}H_{32}F_2N_2O_5$  (MW =502.23); mass spectroscopy (MH+) = 447.1

Step B



The hydantoin from **Step A** (0.410 g, 0.00082 mol) is dissolved in methylene chloride (5 ml) and treated with trifluoroacetic acid (0.315 ml, 0.0041 mol) and stirred overnight. The solvent is concentrated and the product is vacuum dried.  $C_{23}H_{24}F_2N_2O_5$  (MW = 446.17); mass spectroscopy (MH<sup>+</sup>) = 447.2

Step C

10

The acid from **Step B** (0.428 g, 0.00096 mol) is dissolved in ethanol (20 ml) and treated with sodium borohydride (0.363 g, 0.0096 mol). One hour later, additional sodium borohydride (0.363 g) is added and the reaction is stirred overnight. Fifty milliliters of 5N HCl is added to the reaction mixture followed by water (50 ml). The aqueous solution is extracted with ethyl acetate (2X, 50 ml). The organic layer is washed, dried, and concentrated. Purification by flash chromatography (100% ethyl acetate) gives the desired acid as a burgundy solid (150 g, 36 %).  $C_{23}H_{24}F_2N_2O_4$  (MW = 430.17); mass spectroscopy (MH<sup>+</sup>) = 431.1

20

**Example 21**Step A

5

A DMF solution (20 mL) of the hydantoin from **Example 14** (455.2 mg, 1.21 mmol) is treated sequentially with benzyl bromide (160  $\mu$ L, 1.35 mmol),  $K_2CO_3$  (0.44 g, 3.2 mmol), and  $MgSO_4$  (0.48 g, 4.0 mmol). The resulting mixture is stirred at room temperature overnight. The reaction is quenched by the slow addition of 1 N HCl. The resulting mixture is extracted with ethyl acetate (2x75 mL). The combined organic extracts are washed with brine then dried over  $Na_2SO_4$  and concentrated. Purification by chromatography (2:1 hexanes:ethyl acetate) gives the desired alkylated hydantoin as a white solid (512.0 mg, 91%).

15

$C_{27}H_{34}N_2O_5$  (MW = 466.58); mass spectroscopy:  $(MH^+ - t\text{-butyl}) = 411.1$ ,  $(MH^-) = 465.3$



Step B


A  $\text{CH}_2\text{Cl}_2$  solution (15 mL) of the ester from **Step A** (495.5 mg, 1.06 mmol) is cooled to 0 °C and treated with TFA (2 mL, 5 26 mmol). The mixture is warmed to room temperature and stirred overnight. The solvent is concentrated to give the crude acid. Trace amounts of TFA are removed by concentration from toluene (3x15 mL) after which the product is used in the subsequent reaction without further 10 purification.

$\text{C}_{23}\text{H}_{26}\text{N}_2\text{O}_5$  (MW = 410.47); mass spectroscopy:  $(\text{MH}^+) = 411.1$ ,  $(\text{MH}^-) = 409.2$


Step C

15 The hydantoin from **Step B** is dissolved in ethanol (20 mL) and treated with  $\text{NaBH}_4$  (400 mg, 10.5 mmol). Additional  $\text{NaBH}_4$  is added to the reaction mixture after 3 hr (200 mg, 4.8 mmol). The resulting mixture is stirred at room 20 temperature overnight. Additional  $\text{NaBH}_4$  is added to the reaction mixture (370 mg, 9.7 mmol) and stirring is continued for about two days. The reaction is quenched by the careful addition of 5 N HCl. The mixture is stirred for 30 minutes then diluted with  $\text{H}_2\text{O}$  and extracted with ethyl 25 acetate. The combined organic extracts are washed, dried, and concentrated. Purification by chromatography (gradient: 100%  $\text{CH}_2\text{Cl}_2$  to 5% methanol in  $\text{CH}_2\text{Cl}_2$ ) gives the desired imidazolone as a glass-like solid (93.2 mg, 22% for two steps).

30  $\text{C}_{23}\text{H}_{26}\text{N}_2\text{O}_4$  (MW = 394.47); mass spectroscopy:  $(\text{MH}^+) = 395.1$ ,  $(\text{MH}^-) = 393.2$

**Example 22**Step A

The corresponding hydantoin is prepared using substantially  
5 the procedures described herein above.



Yield: 84%.

$C_{31}H_{36}N_2O_5$  (MW = 516.64); mass spectroscopy:  $(MH^+ - t\text{-butyl}) =$   
461.2,  $(MH^-) = 515.4$


10

Step B

A  $CH_2Cl_2$  solution (15 mL) of the ester from **Step A** (387.8  
mg, 0.751 mmol) is cooled to 0 °C and treated with TFA (2  
mL, 26 mmol). The mixture is warmed to room temperature and  
15 stirred overnight. The solvent is concentrated to give the  
crude acid. Trace amounts of TFA are removed by  
concentration from toluene (3x15 mL) after which the product  
is used in the subsequent reaction without further  
purification.

$C_{27}H_{28}N_2O_5$  (MW = 460.53); mass spectroscopy:  $(MH^+) = 461.2$ ,  
 $(MH^-) = 459.3$


Step C



5 The hydantoin from **Step B** is dissolved in ethanol:THF (2:1, 30 mL) and treated with  $NaBH_4$  (313 mg, 8.2 mmol). Additional  $NaBH_4$  is added to the reaction mixture after 3 hr (180 mg, 4.7 mmol). The resulting mixture is stirred at room temperature overnight. The reaction is quenched by the  
10 careful addition of 5 N HCl. The mixture is stirred for 30 minutes then diluted with  $H_2O$  (75 mL) and extracted with ethyl acetate (2x75 mL). The combined organic extracts are washed, dried, and concentrated. Purification by chromatography (gradient: 100%  $CH_2Cl_2$  to 5% methanol in  
15  $CH_2Cl_2$ ) gives the desired imidazolone as a white foam (109.0 mg, 33% for two steps).


$C_{27}H_{28}N_2O_4$  (MW = 444.54); mass spectroscopy:  $(MH^+) = 445.2$ ,  
 $(MH^-) = 443.1$

The following Examples are prepared using substantially the  
20 procedures described herein before.

**Example 23**


C<sub>24</sub>H<sub>28</sub>N<sub>2</sub>O<sub>4</sub> (MW = 408.50); mass spectroscopy: (MH<sup>+</sup>) = 409.2,  
(MH<sup>-</sup>) = 407.1

5


**Example 24**

C<sub>26</sub>H<sub>32</sub>N<sub>2</sub>O<sub>4</sub> (MW = 436.56); mass spectroscopy: (MH<sup>+</sup>) = 437.2,  
(MH<sup>-</sup>) = 435.1

10


**Example 25**

15

Step A


Following the procedure of **Example 15, Step A**, the product was obtained using the hydantoin from **Example 14** (0.5g, 1.33 mmol) and 4-ethylbenzyl chloride (Aldrich, 0.2ml, 1.46mol). This reaction was heated to 45 °C to affect reaction completion, yielding, after purification, the product (0.41g).

$C_{29}H_{38}N_2O_5$  (MW = 494.6); MS ( $M^+$ , 495.3,  $M+18$ , 512.3)

Step B

The product from **Example 25, Step A** (0.4g, 0.8mmol) was stirred in methylene chloride (10ml) with trifluoroacetic acid (Aldrich, 2 ml), under a drying tube, overnight. The solvent was evaporated and the obtained residue was placed under vacuum to yield the desired carboxylic acid (0.36g).

$C_{25}H_{30}N_2O_5$  (MW = 438.5); MS ( $M^-$ , 437.1)

Step C

The product from **Example 25, Step B** (0.33g, 0.75mmol) was stirred in ethanol (30ml) and sodium borohydride (0.28g, 7.5mmol) added slowly. After stirring overnight, 5 N


hydrochloric acid (5ml) was added slowly and allowed to stir for 1 hr. Water (50ml) added and the mixture was extracted twice with ethyl acetate. The organic layers were combined, washed with aqueous brine and dried over sodium sulfate.

5 Evaporation of the solvent and subsequent purification by flash chromatography (methylene chloride:methanol) gave the desired product (0.120g).


$C_{25}H_{30}N_2O_4$  (MW = 422.5); MS ( $M^+$ , 423.3,  $M^-$ , 421.1)

10

### Example 26



### Step A




A DMF solution (75 mL) of the hydantoin from **Example 13** (1.5 g, 6.0 mmol) is treated sequentially with benzyl bromide (1.8 mL, 15 mmol),  $K_2CO_3$  (3.7 g, 27 mmol), and  $MgSO_4$  (4.2 g, 35 mmol). The resulting mixture is stirred at room temperature overnight. The reaction is quenched by the slow addition of 1 N HCl (100 mL). The resulting mixture is extracted with ethyl acetate (2x100 mL). The combined organic extracts are washed with brine then dried over  $Na_2SO_4$  and concentrated. Purification by chromatography

(5:1 hexanes:ethyl acetate) gives the desired alkylated hydantoin as a colorless oil (1.33 g, 52%).


$C_{27}H_{28}N_2O_3$  (MW = 428.21); mass spectroscopy:  $(MH^+) = 429.2$

5

Step B


To a slurry of LAH (122 mg, 3.2 mmol) in THF (10 mL) at 0 °C under  $N_2$  is added the hydantoin from **Step A** (1.02 g, 2.38 mmol) as a solution in THF (10 mL). After 10 minutes the reaction is quenched by the addition of 5 N HCl (4 mL) in THF (2 mL), stirred for 30 minutes, then diluted with  $H_2O$  (75 mL). The resulting mixture is extracted with ethyl acetate (2x75 mL). The combined organic extracts are washed with brine, dried and concentrated. Purification by chromatography (gradient: 5:1 to 1:1 hexanes:ethyl acetate) gives the desired imidazolone as a colorless oil (0.873 g, 89%).

20

Step C

The imidazolone from **Step B** (0.81 g, 2.0 mmol) is dissolved in CH<sub>2</sub>Cl<sub>2</sub> and cooled to 0 °C under an atmosphere of N<sub>2</sub>. A solution of BBr<sub>3</sub> (600 µL, 6.3 mmol) in CH<sub>2</sub>Cl<sub>2</sub> (15 mL) is added dropwise, then the reaction mixture is warmed to room temperature. After 1 hr, the solution is again cooled to 0 °C and quenched by the slow addition of a methanol (10 mL) solution in CH<sub>2</sub>Cl<sub>2</sub> (50 mL). The resulting mixture is extracted with H<sub>2</sub>O (150 mL). The organic extract is washed with brine, dried over Na<sub>2</sub>SO<sub>4</sub>, and concentrated to give the desired phenol as a foam-like solid (0.78 g, 98%).  
C<sub>26</sub>H<sub>26</sub>N<sub>2</sub>O<sub>2</sub> (MW = 398.51); mass spectroscopy: (MH<sup>+</sup>) = 399.2, (MH<sup>-</sup>) = 397.3

Step D




An ethanol solution (40 mL) of the phenol from **Step C** (0.78 g, 1.9 mmol) is treated sequentially with ethyl 2-bromoisobutyrate (1.5 mL, 10.2 mmol), K<sub>2</sub>CO<sub>3</sub> (1.4 g, 10.1 mmol), and MgSO<sub>4</sub> (1.2 g, 10 mmol). The resulting mixture is heated to 50 °C for two days. The reaction is quenched by the slow addition of 1 N HCl (15 mL) then poured into 0.5 N HCl (100 mL). The resulting mixture is extracted with ethyl acetate (2x70 mL). The combined organic extracts are washed, then dried and concentrated. Purification by chromatography (gradient: 3:1 to 1:1 hexanes:ethyl acetate)



gives the desired ester as a slightly yellow oil (0.79 g, 81%).

$C_{32}H_{36}N_2O_4$  (MW = 512.65); mass spectroscopy:  $(MH^+) = 513.3$

5

Step E

A dioxane solution (15 mL) of the ester from **Step D** (0.741, 1.45 mmol) is treated with an aqueous solution (5 mL) of LiOH (120.4 mg, 5.0 mmol). After stirring overnight the solvent is concentrated and the resulting oil is diluted with  $H_2O$  (70 mL) and extracted with  $Et_2O$  (70 mL). The aqueous extract is acidified with 1 N HCl and extracted with ethyl acetate (2x70 mL). The combined organic extracts are washed, dried, and concentrated to give the desired carboxylic acid as a slightly yellow foam-like solid (0.601 g, 86%)

10


15

$C_{30}H_{32}N_2O_4$  (MW = 484.60); mass spectroscopy:  $(MH^+) = 485.3$ ,  $(MH^-) = 483.3$


20

**Example 27**


**3-{4-[2-(1-Benzyl-3-methyl-2-oxo-imidazolidin-4-yl)-ethoxy]-phenyl}-2-(2-fluoro-phenoxy)-2-methyl-propionic acid**



Step A



**2-(2-Fluoro-phenoxy)-propionic acid ethyl ester:** Cesium carbonate (65.69g, 201.61mmol) is added to a solution of 2-fluorophenol (9mL, 100.84mmol, d=1.256) in anhydrous DMF (300mL) at room temperature under an atmosphere of nitrogen. After five minutes, ethyl 2-bromopropionate (13.1mL, 100.84mmol, d=1.394) is added rapidly dropwise and the resultant mixture is allowed to stir at 90°C for 18h. The reaction mixture is diluted with diethyl ether, then extracted twice with 1N HCl and twice with water. The organic layer is dried, concentrated, and purified by flash chromatography (25% ether in hexanes) to provide the titled compound (20.14g, 94%). <sup>1</sup>H NMR (400MHz, CDCl<sub>3</sub>): δ 7.07-6.96 (m, 2H), 6.93-6.87 (m, 2H), 4.73, 4.71 (ABq, 1H, J = 6.7 Hz), 4.21-4.15 (m, 2H), 1.61 (d, 3H, J = 6.7 Hz), 1.21 (t, 3H, J = 6.7 Hz). MS [EI+] 507 (M+H)<sup>+</sup>. R<sub>f</sub>=0.39 in 20% acetone in hexanes.


Step B

**3-(4-Benzyloxy-phenyl)-2-(2-fluoro-phenoxy)-3-hydroxy-2-methyl-propionic acid ethyl ester:** A solution of LDA


5 (49.6mL, 99.15mmol, 2M in cyclohexane) in anhydrous THF (150mL) is cooled to  $-78^{\circ}\text{C}$  in a dry ice/acetone bath and added to a solution of 2-(2-Fluoro-phenoxy)-propionic acid ethyl ester in anhydrous THF (150mL) also cooled to  $-78^{\circ}\text{C}$  under an atmosphere of nitrogen. After five minutes, 4-  
10 benzyloxybenzaldehyde (10.52g, 99.15mmol) is added in one portion. After stirring for one minute, the reaction mixture is quenched with acetic acid (9.5mL, 165.3mmol,  $d=1.049$ ) and a saturated solution of aqueous  $\text{NH}_4\text{Cl}$  (100mL). The biphasic mixture is allowed to warm to room temperature  
15 and is diluted with diethyl ether (1L). The organic layer is washed, dried, and concentrated. The residue is purified by flash chromatography (20% acetone in hexanes) to provide a mixture of diastereomers of 3-(4-Benzyloxy-phenyl)-2-(2-fluoro-phenoxy)-3-hydroxy-2-methyl-propionic acid ethyl  
20 ester (16.21g, 69%).  $^1\text{H}$  NMR (400MHz,  $\text{CDCl}_3$ ):  $\delta$  7.45-7.33 (m, 7H), 7.09-6.95 (m, 6H), 5.07 (s, 2H), 4.21-4.17 (m, 2H), 3.10 (d, 1H,  $J = 5.4$  Hz), 1.39 (s, 3H), 1.20 (t, 3H,  $J = 6.1$  Hz)  $R_f=0.06$  in 25% acetone in hexanes.

25


Step C



**3-(4-Benzyloxy-phenyl)-2-(2-fluoro-phenoxy)-2-methyl-propionic acid ethyl ester:** Boron trifluoride etherate (21.62mL, 171.85mmol, d=1.128) is added dropwise rapidly to a 0°C solution of 3-(4-Benzyloxy-phenyl)-2-(2-fluoro-phenoxy)-2-methyl-propionic acid ethyl ester (16.21g, 38.19mmol) and triethylsilane (27.45mL, 171.85mmol, d=0.728) in anhydrous CH<sub>2</sub>Cl<sub>2</sub> (150mL). The mixture is stirred for 90 minutes, gradually warming to ambient temperature. The reaction mixture is quenched with a saturated solution of aqueous sodium carbonate and extracted with CH<sub>2</sub>Cl<sub>2</sub>. The organic layer is dried over MgSO<sub>4</sub>, concentrated *in vacuo*, and purified by flash chromatography (20 % acetone in hexanes) to provide the titled compound (12.8g, 82%). <sup>1</sup>H NMR (400MHz, CDCl<sub>3</sub>): δ 7.46 (d, 2H, J = 6.7 Hz), 7.42 (d, 2H, J = 6.7 Hz), 7.39-7.33 (m, 1H), 7.26 (d, 2H, J = 8.2 Hz), 7.11-7.06 (m, 1H), 7.01-6.93 (m, 5H), 4.23 (q, 2H, J = 7.2 Hz), 3.31, 3.21 (AB<sub>q</sub>, 2H, J = 13.4 Hz), 1.45 (s, 3H), 1.26 (t, 3H, J = 7.2 Hz). R<sub>f</sub>=0.25 in 20% acetone in hexanes.


Step D

**2-(2-Fluoro-phenoxy)-3-(4-hydroxy-phenyl)-2-methyl-propionic acid ethyl ester:** 3-(4-Benzyloxy-phenyl)-2-(2-fluoro-phenoxy)-2-methyl-propionic acid ethyl ester (12.8g, 31.34mmol) is dissolved in ethanol (500mL), treated with 10% palladium on carbon (6.0g), and stirred under an atmosphere of hydrogen for 90 minutes. The suspension is filtered through celite and concentrated to provide 2-(2-Fluoro-phenoxy)-3-(4-hydroxy-phenyl)-2-methyl-propionic acid ethyl ester (9.65g, 97%). The title compound is separated by chiral chromatography (Column OJ, 40% IPA in heptane, 1mL/Mmin, 240nm UV.  $^1\text{H}$  NMR (400MHz,  $\text{CDCl}_3$ ):  $\delta$  7.13 (d, 2H,  $J = 8.0$  Hz), 7.07-7.02 (m, 1H), 6.98-6.88 (m, 3H), 6.74 (d, 2H,  $J = 8.0$  Hz), 6.05 (s, 1H), 4.20 (q, 2H,  $J = 6.9$  Hz), 3.24, 3.16 (AB<sub>q</sub>, 2H,  $J = 13.4$  Hz), 1.39 (s, 3H), 1.23 (t, 3H,  $J = 6.8$  Hz).

Step E


2-(2-Fluoro-phenoxy)-3-(4-{2-[1-(4-methoxy-benzyl)-3-methyl-2-oxo-imidazolidin-4-yl]-ethoxy}-phenyl)-2-methyl-propionic acid ethyl ester : Cesium carbonate (8.23g, 25.3mmol) is added to a solution of 2-(2-Fluoro-phenoxy)-3-(4-hydroxy-phenyl)-2-methyl-propionic acid ethyl ester (2.68g, 8.43mmol) and toluene-4-sulfonic acid 2-[1-(4-methoxy-benzyl)-3-methyl-2-oxo-imidazolidin-4-yl]-ethyl ester (3.88g, 9.27mmol) in DMF (30mL). The resultant mixture is stirred at 65°C under an atmosphere of nitrogen for 18 h, then diluted with ethyl acetate. The organic layer is washed with 1N HCl, water, and brine. The organic layer is dried, concentrated *in vacuo*, and purified by flash chromatography (17% acetone in hexanes) to provide the title compound (4.4g, 93%). <sup>1</sup>H NMR 400MHz, CDCl<sub>3</sub>): δ 7.19-7.15 (m, 4H), 7.06-7.01 (m, 1H), 6.97-6.87 (m, 3H), 6.83 (d, 2H, *J* = 8.6 Hz), 6.74 (d, 2H, *J* = 8.6 Hz), 4.31, 4.27 (AB<sub>q</sub>, 2H, *J* = 14.8 Hz), 4.21, 4.17 (AB<sub>q</sub>, 2H, *J* = 7.2 Hz), 3.96-3.92 (m, 2H), 3.76 (s, 3H), 3.60-3.53 (m, 1H), 3.33 (t, 1H, *J* = 8.6 Hz), 3.25, 3.13 (AB<sub>q</sub>, 2H, *J* = 13.8 Hz), 2.994-2.83 (m, 2H), 2.82 (s, 3H), 2.24-2.16 (m, 1H), 1.89-1.80 (m, 1H), 1.37 (s, 3H), 1.22 (t, 3H, *J* = 7.2 Hz). MS [EI+] 565 (M+H)<sup>+</sup>. R<sub>f</sub>=0.56 in 50% acetone in hexanes.

#### Step F



2-(2-Fluoro-phenoxy)-2-methyl-3-{4-[2-(3-methyl-2-oxo-imidazolidin-4-yl)-ethoxy]-phenyl}-propionic acid ethyl ester: Trifluoroacetic acid (70mL) is added dropwise to a solution of triethylsilane (2.5mL, 15.73mmol,  $d=0.728$ ) and  
5 2-(2-Fluoro-phenoxy)-3-(4-{2-[1-(4-methoxy-benzyl)-3-methyl-2-oxo-imidazolidin-4-yl]-ethoxy}-phenyl)-2-methyl-propionic acid ethyl ester (4.44g, 7.86mmol). The reaction mixture is stirred 2h, then concentrated *in vacuo*. The residue is diluted with ethyl acetate, then washed with a saturated  
10 solution of aqueous sodium carbonate, water, and brine. The organic layer is dried and concentrated to provide the title compound (3.5g, 100%).  $^1\text{H}$  NMR (400MHz,  $\text{CDCl}_3$ ):  $\delta$  7.20 (d, 2H,  $J = 8.7$  Hz), 7.06-7.01 (m, 1H), 6.97-6.87 (m, 3H), 6.79 (d, 2H,  $J = 8.7$  Hz), 5.04-4.98 (m, 1H), 4.20 (q, 2H,  $J = 6.8$   
15 Hz), 4.02 (t, 2H,  $J = 6.8$  Hz), 3.76-3.69 (m, 1H), 3.55 (t, 1H,  $J = 8.7$  Hz), 3.26, 3.14 (ABq, 2H,  $J = 14.5$  Hz), 3.21 (t, 1H,  $J = 8.7$  Hz), 2.78 (s, 3H), 2.27-2.20 (m, 1H), 1.99-1.90 (m, 1H), 1.38 (s, 3H), 1.23 (t, 3H,  $J = 6.8$  Hz), 0.96 (t, 2H,  $J = 7.7$  Hz). MS [EI+] 445 (M+H) $^+$ .  $R_f=0.50$  in 10%  
20 methanol in methylene chloride.


#### Step G



3-{4-[2-(1-Benzyl-3-methyl-2-oxo-imidazolidin-4-yl)-ethoxy]-  
25 phenyl}-2-(2-fluoro-phenoxy)-2-methyl-propionic acid ethyl

**ester:** Benzyl bromide (0.02mL, 0.172mmol, d=1.438) and tetrabutyl ammonium iodide (catalytic amount) are added to a 0°C suspension of 2-(2-Fluoro-phenoxy)-2-methyl-3-{4-[2-(3-methyl-2-oxo-imidazolidin-4-yl)-ethoxy]-phenyl}-propionic acid ethyl ester (0.051g, 0.115mmol) and sodium hydride (0.011g, 0.287mmol, 60% suspension on mineral oil), pre-stirred for 1 h at ambient temperature. The reaction mixture is stirred at ambient temperature for 18h, diluted with ethyl acetate, and washed with 1N HCl, water, and brine. The organic layer is dried and concentrated *in vacuo* to provide the title compound, which is used in the next step. MS [EI+] 535 (M+H)<sup>+</sup>.

#### Step H



15

**3-{4-[2-(1-Benzyl-3-methyl-2-oxo-imidazolidin-4-yl)-ethoxy]-phenyl}-2-(2-fluoro-phenoxy)-2-methyl-propionic acid:** A solution of 3-{4-[2-(1-Benzyl-3-methyl-2-oxo-imidazolidin-4-yl)-ethoxy]-phenyl}-2-(2-fluoro-phenoxy)-2-methyl-propionic acid ethyl ester and 5N NaOH (0.2mL) in ethanol (2mL) is refluxed under nitrogen for 1h, cooled to ambient temperature, and concentrated *in vacuo*. The residue is diluted with 1N HCl, extracted with CH<sub>2</sub>Cl<sub>2</sub>, dried, concentrated *in vacuo*, and purified by LCMS to provide the title compound. <sup>1</sup>H NMR (400MHz, CDCl<sub>3</sub>): δ 7.55 (d, 1H, J =


25



8.4 Hz), 7.34-7.21 (m, 6H), 7.10-7.01 (m, 4H), 6.74 (d, 2H,  $J = 8.4$  Hz), 4.46-3.36 (m, 2H), 3.96 (q, 2H,  $J = 6.0$  Hz), 3.74-3.65 (m, 1H), 3.45-3.39 (m, 1H), 3.28, 3.16 (AB<sub>q</sub>, 2H,  $J = 14.7$  Hz), 3.05 (q, 1H;  $J = 8.6$  Hz), 2.26-2.19 (m, 1H), 1.93-1.86 (m, 1H), 1.41 (s, 3H). MS [EI+] 507 (M+H)<sup>+</sup>.

### Example 28

3-(4-{2-[1-(4-Chloro-benzyl)-3-methyl-2-oxo-imidazolidin-4-yl]-ethoxy}-phenyl)-2-(2-fluoro-phenoxy)-2-methyl-propionic acid




4-Chlorobenzyl bromide (0.037g, 0.179mmol) and tetrabutyl ammonium iodide (catalytic amount) are added to a 0°C suspension of 2-(2-Fluoro-phenoxy)-2-methyl-3-{4-[2-(3-methyl-2-oxo-imidazolidin-4-yl)-ethoxy]-phenyl}-propionic acid ethyl ester (0.053g, 0.119mmol) and sodium hydride (0.012g, 0.298mmol, 60% suspension on mineral oil), pre-stirred for 1 h at ambient temperature. The reaction mixture is stirred at ambient temperature for 18h, diluted with ethyl acetate, and washed. The organic layer is dried, concentrated, and purified by LCMS to provide the title compound. <sup>1</sup>H NMR (400MHz, CDCl<sub>3</sub>): δ 7.31-7.21 (m, 4H), 7.16 (d, 2H, *J* = 8.3 Hz), 7.11-7.00 (m, 4H), 6.74 (d, 2H, *J* = 8.3 Hz), 4.33, 4.31 (AB<sub>q</sub>, 2H, *J* = 15.5 Hz), 3.96 (t, 2H, *J* = 6.0

Hz), 3.71-3.64 (m, 1H), 3.39 (t, 1H,  $J = 8.3$  Hz), 3.29, 3.17 (AB<sub>q</sub>, 2H,  $J = 14.3$  Hz), 3.01 (t, 1H,  $J = 8.3$  Hz), 2.84 (s, 3H), 2.25-2.20 (m, 1H), 1.93-1.84 (m, 1H), 1.41 (s, 3H). MS [ES+]  $m/z$  exact mass calcd for C<sub>29</sub>H<sub>31</sub>N<sub>2</sub>O<sub>5</sub> 541.1906, found 541.1913.

### Example 29

2-(2-Fluoro-phenoxy)-2-methyl-3-(4-{2-[3-methyl-2-oxo-1-(4-trifluoromethyl-benzyl)-imidazolidin-4-yl]-ethoxy}-phenyl)-propionic acid




4-Trifluoromethylbenzyl bromide (0.03mL, 0.169mmol,  $d=1.546$ ) and tetrabutyl ammonium iodide (catalytic amount) are added to a 0°C suspension of 2-(2-Fluoro-phenoxy)-2-methyl-3-{4-[2-(3-methyl-2-oxo-imidazolidin-4-yl)-ethoxy]-phenyl}-propionic acid ethyl ester (0.050g, 0.112mmol) and sodium hydride (0.011g, 0.281mmol, 60% suspension on mineral oil), and pre-stirred for 1 h at ambient temperature. The reaction mixture is stirred at ambient temperature for 18h, diluted with ethyl acetate, and washed with 1N HCl, water, and brine. The organic layer is dried, concentrated, and purified by LCMS. <sup>1</sup>H NMR (400MHz, CDCl<sub>3</sub>):  $\delta$  7.54 (d, 1H,  $J = 7.8$  Hz), 7.35 (d, 1H,  $J = 7.8$  Hz), 7.25-7.21 (m, 3H), 7.16 (d, 1H,  $J = 7.8$  Hz), 7.08-6.95 (m, 4H), 6.73 (d, 2H,  $J = 7.8$

Hz), 4.45, 4.37 (AB<sub>q</sub>, 2H, *J* = 15.0 Hz), 3.95 (t, 2H, *J* = 5.9 Hz), 3.68-3.60 (m, 1H), 3.38 (t, 1H, *J* = 8.5 Hz), 3.29, 3.15 (AB<sub>q</sub>, 2H, *J* = 13.7 Hz), 3.02-2.97 (m, 1H), 2.84 (s, 3H), 2.25-2.19 (m, 1H), 1.92-1.83 (m, 1H), 1.39 (s, 3H). MS [ES<sup>+</sup>] *m/z* exact mass calcd for C<sub>30</sub>H<sub>31</sub>N<sub>2</sub>O<sub>5</sub>F<sub>4</sub> 575.2169, found 575.2172.

### Example 30

2-(2-Fluoro-phenoxy)-2-methyl-3-(4-{2-[3-methyl-2-oxo-1-(3-trifluoromethyl-benzyl)-imidazolidin-4-yl]-ethoxy}-phenyl)-propionic acid


#### Step A



2-(2-Fluoro-phenoxy)-2-methyl-3-(4-{2-[3-methyl-2-oxo-1-(3-trifluoromethyl-benzyl)-imidazolidin-4-yl]-ethoxy}-phenyl)-propionic acid ethyl ester : 3-Trifluoromethylbenzyl bromide (0.03mL, 0.175mmol, d=1.565) and tetrabutyl ammonium iodide (catalytic amount) are added to a 0°C suspension of 2-(2-Fluoro-phenoxy)-2-methyl-3-{4-[2-(3-methyl-2-oxo-imidazolidin-4-yl)-ethoxy]-phenyl}-propionic acid ethyl ester (0.052g, 0.117mmol) and sodium hydride (0.012g, 0.292mmol, 60% suspension on mineral oil), and pre-stirred for 1 h at ambient temperature. The reaction mixture is stirred at ambient temperature for 18h, diluted with ethyl

acetate, and washed with 1N HCl, water, and brine. The organic layer is dried, concentrated, and carried into the next step. MS [EI+] 603 (M+H)<sup>+</sup>.

5

Step B


**2-(2-Fluoro-phenoxy)-2-methyl-3-(4-{2-[3-methyl-2-oxo-1-(3-trifluoromethyl-benzyl)-imidazolidin-4-yl]-ethoxy}-phenyl)-propionic acid**

10 **propionic acid** : A solution of 2-(2-Fluoro-phenoxy)-2-methyl-3-(4-{2-[3-methyl-2-oxo-1-(3-trifluoromethyl-benzyl)-imidazolidin-4-yl]-ethoxy}-phenyl)-propionic acid ethyl ester and 5N NaOH (0.2mL) in ethanol (2mL) is refluxed under nitrogen for 1h, cooled to ambient temperature, and  
15 concentrated in vacuo. The residue is diluted with 1N HCl, extracted with CH<sub>2</sub>Cl<sub>2</sub>, dried, concentrated in vacuo, and purified by LCMS to provide the title compound. <sup>1</sup>H NMR (400MHz, CDCl<sub>3</sub>): δ 7.53-7.42 (m, 2H), 7.29-6.93 (m, 7H), 6.85-6.73 (m, 3H), 4.49-4.29 (m, 2H), 3.99-3.95 (m, 2H),  
20 3.81-3.68 (m, 1H), 3.46-3.41 (m, 1H), 3.28, 3.16 (AB<sub>q</sub>, 2H, J = 13.7 Hz), 3.09-3.04 (m, 1H), 2.86 (d, 3H, J = 5.6 Hz), 2.25-2.21 (m, 1H), 1.93-1.87 (m, 1H), 1.40 (s, 3H). MS [ES+] m/z exact mass calcd for C<sub>30</sub>H<sub>31</sub>N<sub>2</sub>O<sub>5</sub>F<sub>4</sub> 575.2169, found 575.2156.

25

## Example 31


2-(2-Fluoro-phenoxy)-3-(4-{2-[1-(3-methoxy-benzyl)-3-methyl-  
2-oxo-imidazolidin-4-yl]-ethoxy}-phenyl)-2-methyl-propionic  
acid

Step A

2-(2-Fluoro-phenoxy)-3-(4-{2-[1-(3-methoxy-benzyl)-3-methyl-  
2-oxo-imidazolidin-4-yl]-ethoxy}-phenyl)-2-methyl-propionic  
acid ethyl ester: 3-Methoxybenzyl bromide (0.02mL,  
0.172mmol, d=1.436) and tetrabutyl ammonium iodide  
(catalytic amount) are added to a 0°C suspension of 2-(2-  
Fluoro-phenoxy)-2-methyl-3-{4-[2-(3-methyl-2-oxo-  
imidazolidin-4-yl)-ethoxy]-phenyl}-propionic acid ethyl  
ester (0.051g, 0.115mmol) and sodium hydride (0.011g,  
0.287mmol, 60% suspension on mineral oil), and pre-stirred  
for 1 h at ambient temperature. The reaction mixture is

stirred at ambient temperature for 18h, diluted with ethyl acetate, and washed with 1N HCl, water, and brine. The organic layer is dried over MgSO<sub>4</sub>, concentrated *in vacuo*, and carried into the next step. MS [EI<sup>+</sup>] 565 (M+H)<sup>+</sup>.

5

Step B

2-(2-Fluoro-phenoxy)-3-(4-{2-[1-(3-methoxy-benzyl)-3-methyl-2-oxo-imidazolidin-4-yl]-ethoxy}-phenyl)-2-methyl-propionic


10 acid ethyl ester : A solution of 2-(2-Fluoro-phenoxy)-3-(4-{2-[1-(3-methoxy-benzyl)-3-methyl-2-oxo-imidazolidin-4-yl]-ethoxy}-phenyl)-2-methyl-propionic acid ethyl ester and 5N NaOH (0.2mL) in ethanol (2mL) is refluxed under nitrogen for 1h, cooled to ambient temperature, and concentrated. The

15 residue is diluted with 1N HCl, extracted with CH<sub>2</sub>Cl<sub>2</sub>, dried, concentrated, and purified by LCMS to provide the title compound. <sup>1</sup>H NMR (400MHz, CDCl<sub>3</sub>): δ 7.33-7.20 (m, 6H), 7.10-7.01 (m, 3H), 6.81-6.73 (m, 3H), 4.36, 4.30 (AB<sub>q</sub>, 2H, *J* = 14.9 Hz), 3.95 (t, 2H, *J* = 6.5 Hz), 3.71-3.68 (m, 1H),

20 3.45-3.40 (m, 1H), 3.29, 3.17 (AB<sub>q</sub>, 2H, *J* = 14.0 Hz), 3.08-3.03 (m, 1H), 2.85 (s, 3H), 2.24-2.20 (m, 1H), 1.94-1.85 (m, 1H), 1.41 (s, 3H). MS [ES<sup>+</sup>] *m/z* exact mass calcd for C<sub>30</sub>H<sub>34</sub>N<sub>2</sub>O<sub>6</sub> 537.2401, found 537.2418.

## Example 32

**3-{4-[2-(1-Benzyl-3-methyl-2-oxo-imidazolidin-4-yl)-ethoxy]-phenyl}-2-(2-fluoro-phenoxy)-2-methyl-propionic acid**


Step A

5  
**2-(2-Fluoro-phenoxy)-3-(4-{2-[1-(4-methoxy-benzyl)-3-methyl-2-oxo-imidazolidin-4-yl]-ethoxy}-phenyl)-2-methyl-propionic acid ethyl ester** Cesium carbonate (6.12g, 18.76mmol) is added to a solution of 2-(2-Fluoro-phenoxy)-3-(4-hydroxy-phenyl)-2-methyl-propionic acid ethyl ester 1.99g, 6.26mmol)  
 10 and toluene-4-sulfonic acid 2-[1-(4-methoxy-benzyl)-3-methyl-2-oxo-imidazolidin-4-yl]-ethyl ester (2.88g, 6.88mmol) in DMF (20mL). The resultant mixture is stirred at 65°C under an atmosphere of nitrogen for 18 h, then  
 15 diluted with ethyl acetate. The organic layer is washed with 1N HCl, water, and brine. The organic layer is dried over MgSO<sub>4</sub>, concentrated in vacuo, and purified by flash column chromatography (17% acetone in hexanes) to provide the title compound (3.4g, 97%). <sup>1</sup>H NMR (400MHz, CDCl<sub>3</sub>): δ  
 20 7.19 (d, 2H, J = 5.4 Hz), 7.17 (d, 2H, J = 5.4 Hz), 7.07-7.01 (m, 1H), 6.97-6.87 (m, 3H), 6.83 (d, 2H, J = 8.6 Hz), 6.74 (d, 2H, J = 8.6 Hz), 4.32, 4.28 (AB<sub>q</sub>, 2H, J = 14.6 Hz), 4.20 (q, 2H, J = 7.5 Hz), 3.94 (t, 2H, J = 5.9 Hz), 3.77 (s, 3H), 3.60-3.53 (m, 1H), 3.33 (t, 1H, J = 8.6 Hz), 3.26, 3.14  
 25 (AB<sub>q</sub>, 2H, J = 14.0 Hz), 2.92 (t, 1h, J = 8.6 Hz), 2.82 (s, 3H), 2.24-2.17 (m, 1H), 1.89-1.82 (m, 1H), 1.38 (s, 3H),

1.23 (t, 3H,  $J = 7.5$  Hz). MS [EI+] 565 (M+H)<sup>+</sup>.  $R_f=0.48$  in 50% acetone in hexanes.


Step B

5




**2-(2-Fluoro-phenoxy)-2-methyl-3-{4-[2-(3-methyl-2-oxo-imidazolidin-4-yl)-ethoxy]-phenyl}-propionic acid ethyl ester:** Trifluoroacetic acid (55mL) is added dropwise to a solution of triethylsilane (1.95mL, 12.18mmol,  $d=0.728$ ) and  
10 2-(2-Fluoro-phenoxy)-3-(4-{2-[1-(4-methoxy-benzyl)-3-methyl-2-oxo-imidazolidin-4-yl]-ethoxy}-phenyl)-2-methyl-propionic acid ethyl ester (3.44g, 6.09mmol). The reaction mixture is stirred under nitrogen at ambient temperature for 2h, then concentrated. The residue is diluted with ethyl  
15 acetate, then washed. The organic layer is dried and concentrated *in vacuo* to provide the title compound (2.7g, 78%). <sup>1</sup>H NMR (400MHz, CDCl<sub>3</sub>):  $\delta$  7.20 (d, 2H,  $J = 8.7$  Hz), 7.06-7.01 (m, 1H), 6.97-6.87 (m, 3H), 6.79 (d, 2H,  $J = 8.7$  Hz), 5.04-4.98 (m, 1H), 4.20 (q, 2H,  $J = 6.8$  Hz), 4.02 (t, 2H,  $J = 6.8$  Hz), 3.76-3.69 (m, 1H), 3.55 (t, 1H,  $J = 8.7$  Hz), 3.26, 3.14 (AB<sub>q</sub>, 2H,  $J = 14.5$  Hz), 3.21 (t, 1H,  $J = 8.7$  Hz), 2.78 (s, 3H), 2.27-2.20 (m, 1H), 1.99-1.90 (m, 1H), 1.38 (s, 3H), 1.23 (t, 3H,  $J = 6.8$  Hz), 0.96 (t, 2H,  $J = 7.7$  Hz). MS [EI+] 445 (M+H)<sup>+</sup>.  
20  
25  $R_f=0.38$  in 10% methanol in methylene chloride.



Step C


- 3-{4-[2-(1-Benzyl-3-methyl-2-oxo-imidazolidin-4-yl)-ethoxy]-phenyl}-2-(2-fluoro-phenoxy)-2-methyl-propionic acid ethyl ester:** Benzyl bromide (0.02mL, 0.179mmol, d=1.438) and tetrabutyl ammonium iodide (catalytic amount) are added to a 0°C suspension of 2-(2-Fluoro-phenoxy)-2-methyl-3-{4-[2-(3-methyl-2-oxo-imidazolidin-4-yl)-ethoxy]-phenyl}-propionic acid ethyl ester (0.053g, 0.119mmol) and sodium hydride (0.012g, 0.298mmol, 60% suspension on mineral oil), pre-stirred for 1 h at ambient temperature. The reaction mixture is stirred at ambient temperature for 18h, diluted with ethyl acetate, and washed with 1N HCl, water, and brine. The organic layer is dried over MgSO<sub>4</sub> and concentrated in vacuo to provide the title compound, which is carried into the next step. MS [EI<sup>+</sup>] 535 (M+H)<sup>+</sup>.

Step H

3-{4-[2-(1-Benzyl-3-methyl-2-oxo-imidazolidin-4-yl)-ethoxy]-phenyl}-2-(2-fluoro-phenoxy)-2-methyl-propionic acid: A solution of 3-{4-[2-(1-Benzyl-3-methyl-2-oxo-imidazolidin-4-yl)-ethoxy]-phenyl}-2-(2-fluoro-phenoxy)-2-methyl-propionic acid ethyl ester and 5N NaOH (0.2mL) in ethanol (2mL) is refluxed under nitrogen for 1h, cooled to ambient temperature, and concentrated *in vacuo*. The residue is diluted with 1N HCl, extracted with CH<sub>2</sub>Cl<sub>2</sub>, dried, concentrated, and purified by LCMS to provide the title compound. <sup>1</sup>H NMR (400MHz, CDCl<sub>3</sub>): δ 7.55 (d, 1H, *J* = 8.4 Hz), 7.34-7.21 (m, 6H), 7.10-7.01 (m, 4H), 6.74 (d, 2H, *J* = 8.4 Hz), 4.46-3.36 (m, 2H), 3.96 (q, 2H, *J* = 6.0 Hz), 3.74-3.65 (m, 1H), 3.45-3.39 (m, 1H), 3.28, 3.16 (AB<sub>q</sub>, 2H, *J* = 14.7 Hz), 3.05 (q, 1H, *J* = 8.6 Hz), 2.26-2.19 (m, 1H), 1.93-1.86 (m, 1H), 1.41 (s, 3H). . MS [ES+] *m/z* exact mass calcd for C<sub>29</sub>H<sub>32</sub>N<sub>2</sub>O<sub>5</sub>F 507.2295, found 507.2308.

### Example 33

3-(4-{2-[1-(4-Chloro-benzyl)-3-methyl-2-oxo-imidazolidin-4-yl]-ethoxy}-phenyl)-2-(2-fluoro-phenoxy)-2-methyl-propionic acid




4-Chlorobenzyl bromide (0.035g, 0.172mmol) and tetrabutyl ammonium iodide (catalytic amount) are added to a 0°C suspension of 2-(2-Fluoro-phenoxy)-2-methyl-3-{4-[2-(3-


methyl-2-oxo-imidazolidin-4-yl)-ethoxy]-phenyl)-propionic acid ethyl ester (0.051g, 0.115mmol) and sodium hydride (0.011g, 0.287mmol, 60% suspension on mineral oil), pre-stirred for 1 h at ambient temperature. The reaction mixture is stirred at ambient temperature for 18h, diluted with ethyl acetate, and washed. The organic layer is dried, concentrated, and purified by LCMS to provide the title compound. <sup>1</sup>H NMR (400MHz, CDCl<sub>3</sub>): δ 7.31-7.22 (m, 5H), 7.15-7.01 (m, 5H), 6.74 (d, 2H, J = 7.8 Hz), 4.34 (d, 2H, J = 14.3 Hz), 3.96 (s, 2H), 3.73-3.71 (m, 1H), 3.46-3.40 (m, 1H), 3.28, 3.16 (AB<sub>q</sub>, 2H, J = 14.3 Hz), 3.08-3.06 (m, 1H), 2.85 (s, 3H), 2.23-2.20 (m, 1H), 1.91-1.89 (m, 1H), 1.41 (s, 3H). MS [ES+] m/z exact mass calcd for C<sub>29</sub>H<sub>31</sub>N<sub>2</sub>O<sub>5</sub>FCl 541.1906, found 541.1909.

#### Example 34

2-(2-Fluoro-phenoxy)-2-methyl-3-(4-{2-[3-methyl-2-oxo-1-(4-trifluoromethyl-benzyl)-imidazolidin-4-yl]-ethoxy}-phenyl)-propionic acid




Step A



**2-(2-Fluoro-phenoxy)-2-methyl-3-(4-{2-[3-methyl-2-oxo-1-(4-trifluoromethyl-benzyl)-imidazolidin-4-yl]-ethoxy}-phenyl)-propionic acid ethyl ester :** 4-Trifluoromethylbenzyl

5 bromide (0.03mL, 0.169mmol, d=1.546) and tetrabutyl ammonium iodide (catalytic amount) are added to a 0°C suspension of 2-(2-Fluoro-phenoxy)-2-methyl-3-{4-[2-(3-methyl-2-oxo-imidazolidin-4-yl)-ethoxy]-phenyl}-propionic acid ethyl ester (0.050g, 0.113mmol) and sodium hydride (0.011g, 10 0.281mmol, 60% suspension on mineral oil), and pre-stirred for 1 h at ambient temperature. The reaction mixture is stirred at ambient temperature for 18h, diluted with ethyl acetate, and washed with 1N HCl, water, and brine. The organic layer is dried, concentrated, and carried into the 15 next step. MS [EI+] 603 (M+H)<sup>+</sup>.

#### Step B




2-(2-Fluoro-phenoxy)-2-methyl-3-(4-{2-[3-methyl-2-oxo-1-(4-trifluoromethyl-benzyl)-imidazolidin-4-yl]-ethoxy}-phenyl)-propionic acid : A solution of 2-(2-Fluoro-phenoxy)-2-methyl-3-(4-{2-[3-methyl-2-oxo-1-(4-trifluoromethyl-benzyl)-imidazolidin-4-yl]-ethoxy}-phenyl)-propionic acid ethyl ester and 5N NaOH (0.2mL) in ethanol (2mL) is refluxed under nitrogen for 1h, cooled to ambient temperature, and concentrated *in vacuo*. The residue is diluted with 1N HCl, extracted with CH<sub>2</sub>Cl<sub>2</sub>, dried, concentrated, and purified by LCMS to provide the title compound. <sup>1</sup>H NMR (400MHz, CDCl<sub>3</sub>): δ 7.56-7.51 (m 1H), 7.48-7.33 (m, 2H), 7.26-7.21 (m, 3H), 7.11-6.98 (m, 4H), 6.74 (d, 2H, *J* = 8.4 Hz), 4.49-4.34 (m, 2H), 3.98 (t, 2H, *J* = 5.3 Hz), 3.74-3.67 (m, 1H), 3.42 (t, 1H, *J* = 8.4 Hz), 3.35, 3.23 (AB<sub>q</sub>, 2H, *J* = 14.5 Hz), 3.08-3.03 (m, 1H), 2.86 (s, 3H), 2.27-2.20 (m, 1H), 1.95-1.88 (m, 1H), 1.42 (s, 3H). MS [ES<sup>+</sup>] *m/z* exact mass calc'd for C<sub>30</sub>H<sub>31</sub>N<sub>2</sub>O<sub>5</sub>F<sub>4</sub> 575.2169, found 575.2175.

20


25

**Example 35**

**2-(2-Fluoro-phenoxy)-2-methyl-3-(4-{2-[3-methyl-2-oxo-1-(3-trifluoromethyl-benzyl)-imidazolidin-4-yl]-ethoxy}-phenyl)-propionic acid**

Step A


**2-(2-Fluoro-phenoxy)-2-methyl-3-(4-{2-[3-methyl-2-oxo-1-(3-trifluoromethyl-benzyl)-imidazolidin-4-yl]-ethoxy}-phenyl)-propionic acid ethyl ester** : 3-Trifluoromethylbenzyl bromide (0.03mL, 0.165mmol, d=1.565) and tetrabutyl ammonium iodide (catalytic amount) are added to a 0°C suspension of 2-(2-Fluoro-phenoxy)-2-methyl-3-{4-[2-(3-methyl-2-oxo-imidazolidin-4-yl)-ethoxy]-phenyl}-propionic acid ethyl ester (0.049g, 0.110mmol) and sodium hydride (0.011g, 0.276mmol, 60% suspension on mineral oil), pre-stirred for 1 h at ambient temperature. The reaction mixture is stirred at ambient temperature for 18h, diluted with ethyl acetate, and washed. The organic layer is dried, concentrated, and carried into the next step. MS [EI+] 603 (M+H)<sup>+</sup>.

Step B

2-(2-Fluoro-phenoxy)-2-methyl-3-(4-{2-[3-methyl-2-oxo-1-(3-  
 5 trifluoromethyl-benzyl)-imidazolidin-4-yl]-ethoxy}-phenyl)-  
 propionic acid ethyl ester : A solution of 2-(2-Fluoro-  
 phenoxy)-2-methyl-3-(4-{2-[3-methyl-2-oxo-1-(3-  
 trifluoromethyl-benzyl)-imidazolidin-4-yl]-ethoxy}-phenyl)-  
 propionic acid ethyl ester and 5N NaOH (0.2mL) in ethanol  
 10 (2mL) is refluxed under nitrogen for 1h, cooled to ambient  
 temperature, and concentrated in vacuo. The residue is  
 diluted with 1N HCl, extracted with CH<sub>2</sub>Cl<sub>2</sub>, dried,  
 concentrated, and purified by LCMS to provide the title  
 compound. <sup>1</sup>H NMR (400MHz, CDCl<sub>3</sub>): δ 7.51-7.41 (m, 2H), 7.28-  
 15 7.19 (m, 3H), 7.09-6.99 (m, 4H), 6.81-6.73 (m, 3H), 4.48-  
 4.28 (m, 2H), 3.98-3.94 (m, 2H), 3.73-3.65 (m, 1H), 3.42 (t,  
 1H, J = 8.7 Hz), 3.29, 3.15 (AB<sub>q</sub>, 2H, J = 13.6 Hz), 3.06-  
 3.01 (m, 1H), 2.85 (d, 3H, J = 5.6 Hz), 2.25-2.21 (m, 1H),  
 1.93-1.87 (m, 1H), 1.40 (s, 3H). MS [ES+] m/z exact mass  
 20 calcd for C<sub>30</sub>H<sub>31</sub>N<sub>2</sub>O<sub>5</sub>F<sub>4</sub> 575.2169, found 575.2172.

**Example 36**

2-(2-Fluoro-phenoxy)-3-(4-{2-[1-(3-methoxy-benzyl)-3-methyl-  
 2-oxo-imidazolidin-4-yl]-ethoxy}-phenyl) -2-methyl-propionic  
 25 acid ethyl ester

Step A


5    **2-(2-Fluoro-phenoxy)-3-(4-{2-[1-(3-methoxy-benzyl)-3-methyl-2-oxo-imidazolidin-4-yl]-ethoxy}-phenyl)-2-methyl-propionic acid ethyl ester:** 3-Methoxybenzyl bromide (0.03mL, 0.179mmol, d=1.436) and tetrabutyl ammonium iodide (catalytic amount) are added to a 0°C suspension of 2-(2-Fluoro-phenoxy)-2-methyl-3-{4-[2-(3-methyl-2-oxo-imidazolidin-4-yl)-ethoxy]-phenyl}-propionic acid ethyl ester (0.053g, 0.119mmol) and sodium hydride (0.012g, 0.298mmol, 60% suspension on mineral oil), and pre-stirred for 1 h at ambient temperature. The reaction mixture is stirred at ambient temperature for 18h, diluted with ethyl acetate, and washed. The organic layer is dried, concentrated, and carried into the next step. MS [EI+] 565 (M+H)<sup>+</sup>.

10

15

Step B






2-(2-Fluoro-phenoxy)-3-(4-{2-[1-(3-methoxy-benzyl)-3-methyl-2-oxo-imidazolidin-4-yl]-ethoxy}-phenyl)-2-methyl-propionic acid ethyl ester : A solution of 2-(2-Fluoro-phenoxy)-3-(4-{2-[1-(3-methoxy-benzyl)-3-methyl-2-oxo-imidazolidin-4-yl]-ethoxy}-phenyl)-2-methyl-propionic acid ethyl ester and 5N NaOH (0.2mL) in ethanol (2mL) is refluxed under nitrogen for 1h, cooled to ambient temperature, and concentrated in vacuo. The residue is diluted with 1N HCl, extracted with CH<sub>2</sub>Cl<sub>2</sub>, dried through a Varian ChemElut cartridge, concentrated in vacuo, and purified by LCMS to provide the title compound. <sup>1</sup>H NMR (400MHz, CDCl<sub>3</sub>): δ 7.26-7.19 (m, 3H), 7.08-6.99 (m, 4H), 6.83-6.78 (m, 3H), 6.74 (d, 2H, *J* = 8.4 Hz), 4.36, 4.30 (AB<sub>q</sub>, 2H, *J* = 14.9 Hz), 3.95 (t, 2H, *J* = 6.5 Hz), 3.77 (s, 3H), 3.65-3.58 (m, 1H), 3.37 (t, 1H, *J* = 8.8 Hz), 3.28, 3.16 (AB<sub>q</sub>, 2H, *J* = 14.4 Hz), 2.97 (t, 1H, *J* = 8.8 Hz), 2.83 (s, 3H), 2.25-2.17 (m, 1H), 1.92-1.83 (m, 1H), 1.40 (s, 3H). MS [ES+] *m/z* exact mass calcd for C<sub>30</sub>H<sub>34</sub>N<sub>2</sub>O<sub>6</sub>F 537.2401, found 537.2413.

#### Example 37

3-{4-[2-(1-Benzyl-3-methyl-2-oxo-imidazolidin-4-yl)-ethoxy]-phenyl}-2-butoxy-2-methyl-propionic acid

Step A**3-(4-Benzyloxy-phenyl)-2-methyl-acrylic acid ethyl ester :**


Triethyl-2-phosphonopropionate (22.2mL, 103.65mmol, d=1.111)

5 is added to a 0°C slurry of sodium hydride (4.15g, 103.65mmol, 60% dispersion on mineral oil) in anhydrous THF (150mL), then allowed to warm to ambient temperature over thirty minutes. The slurry is re-cooled to 0°C and 4-benzyloxybenzaldehyde (20g, 94.23mmol) is added in one

10 portion as a solid. The reaction mixture is stirred 18h at ambient temperature, then diluted with diethyl ether. The organic layer is washed with a saturated solution of aqueous ammonium chloride, water, and brine, then dried over MgSO<sub>4</sub> and concentrated *in vacuo*. The residue is purified by flash

15 chromatography to provide the title compound (24.3g, 87%).  
<sup>1</sup>H NMR (400MHz, CDCl<sub>3</sub>): δ 7.67 (s, 1H), 7.46-7.35 (m, 6H), 7.01 (d, 2H, *J* = 8.1 Hz), 5.10 (s, 2H), 4.28 (q, 2H, *J* = 7.2 Hz), 2.15 (s, 3H), 1.36 (t, 3H, *J* = 7.2 Hz). R<sub>f</sub>=0.31 in 20% acetone in hexanes.


20

Step B

**3-(4-Benzyloxy-phenyl)-2,3-dihydroxy-2-methyl-propionic acid ethyl ester :** 3-(4-Benzyloxy-phenyl)-2-methyl-acrylic acid

ethyl ester (13.84g, 46.70mmol) is added to a solution of acetone (235mL), 4-methylmorpholine N-oxide (6.02g, 51.37mmol), tert-butyl alcohol (24mL), and 4% osmium tetroxide in water (14.7mL), then stirred 20h. The reaction is quenched with sodium hydrosulphite (4.7g, 26.81mmol), diluted with ethyl acetate, washed with water. The organic layer is dried, concentrated, and purified by flash chromatography to provide the title compound (12.9g, 83%).  
<sup>1</sup>H NMR (400MHz, CDCl<sub>3</sub>): δ 7.44-7.32 (m, 7H), 6.82 (d, 2H, *J* = 9.3 Hz), 5.06 (s, 2H), 4.78 (d, 1H, *J* = 6.0 Hz), 4.32-4.27 (m, 2H), 3.52 (s, 1H), 2.66 (d, 1H, *J* = 7.3 Hz), 1.33 (t, 3H, *J* = 7.3 Hz), 1.17 (s, 3H). *R*<sub>f</sub>=0.48 in 50% acetone in hexanes.

15


Step C

**3-(4-Benzyloxy-phenyl)-2-hydroxy-2-methyl-propionic acid ethyl ester** : Boron trifluoride etherate was added to a 0°C solution of 3-(4-Benzyloxy-phenyl)-2,3-dihydroxy-2-methyl-propionic acid ethyl ester (12.9g, 38.96mmol) and triethylsilane (28mL, 175.3mmol, d=0.728) in anhydrous CH<sub>2</sub>Cl<sub>2</sub>, then the reaction is allowed to gradually warm to ambient temperature over 3h. The reaction is quenched with a saturated aqueous solution of sodium bicarbonate and extracted with additional CH<sub>2</sub>Cl<sub>2</sub>. The organic layer is dried, concentrated, and separated by chiral chromatography to provide the title compound (11.7g, 96%).  
<sup>1</sup>H NMR (400MHz, CDCl<sub>3</sub>): δ 7.44-7.33 (m, 5H), 7.12 (d, 2H, *J* = 6.85 Hz), 6.90

(d, 2H,  $J = 6.85$  Hz), 5.04 (s, 2H), 4.22-4.14 (m, 2H), 3.03 (d, 1H,  $J = 13.69$  Hz), 2.87 (d, 1H,  $J = 13.69$  Hz), 1.48 (s, 3H), 1.27 (t, 3H,  $J = 7.34$  Hz).  $R_f=0.60$  in 50% acetone in hexanes.

5


#### Step D



10 **2-Hydroxy-3-(4-hydroxy-phenyl)-2-methyl-propionic acid ethyl ester** : 3-(4-Benzyloxy-phenyl)-2-hydroxy-2-methyl-propionic acid ethyl ester (5.9g, 18.77mmol) is dissolved in ethanol (375mL), treated with 10% palladium on carbon (2.95g), and stirred under an atmosphere of hydrogen for 2 h. The  
 15 suspension is filtered and concentrated to provide the title compound.  $^1\text{H}$  NMR (400MHz,  $\text{CDCl}_3$ ):  $\delta$  7.00 (d, 2H,  $J = 8.3$  Hz), 6.69 (d, 2H,  $J = 8.3$  Hz), 4.18-4.13 (m, 2H), 2.99, 2.83 (ABq, 2H,  $J = 13.7$  Hz), 1.47 (s, 3H), 1.24 (t, 3H,  $J = 7.34$  Hz).  $R_f=0.58$  in 50% acetone in hexanes.


20

#### Step E




**2-Hydroxy-3-(4-{2-[1-(4-methoxy-benzyl)-3-methyl-2-oxo-imidazolidin-4-yl]-ethoxy}-phenyl)-2-methyl-propionic acid ethyl ester:** Cesium carbonate (21.9g, 67.3mmol) is added to a solution of 2-hydroxy-3-(4-{2-[1-(4-methoxy-benzyl)-3-methyl-2-oxo-imidazolidin-4-yl]-ethoxy}-phenyl)-2-methyl-propionic acid ethyl ester (5.03g, 22.4mmol) and toluene-4-sulfonic acid 2-[1-(4-methoxy-benzyl)-3-methyl-2-oxo-imidazolidin-4-yl]-ethyl ester (10.33g, 24.7mmol) in DMF (100mL). The resultant mixture is stirred at 55°C under an atmosphere of nitrogen for 18 h, then diluted with ethyl acetate. The organic layer is washed. The organic layer is dried, concentrated, and purified by flash chromatography to provide the title compound (735g, 70%). <sup>1</sup>H NMR (400MHz, CDCl<sub>3</sub>): δ 7.16 (d, 2H, *J* = 8.7 Hz), 7.07 (d, 2H, *J* = 7.9 Hz), 6.82 (d, 2H, *J* = 7.9 Hz), 6.70 (d, 2H, *J* = 8.7 Hz), 4.31, 4.27 (AB<sub>q</sub>, 2H, *J* = 15.1 Hz), 4.19-4.11 (m, 2H), 3.91 (t, 2H, *J* = 7.2 Hz), 3.77 (s, 3H), 3.59-3.52 (m, 1H), 3.32 (t, 1H, *J* = 8.7 Hz), 3.01-2.83 (m, 6H), 2.81 (s, 3H), 2.23-2.15 (m, 1H), 1.88-1.80 (m, 1H), 1.45 (s, 3H), 1.26 (t, 6H, *J* = 6.65 Hz), 0.87 (t, 3H, *J* = 6.5 Hz). *R*<sub>f</sub>=0.41 in 50% acetone in hexanes.

#### Step F




**2-Butoxy-3-(4-{2-[1-(4-methoxy-benzyl)-3-methyl-2-oxo-imidazolidin-4-yl]-ethoxy}-phenyl)-2-methyl-propionic acid ethyl ester :** A slurry of 2-hydroxy-3-(4-{2-[1-(4-methoxy-benzyl)-3-methyl-2-oxo-imidazolidin-4-yl]-ethoxy}-phenyl)-2-methyl-propionic acid ethyl ester (7.35g, 15.62mmol) and sodium hydride (2.5g, 62.5mmol, 60% dispersion on mineral oil) in DMF (50mL) is stirred for 45 minutes, then cooled to 0° C. Iodobutane (36mL, 312mmol, d=1.617) and 18-crown-6 (16.5g, 62.5mmol) are added, then the reaction mixture is stirred 18h at ambient temperature and diluted with ethyl acetate. The organic layer is washed, then dried concentrated in vacuo, and diluted with ethanol (400mL) and 5N NaOH (40mL). The reaction mixture is refluxed for 1h, cooled to ambient temperature, and concentrated in vacuo. The residue is washed, dried, concentrated, and re-dissolved in ethanol (300mL). Concentrated H<sub>2</sub>SO<sub>4</sub> (30mL) is added to the reaction solution, then the mixture is refluxed 1h, cooled to ambient temperature, concentrated, and diluted with ethyl acetate. The organic layer is washed, dried, and concentrated to provide the title compound (5.3g, 65%). <sup>1</sup>H NMR (400MHz, CDCl<sub>3</sub>): δ 7.17 (d, 2H, J = 8.8 Hz), 7.09 (d, 2H, J = 8.8 Hz), 6.84 (d, 2H, J = 8.8 Hz), 6.70 (d, 2H, J = 8.8 Hz), 4.32, 4.28 (AB<sub>q</sub>, 2H, J = 14.7 Hz), 4.17-4.11 (m, 2H), 3.93 (t, 2H, J = 5.9 Hz), 3.78 (s, 3H), 3.61-3.53 (m, 3H), 3.40-3.31 (m, 3H), 2.97-2.89 (m, 3H), 2.97-2.89 (m, 3H), 2.81 (s, 3H), 2.25-2.15 (m, 1H), 1.89-1.81 (m, 1H), 1.59-1.49 (m, 2H), 1.42-1.33 (m, 2H), 1.29 (s, 3H), 1.24 (t, 3H, J = 7.2 Hz), 0.91 (t, 3H, J = 7.2 Hz). R<sub>f</sub>=0.43 in 50% acetone in hexanes.

Step G


**2-Butoxy-2-methyl-3-{4-[2-(3-methyl-2-oxo-imidazolidin-4-yl)-ethoxy]-phenyl}-propionic acid ethyl ester :**

Trifluoroacetic acid (110mL) is added dropwise to a solution of 2-Butoxy-3-(4-{2-[1-(4-methoxy-benzyl)-3-methyl-2-oxo-imidazolidin-4-yl]-ethoxy}-phenyl)-2-methyl-propionic acid ethyl ester (5.31g, 10.1mmol) and triethylsilane (3.22mL, 20.2mmol, d=0.728). The reaction mixture is stirred at ambient temperature for 4h, concentrated in vacuo, and diluted with ethyl acetate. The solution is washed, then dried, concentrated, and purified by flash chromatography to provide the title compound (4.1g, 100%). <sup>1</sup>H NMR (400MHz, CDCl<sub>3</sub>): δ 7.09 (d, 2H, J = 8.4 Hz), 6.74 (d, 2H, J = 8.4 Hz), 5.36 (s, 1H), 4.17-4.07 (m, 2H), 4.02-3.96 (m, 2H), 3.76-3.68 (m, 2H), 3.55 (t, 1H, J = 8.8 Hz), 3.40-3.31 (m, 2H), 3.20 (t, 1H, J = 8.0 Hz), 2.95, 2.91 (AB<sub>q</sub>, 2H, J = 13.6 Hz), 2.76 (s, 3H), 2.24-2.18 (m, 1H), 1.97-1.88 (m, 1H), 1.58-1.51 (m, 2H), 1.41-1.31 (m, 2H), 1.27 (s, 3H), 1.22 (t, 3H, J = 7.2 Hz), 0.89 (t, 3H, J = 7.2 Hz). R<sub>f</sub>=0.44 in 10% CH<sub>3</sub>OH in CH<sub>2</sub>Cl<sub>2</sub>.

Step H

**3-{4-[2-(1-Benzyl-3-methyl-2-oxo-imidazolidin-4-yl)-ethoxy]-phenyl}-2-butoxy-2-methyl-propionic acid ethyl ester :**

- 5 Benzyl bromide (0.0.21mL, 0.174mmol, d=1.438) and tetrabutyl ammonium iodide (catalytic amount) are added to a 0°C suspension of 2-butoxy-2-methyl-3-{4-[2-(3-methyl-2-oxo-imidazolidin-4-yl)-ethoxy]-phenyl}-propionic acid ethyl ester (0.047g, 0.116mmol) and sodium hydride (0.012g, 10 0.29mmol, 60% suspension on mineral oil), pre-stirred for 1 h at ambient temperature. The reaction mixture is stirred at ambient temperature for 48h, diluted with ethyl acetate, and washed with 1N HCl and water. The organic layer is dried, concentrated, and used in the next step. MS [EI+] 497  
15 (M+H)<sup>+</sup>.

Step I



3-{4-[2-(1-Benzyl-3-methyl-2-oxo-imidazolidin-4-yl)-ethoxy]-phenyl}-2-butoxy-2-methyl-propionic acid : A solution of 3-{4-[2-(1-Benzyl-3-methyl-2-oxo-imidazolidin-4-yl)-ethoxy]-phenyl}-2-butoxy-2-methyl-propionic acid ethyl ester and 5N NaOH (0.3mL) in ethanol (3mL) is refluxed under nitrogen for 1h, cooled to ambient temperature, and concentrated. The residue is diluted with 1N HCl, extracted with CH<sub>2</sub>Cl<sub>2</sub>, dried, concentrated, and purified by LCMS to provide the title compound. <sup>1</sup>H NMR (400MHz, CDCl<sub>3</sub>): δ 7.33-7.22 (m, 5H), 7.07 (d, 2H, J = 8.6 Hz), 6.71 (d, 2H, J = 8.6 Hz), 4.39, 4.35 (AB<sub>q</sub>, 2H, J = 15.3 Hz), 3.94 (t, 2H, J = 5.7 Hz), 3.69-3.62 (m, 1H), 3.58-3.52 (m, 1H), 3.50-3.44 (m, 1H), 3.40 (t, 1H, J = 9.1 Hz), 3.04-3.01 (m, 1H), 3.02, 2.94 (AB<sub>q</sub>, 2H, J = 13.9 Hz), 2.85 (s, 3H), 2.25-2.18 (m, 1H), 1.92-1.85 (m, 1H), 1.62-1.54 (m, 1H), 1.46 (s, 3H), 1.42-1.32 (m, 2H), 0.92 (t, 3H, J = 7.2 Hz). MS [EI+] 469 (M+H)<sup>+</sup>, [EI-] 467 (M-H)<sup>+</sup>.

### Example 38

2-Butoxy-3-(4-{2-[1-(3-methoxy-benzyl)-3-methyl-2-oxo-imidazolidin-4-yl]-ethoxy}-phenyl)-2-methyl-propionic acid

#### Step A

2-Butoxy-3-(4-{2-[1-(3-methoxy-benzyl)-3-methyl-2-oxo-imidazolidin-4-yl]-ethoxy}-phenyl)-2-methyl-propionic acid ethyl ester: 3-Methoxybenzyl bromide (0.024mL, 0.170mmol, d=1.436) and tetrabutyl ammonium iodide (catalytic amount) are added to a 0°C suspension of 2-butoxy-2-methyl-3-{4-[2-(3-methyl-2-oxo-imidazolidin-4-yl)-ethoxy]-phenyl}-propionic acid ethyl ester (0.046g, 0.114mmol) and sodium hydride (0.011g, 0.28mmol, 60% suspension on mineral oil), pre-stirred for 1 h at ambient temperature. The reaction

mixture is stirred at ambient temperature 48h, diluted with ethyl acetate, and washed with 1N HCl and water. The organic layer is dried, concentrated, and used in the next step. MS [EI+] 527 (M+H)<sup>+</sup>.

5

#### Step B


**2-Butoxy-3-(4-{2-[1-(3-methoxy-benzyl)-3-methyl-2-oxo-imidazolidin-4-yl]-ethoxy}-phenyl)-2-methyl-propionic acid :**

A solution of 2-Butoxy-3-(4-{2-[1-(3-methoxy-benzyl)-3-methyl-2-oxo-imidazolidin-4-yl]-ethoxy}-phenyl)-2-methyl-propionic acid ethyl ester and 5N NaOH (0.3mL) in ethanol (3mL) is refluxed under nitrogen for 1h, cooled to ambient temperature, and concentrated in vacuo. The residue is diluted with 1N HCl, extracted, dried, concentrated, and purified by LCMS to provide the title compound. <sup>1</sup>H NMR (400MHz, CDCl<sub>3</sub>): δ 7.22 (t, 1H, J = 8.0 Hz), 7.07 (d, 2H, J = 8.5 Hz), 6.83-6.79 (m, 3H), 6.72 (d, 2H, J = 8.5 Hz), 4.38, 4.30 (AB<sub>q</sub>, 2H, J = 15.1 Hz), 3.95 (t, 2H, J = 6.0 Hz), 3.78 (s, 3H), 3.69-3.62 (m, 1H), 3.60-3.54 (m, 1H), 3.50-3.45 (m, 1H), 3.50-3.45 (m, 1H), 3.40 (t, 1H, J = 9.0 Hz), 3.03-3.00 (m, 1H), 3.01, 2.95 (AB<sub>q</sub>, 2H, J = 14.6 Hz), 2.85 (s, 3H), 2.26-2.19 (m, 1H), 1.92-1.85 (m, 1H), 1.62-1.55 (m, 2H), 1.48 (s, 3H), 1.42 (m, 2H), 0.93 (t, 3H, J = 7.0 Hz). MS [EI+] 499 (M+H)<sup>+</sup>, [EI-] 497 (M-H)<sup>+</sup>.

25

#### **Example 39**

**2-Butoxy-3-(4-{2-[1-(4-chloro-benzyl)-3-methyl-2-oxo-imidazolidin-4-yl]-ethoxy}-phenyl)-2-methyl-propionic acid**




4-Phenylbenzyl chloride (0.031g, 0.152mmol) and  
tetrabutyl ammonium iodide (catalytic amount) are added to a  
5 0°C suspension of 2-butoxy-2-methyl-3-{4-[2-(3-methyl-2-oxo-  
imidazolidin-4-yl)-ethoxy]-phenyl}-propionic acid ethyl  
ester (0.041g, 0.101mmol) and sodium hydride (0.010g,  
0.25mmol, 60% suspension on mineral oil), pre-stirred for 1  
h at ambient temperature. The reaction mixture is stirred  
10 at ambient temperature 48h, diluted with ethyl acetate, and  
washed with 1N HCl and water. The organic layer is dried,  
concentrated, and purified by LCMS to provide the title  
compound. <sup>1</sup>H NMR (400MHz, CDCl<sub>3</sub>): δ 7.58-7.53 (m, 5H), 7.43  
(t, 2H, *J* = 7.9 Hz), 7.35 (d, 1H, *J* = 7.5 Hz), 7.33 (d, 2H,  
15 *J* = 7.9 Hz), 7.05 (d, 2H, *J* = 8.8 Hz), 6.72 (d, 2H, *J* = 8.8  
Hz), 4.44, 4.38 (AB<sub>q</sub>, 2H, *J* = 14.7 Hz), 3.96 (t, 3H, *J* = 6.3  
Hz), 3.71-3.64 (m, 1H), 3.58-3.53 (m, 1H), 3.49-3.42 (m,  
2H), 3.06, 3.04 (AB<sub>q</sub>, 2H, *J* = 13.7 Hz), 2.99, 2.93 (AB<sub>q</sub>, 2H,  
*J* = 13.7 Hz), 2.86 (s, 3H), 2.27-2.21 (m, 1H), 1.93-1.87 (m,  
20 1H), 1.60-1.53 (m, 2H), 1.47 (s, 3H), 1.40-1.31 (m, 2H),  
0.92 (t, 3H, *J* = 7.4 Hz). MS [EI+] 545 (M+H)<sup>+</sup>, [EI-] 543 (M-  
H)<sup>+</sup>.

**Example 40**

**2-Butoxy-2-methyl-3-(4-{2-[3-methyl-2-oxo-1-(3-trifluoromethyl-benzyl)-imidazolidin-4-yl]-ethoxy}-phenyl)-propionic acid**

5

Step A

**2-Butoxy-2-methyl-3-(4-{2-[3-methyl-2-oxo-1-(3-trifluoromethyl-benzyl)-imidazolidin-4-yl]-ethoxy}-phenyl)-propionic acid ethyl ester:** 3-Trifluoromethylbenzyl bromide (0.037g, 0.157mmol) and tetrabutyl ammonium iodide (catalytic amount) are added to a 0°C suspension of 2-Butoxy-2-methyl-3-(4-[2-(3-methyl-2-oxo-imidazolidin-4-yl)-ethoxy]-phenyl)-propionic acid ethyl ester (0.043g, 0.105mmol) and sodium hydride (0.010g, 0.26mmol, 60% suspension on mineral oil), pre-stirred for 1 h at ambient temperature. The reaction mixture is stirred at ambient temperature for 48h, diluted with ethyl acetate, and washed with 1N HCl and water. The organic layer is dried over MgSO<sub>4</sub>, concentrated *in vacuo*, and used in the next step. MS [EI+] 565 (M+H)<sup>+</sup>.

Step B

**2-Butoxy-2-methyl-3-(4-{2-[3-methyl-2-oxo-1-(3-trifluoromethyl-benzyl)-imidazolidin-4-yl]-ethoxy}-phenyl)-propionic acid** : A solution of 2-Butoxy-2-methyl-3-(4-{2-[3-methyl-2-oxo-1-(3-trifluoromethyl-benzyl)-imidazolidin-4-yl]-ethoxy}-phenyl)-propionic acid ethyl ester and 5N NaOH (0.3mL) in ethanol (3mL) is refluxed under nitrogen for 1h, cooled to ambient temperature, and concentrated. The residue is diluted with 1N HCl, extracted with CH<sub>2</sub>Cl<sub>2</sub>, dried, concentrated, and purified by LCMS to provide the title compound. <sup>1</sup>H NMR (400MHz, CDCl<sub>3</sub>): δ 7.51-7.38 (m, 4H), 7.04 (d, 2H, J = 8.8 Hz), 6.69 (d, 2H, J = 8.8 Hz), 4.46, 4.34 (AB<sub>q</sub>, 2H, J = 14.9 Hz), 3.94 (t, 2H, J = 5.7 Hz), 3.68-3.60 (m, 1H), 3.57-3.52 (m, 1H), 3.48-3.43 (m, 1H), 3.38 (t, 1H, J = 8.8 Hz), 2.98, 2.92 (AB<sub>q</sub>, 2H, J = 14.4 Hz), 2.83 (s, 3H), 2.25-2.18 (m, 1H), 1.91-1.83 (m, 1H), 1.59-1.52 (m, 2H), 1.45 (s, 3H), 1.39-1.30 (m, 2H), 0.91 (t, 3H, J = 7.2 Hz). MS [EI+] 537 (M+H)<sup>+</sup>, [EI-] 535 (M-H)<sup>+</sup>.

Example 41

**2-Butoxy-3-(4-{2-[1-(4-chloro-benzyl)-3-methyl-2-oxo-imidazolidin-4-yl]-ethoxy}-phenyl)-2-methyl-propionic acid**: 4-Chlorobenzyl chloride (0.04g, 0.174mmol) and tetrabutyl ammonium iodide (catalytic amount) are added to a 0°C suspension of 2-butoxy-2-methyl-3-(4-[2-(3-methyl-2-oxo-imidazolidin-4-yl)-ethoxy]-phenyl)-propionic acid ethyl ester (0.047g, 0.116mmol) and sodium hydride (0.012g, 0.29mmol, 60% suspension on mineral oil), pre-stirred for 1 h at ambient temperature. The reaction mixture is stirred at ambient temperature 48h, diluted with ethyl acetate, and washed with 1N HCl and water. The organic layer is dried, concentrated, and purified by LCMS to provide the title

compound.  $^1\text{H}$  NMR (400MHz,  $\text{CDCl}_3$ ):  $\delta$  7.27 (d, 2H,  $J = 8.7$  Hz), 7.17 (d, 2H,  $J = 8.7$  Hz), 7.07 (d, 2H,  $J = 8.7$  Hz), 7.71 (d, 2H,  $J = 8.7$  Hz), 4.34, 4.32 ( $\text{AB}_q$ , 2H,  $J = 15.0$  Hz), 3.95 (t, 3H,  $J = 5.9$  Hz), 3.69-3.62 (m, 1H), 3.61-3.55 (m, 1H), 3.51-3.45 (m, 1H), 3.38 (t, 1H,  $J = 8.7$  Hz), 3.04-2.93 (m, 3H), 2.84 (s, 3H), 2.26-2.18 (m, 1H), 1.92-1.84 (m, 1H), 1.62-1.54 (m, 2H), 1.49 (s, 3H), 1.42-1.33 (m, 3H), 0.93 (t, 3H,  $J = 7.3$  Hz). MS  $[\text{EI}^+]$  503 ( $\text{M}+\text{H}$ ) $^+$ ,  $[\text{EI}^-]$  501 ( $\text{M}-\text{H}$ ) $^+$ .

10

**Example 42****2-Butoxy-3-(4-{2-[1-(4-methoxy-benzyl)-3-methyl-2-oxo-imidazolidin-4-yl]-ethoxy}-phenyl)-2-methyl-propionic acid:**

A solution of 2-Butoxy-2-methyl-3-{4-[2-(3-methyl-2-oxo-imidazolidin-4-yl)-ethoxy]-phenyl}-propionic acid ethyl ester and 5N NaOH (0.2mL) in ethanol (2mL) is refluxed under nitrogen for 1h, cooled to ambient temperature, and concentrated. The residue is diluted with 1N HCl, extracted with  $\text{CH}_2\text{Cl}_2$ , dried, concentrated, and purified by LCMS to provide the title compound (0.024g, 84%).  $^1\text{H}$  NMR (400MHz,  $\text{CDCl}_3$ ):  $\delta$  7.16 (d, 2H,  $J = 140.1$  Hz), 7.08 (d, 2H,  $J = 8.4$  Hz), 6.83 (d, 2H,  $J = 10.1$  Hz), 6.70 (d, 2H,  $J = 8.4$  Hz), 4.32, 4.28 ( $\text{AB}_q$ , 2H,  $J = 14.7$  Hz), 3.92 (t, 2H,  $J = 6.1$  Hz), 3.78 (s, 3H), 3.61-3.44 (m, 3H), 3.31 (t, 1H,  $J = 8.5$  Hz), 3.00, 2.92 ( $\text{AB}_q$ , 2H,  $J = 14.0$  Hz), 2.93 (t, 1H,  $J = 8.5$  Hz), 2.82 (s, 3H), 2.23-2.16 (m, 1H), 1.89-1.81 (m, 1H), 1.61-1.54 (m, 2H), 1.43 (s, 3H), 1.41-1.32 (m, 2H), 0.91 (t, 3H,  $J = 7.3$  Hz). MS  $[\text{EI}^+]$  499 ( $\text{M}+\text{H}$ ) $^+$ ,  $[\text{EI}^-]$  497 ( $\text{M}-\text{H}$ ) $^+$ .

30

**Example 43****2-Ethoxy-3-(4-{2-[1-(4-methoxy-benzyl)-3-methyl-2-oxo-imidazolidin-4-yl]-ethoxy}-phenyl)-2-methyl-propionic acid**

Step A

**2-Ethoxy-3-(4-{2-[1-(4-methoxy-benzyl)-3-methyl-2-oxo-imidazolidin-4-yl]-ethoxy}-phenyl)-2-methyl-propionic acid ethyl ester** : A slurry of 2 2-Hydroxy-3-(4-{2-[1-(4-methoxy-benzyl)-3-methyl-2-oxo-imidazolidin-4-yl]-ethoxy}-phenyl)-2-methyl-propionic acid ethyl ester (3.75g, 7.97mmol) and sodium hydride (1.28g, 31.88mmol, 60% dispersion on mineral oil) in DMF (30mL) is stirred for 45minutes, then cooled to 0° C. Iodoethane (12.75mL, 159.38mmol, d=1.95) and 18-crown-6 (8.43g, 31.88mmol) are added, then the reaction mixture is stirred 18h at ambient temperature, then diluted with ethyl acetate. The organic layer is washed, dried, concentrated, and purified by flash chromatography (20% acetone in hexanes) to provide the title compound (3.23g, 80%). <sup>1</sup>H NMR (400MHz, CDCl<sub>3</sub>): δ 7.17(d, 2H, J = 8.6 Hz), 7.08 (d, 2H, J = 8.6 Hz), 6.83 (d, 2H, J = 9.1 Hz), 6.70 (d, 2H, J = 9.1 Hz), 4.31, 4.27 (AB<sub>q</sub>, 2H, J = 14.5 Hz), 4.17-4.11 (m, 2H), 3.93 (t, 2H, J = 6.4 Hz), 3.77 (s, 3H), 3.60-3.53 (m, 1H), 3.48-3.39 (m, 2H), 3.32 (t, 1H, J = 8.6 Hz), 2.98-2.89 (m, 3H), 2.81(s, 3H), 2.24-2.16 (m, 1H), 1.88-1.80 (m, 1H), 1.29 (s, 3H), 1.25-1.15 (m, 6H). R<sub>f</sub>=0.54 in 50% acetone in hexanes.


Step B

**2-Ethoxy-3-(4-{2-[1-(4-methoxy-benzyl)-3-methyl-2-oxo-imidazolidin-4-yl]-ethoxy}-phenyl)-2-methyl-propionic acid** : A solution of 2-ethoxy-3-(4-{2-[1-(4-methoxy-benzyl)-3-methyl-2-oxo-imidazolidin-4-yl]-ethoxy}-phenyl)-2-methyl-propionic acid ethyl ester (0.038g, 0.076mmol) and 5N NaOH (0.2mL) is refluxed for 1h, cooled to ambient temperature, and concentrated. The residue is diluted with CH<sub>2</sub>Cl<sub>2</sub>, washed, dried, and concentrated in vacuo to provide the

title compound (.027g, 77%).  $^1\text{H}$  NMR (400MHz,  $\text{CDCl}_3$ ):  $\delta$  7.16 (d, 2H,  $J = 8.7$  Hz), 7.07 (d, 2H,  $J = 8.1$  Hz), 6.83 (d, 2H,  $J = 8.7$  Hz), 6.70 (d, 2H,  $J = 8.1$  Hz), 4.31, 4.27 (Abq, 2H,  $J = 14.9$  Hz), 3.92 (t, 2H,  $J = 6.2$  Hz), 3.78 (s, 3H), 3.59-3.52 (m, 3H), 3.33 (t, 1H,  $J = 8.2$  Hz), 3.03-2.91 (m, 3H), 2.81 (s, 3H), 2.21-2.17 (m, 1H), 1.90-1.81 (m, 1H), 1.43 (s, 3H), 1.20 (t, 3H,  $J = 7.0$  Hz). MS [ES+]  $m/z$  exact mass calcd for  $\text{C}_{26}\text{H}_{35}\text{N}_2\text{O}_6$  471.2495, found 471.2507.

#### Example 44

**2-Ethoxy-2-methyl-3-(4-(2-[3-methyl-2-oxo-1-(4-trifluoromethyl-benzyl)-imidazolidin-4-yl]-ethoxy)-phenyl)-propionic acid**




#### Step A

**2-Hydroxy-2-methyl-3-(4-(2-[3-methyl-2-oxo-1-(4-trifluoromethyl-benzyl)-imidazolidin-4-yl]-ethoxy)-phenyl)-propionic acid ethyl ester** : Cesium carbonate (8.72g, 26.76mmol) is added to a solution of 2-hydroxy-3-(4-(2-[1-(4-methoxy-benzyl)-3-methyl-2-oxo-imidazolidin-4-yl]-ethoxy)-phenyl)-2-methyl-propionic acid ethyl ester (2.0g, 8.92mmol) and Toluene-4-sulfonic acid 2-[3-methyl-2-oxo-1-(4-trifluoromethyl-benzyl)-imidazolidin-4-yl]-ethyl ester (4.34g, 9.51mmol) in DMF (50mL). The resultant mixture is



stirred at 55°C under an atmosphere of nitrogen for 18 h,  
 then diluted with ethyl acetate. The organic layer is  
 washed, dried, concentrated, and purified by flash  
 chromatography (25% acetone in hexanes) to provide the title  
 5 compound. <sup>1</sup>H NMR (400MHz, CDCl<sub>3</sub>): δ 7.95 (s, 1H), 7.50 (d,  
 2H, J = 8.2 Hz), 7.32 (d, 2H, J = 8.2 Hz), 7.03 (d, 2H, J =  
 8.2 Hz), 6.65 (d, 2H, J = 8.2 Hz), 4.40, 4.32 (AB<sub>q</sub>, 2H, J =  
 14.7 Hz), 4.14-4.07 (m, 2H), 3.91 (t, 2H, J = 5.7 Hz), 3.61-  
 3.55 (m, 1H), 3.33 (t, 1H, J = 8.2 Hz), 2.96-2.79 (m, 6H),  
 10 2.22-2.15 (m, 1H), 1.87-1.80 (m, 1H), 1.40 (s, 3H), 1.21 (t,  
 3H, J = 7.4 Hz). R<sub>f</sub>=0.38 in 50% acetone in hexanes.

### Step B



15

**2-Ethoxy-2-methyl-3-(4-{2-[3-methyl-2-oxo-1-(4-trifluoromethyl-benzyl)-imidazolidin-4-yl]-ethoxy}-phenyl)-propionic acid ethyl ester** : A slurry of 2-hydroxy-3-(4-hydroxy-phenyl)-2-methyl-propionic acid ethyl ester  
 20 (0.509g, 1.00mmol) and sodium hydride (0.16g, 4.00mmol, 60% dispersion on mineral oil) in DMF (3mL) is stirred for 45minutes, then cooled to 0° C. Iodoethane (1.60mL, 20.02mmol, d=1.95) and 18-crown-6 (1.06g, 4.00mmol) are added, then the reaction mixture is stirred 18h at ambient  
 25 temperature, then diluted with ethyl acetate. The organic layer is washed, dried, concentrated, and purified by flash chromatography (20% acetone in hexanes) to provide the title

compound (3.23g, 80%). MS [EI+] 537 (M+H)<sup>+</sup>. R<sub>f</sub>=0.44 in 50% acetone in hexanes.

### Step C

5 **2-Ethoxy-2-methyl-3-(4-{2-[3-methyl-2-oxo-1-(4-trifluoromethyl-benzyl)-imidazolidin-4-yl]-ethoxy}-phenyl)-propionic acid** : A solution of 2-Ethoxy-2-methyl-3-(4-{2-[3-methyl-2-oxo-1-(4-trifluoromethyl-benzyl)-imidazolidin-4-yl]-ethoxy}-phenyl)-propionic acid ethyl ester (0.19g,  
10 0.347mmol) and 5N NaOH (0.5mL) in ethanol (5mL) is refluxed for 1h, cooled to ambient temperature, and concentrated. The residue is diluted with CH<sub>2</sub>Cl<sub>2</sub>, washed, dried, and concentrated *in vacuo* to provide the title compound. <sup>1</sup>H NMR (400MHz, CDCl<sub>3</sub>): δ 7.63 (d, 2H, J = 8.4 Hz), 7.44 (d, 2H, J = 8.4 Hz), 7.08 (d, 2H, J = 8.7 Hz), 6.76 (d, 2H, J = 8.7 Hz), 4.37, 4.35 (AB<sub>q</sub>, 2H, J = 16.4 Hz), 3.99-3.89 (m, 2H), 3.63-3.53 (m, 1H), 3.43-3.23 (m, 2H), 2.98 (t, 1H, J = 8.8 Hz), 2.86, 2.84 (AB<sub>q</sub>, 2H, J = 15.4 Hz), 2.71 (s, 3H), 2.51-2.49 (m, 3H), 2.23-2.13 (m, 1H), 1.86-1.74 (m, 1H), 1.15 (s, 20 3H), 1.08 (t, 3H, J = 6.9 Hz). MS [ES+] m/z exact mass calcd for C<sub>26</sub>H<sub>32</sub>N<sub>2</sub>O<sub>5</sub>F<sub>3</sub> 509.2263, found 509.2288.

### Example 45

25 **2-Methyl-3-(4-{2-[3-methyl-2-oxo-1-(4-trifluoromethyl-benzyl)-imidazolidin-4-yl]-ethoxy}-phenyl)-2-propoxy-propionic acid**

The title compound is prepared using procedures substantially as described herein and purified by LCMS to provide the title compound (0.017g, 34% after two steps). <sup>1</sup>H  
30 NMR (400MHz, CDCl<sub>3</sub>): δ 7.48 (d, 2H, J = 8.0 Hz), 7.29 (d, 2H, J = 7.2 Hz), 7.01 (d, 2H, J = 7.2 Hz), 6.63 (d, 2H, J = 8.0 Hz), 4.45, 4.39 (AB<sub>q</sub>, 2H, J = 15.4 Hz), 3.96 (t, 2H, J =

6.0 Hz), 3.69-3.62 (m, 1H), 3.57-3.52 (m, 1H), 3.46-3.37 (m, 3H), 3.04-2.93 (m, 4H), 2.85 (s, 3H), 2.28-2.20 (m, 1H), 1.93-1.84 (m, 1H), 1.67-1.58 (m, 2H), 1.49 (s, 3H), 0.93 (t, 3H,  $J = 7.4$  Hz). MS [EI+] 523 (M+H)<sup>+</sup>, [EI-] 521 (M-H)<sup>+</sup>.

5

#### Example 46

2-Butoxy-2-methyl-3-(4-{2-[3-methyl-2-oxo-1-(4-trifluoromethyl-benzyl)-imidazolidin-4-yl]-ethoxy}-phenyl)-  
10 propionic acid

The title compound is prepared substantially using the procedures described herein and purified by LCMS to provide the title compound (0.022g, 52% after two steps). <sup>1</sup>H NMR (400MHz, CDCl<sub>3</sub>): δ 7.56 (d, 2H,  $J = 8.3$  Hz), 7.36 (d, 2H,  $J = 8.3$  Hz), 7.07 (d, 2H,  $J = 9.0$  Hz), 6.71 (d, 2H,  $J = 9.0$  Hz), 4.46, 4.38 (AB<sub>q</sub>, 2H,  $J = 15.2$  Hz), 3.96 (t, 2H,  $J = 5.5$  Hz), 3.67-3.61 (m, 1H), 3.60-3.54 (m, 1H), 3.50-3.45 (m, 1H), 3.39 (t, 1H,  $J = 9.0$  Hz), 3.03-2.92 (m, 3H), 2.85 (s, 3H), 2.27-2.20 (m, 1H), 1.93-1.84 (m, 1H), 1.62-1.54 (m, 15 2H), 1.48 (s, 3H), 1.41-1.32 (m, 2H), 0.92 (t, 3H,  $J = 6.9$  Hz). MS [EI+] 537 (M+H)<sup>+</sup>, [EI-] 535 (M-H)<sup>+</sup>.

20

#### Example 47

2-Ethoxy-2-methyl-3-(4-{2-[3-methyl-2-oxo-1-(4-trifluoromethoxy-benzyl)-imidazolidin-4-yl]-ethoxy}-phenyl)-  
25 propionic acid

Trifluoroacetic acid (70mL) is added dropwise to a solution of 2-Ethoxy-2-methyl-3-{4-[2-(3-methyl-2-oxo-imidazolidin-4-yl)-ethoxy]-phenyl}-propionic acid ethyl ester (3.19g, 30 6.40mmol) and triethylsilane (2.04mL, 12.8mmol, d=0.728). The reaction mixture is stirred at ambient temperature for 4 hours, concentrated, and diluted with ethyl acetate. The

solution is washed, then dried, and concentrated in vacuo to provide the title compound (2.2g, 92%). <sup>1</sup>H NMR (400MHz, CDCl<sub>3</sub>): δ 7.11 (d, 2H, J = 9.3 Hz), 6.76 (d, 2H, J = 9.3 Hz), 4.19-4.10 (m, 2H), 4.04-3.98 (m, 2H), 3.77-3.69 (m, 1H), 3.56 (t, 1H, J = 8.8 Hz), 3.51-3.38 (m, 2H), 3.22 (t, 1H, J = 8.8 Hz), 2.96, 2.94 (AB<sub>q</sub>, 2H, J = 14.0 Hz), 2.78 (s, 3H), 2.27-2.20 (m, 1H), 1.99-1.92 (m, 1H). MS [EI+] 379 (M+H)<sup>+</sup>.

#### Step A

10 **2-Ethoxy-2-methyl-3-(4-{2-[3-methyl-2-oxo-1-(4-trifluoromethoxy-benzyl)-imidazolidin-4-yl]-ethoxy}-phenyl)-propionic acid ethyl ester** : 4-(Trifluoromethoxy)benzyl bromide (0.03mL, 0.19mmol) and tetrabutyl ammonium iodide (catalytic amount) are added to a 0°C suspension of 2-Ethoxy-2-methyl-3-  
15 {4-[2-(3-methyl-2-oxo-imidazolidin-4-yl)-ethoxy]-phenyl}-propionic acid ethyl ester (0.048g, 0.127mmol) and sodium hydride (0.013g, 0.32mmol, 60% suspension on mineral oil), pre-stirred for 1 h at ambient temperature. The reaction mixture is stirred at ambient temperature for 90 minutes,  
20 diluted with ethyl acetate, and washed. The organic layer is dried, concentrated, and used in the next step. MS [EI+] 553 (M+H)<sup>+</sup>.

#### Step B

**2-Ethoxy-2-methyl-3-(4-{2-[3-methyl-2-oxo-1-(4-trifluoromethoxy-benzyl)-imidazolidin-4-yl]-ethoxy}-phenyl)-propionic acid** : A solution of 2-ethoxy-2-methyl-3-(4-{2-[3-methyl-2-oxo-1-(4-trifluoromethoxy-benzyl)-imidazolidin-4-yl]-ethoxy}-phenyl)-propionic acid ethyl ester and 5N NaOH (0.3mL) in ethanol (3mL) is refluxed for 1h, cooled to  
30 ambient temperature, and concentrated. The residue is diluted with CH<sub>2</sub>Cl<sub>2</sub>, dried, and concentrated in vacuo to provide the title compound (0.022g, 32% after two steps).

<sup>1</sup>H NMR (400MHz, CDCl<sub>3</sub>): δ 7.24-7.19 (m, 1H), 7.09 (d, 2H, J = 8.3 Hz), 6.83-6.78 (m, 3H), 6.69 (d, 2H, J = 8.3 Hz), 4.38, 4.28 (AB<sub>q</sub>, 2H, J = 14.8 Hz), 3.94-3.91 (m, 3H), 3.77 (s, 3H), 3.60-3.53 (m, 3H), 3.36 (t, 1H, J = 9.0 Hz), 3.04-2.92 (m, 1H), 1.89-1.83 (m, 1H), 1.40 (s, 3H), 1.23 (t, 3H, J = 6.9 Hz). MS [EI+] 525 (M+H)<sup>+</sup>, [EI-] 523 (M-H)<sup>+</sup>. R<sub>f</sub>=0.27 in 10% CH<sub>3</sub>OH in CH<sub>2</sub>Cl<sub>2</sub>.

#### Example 48

10      **2-Ethoxy-3-(4-{2-[1-(3-methoxy-benzyl)-3-methyl-2-oxo-imidazolidin-4-yl]-ethoxy}-phenyl)-2-methyl-propionic acid**  
The title compound is prepared using substantially the procedures described herein, and purified by LCMS to provide the title compound (0.022g, 32% after two steps). <sup>1</sup>H NMR (400MHz, CDCl<sub>3</sub>): δ 7.22 (t, 1H, J = 7.7 Hz), 7.06 (d, 2H, J = 8.3 Hz), 6.83-6.79 (m, 3H), 6.72 (d, 2H, J = 7.1 Hz), 4.37, 4.29 (AB<sub>q</sub>, 2H, J = 14.8 Hz), 3.95 (t, 2H, J = 6.3 Hz), 3.68-3.52 (m, 6H), 3.39 (t, 3H, J = 8.5 Hz), 3.04-2.93 (m, 3H), 2.84 (s, 3H), 2.26-2.20 (m, 1H), 1.92-1.85 (m, 1H), 20 1.49 (s, 3H), 1.24 (t, 3H, J = 7.1 Hz). MS [EI+] 471 (M+H)<sup>+</sup>, [EI-] 469 (M-H)<sup>+</sup>. R<sub>f</sub>=0.27 in 10% CH<sub>3</sub>OH in CH<sub>2</sub>Cl<sub>2</sub>.

#### Example 49

25      **2-Ethoxy-2-methyl-3-(4-{2-[3-methyl-2-oxo-1-(3-trifluoromethoxy-benzyl)-imidazolidin-4-yl]-ethoxy}-phenyl)-propionic acid**


The title compound is prepared using substantially the procedures described purified by LCMS to provide the title compound (0.022g, 32% after two steps). <sup>1</sup>H NMR (400MHz, CDCl<sub>3</sub>): δ 7.33 (t, 1H, J = 8.2 Hz), 7.17 (d, 1H, J = 7.5 Hz), 7.13-7.06 (m, 4H), 6.72 (d, 2H, J = 8.9 Hz), 4.42, 4.34 (AB<sub>q</sub>, 2H, J = 15.3 Hz), 3.96 (t, 2H, J = 5.6 Hz), 3.71-3.52

(m, 3H), 3.41 (t, 1H,  $J = 8.3$  Hz), 3.05-2.93 (m, 3H), 2.85 (s, 3H), 2.27-2.20 (m, 1H), 1.94-1.85 (m, 1H), 1.47 (s, 3H), 1.23 (t, 3H,  $J = 7.3$  Hz). MS [EI+] 525 (M+H)<sup>+</sup>, [EI-] 523 (M-H)<sup>+</sup>.

5

**Example 50**

**2-Ethoxy-2-methyl-3-(4-{2-[3-methyl-2-oxo-1-(3-trifluoromethyl-benzyl)-imidazolidin-4-yl]-ethoxy}-phenyl)-propionic acid**



The title compound is prepared using substantially the  
 10 procedures described herein and purified by LCMS to provide  
 the title compound. <sup>1</sup>H NMR (400MHz, CDCl<sub>3</sub>): δ 7.54-7.42 (m, 4H), 7.07 (d, 2H,  $J = 8.3$  Hz), 6.70 (d, 2H,  $J = 9.3$  Hz), 4.47, 4.37 (AB<sub>q</sub>, 2H,  $J = 15.3$  Hz), 3.96 (t, 2H,  $J = 6.4$  Hz), 3.73-3.65 (m, 1H), 3.64-3.51 (m, 2H), 3.42 (t, 1H,  $J = 8.3$   
 15 Hz), 3.05 (t, 1H,  $J = 7.2$  Hz), 3.01, 2.95 (AB<sub>q</sub>, 2H,  $J = 14.3$  Hz), 2.86 (s, 3H), 2.27-2.20 (m, 1H), 1.94-1.87 (m, 1H), 1.46 (s, 3H), 1.23 (t, 3H,  $J = 7.2$  Hz). MS [EI+] 509 (M+H)<sup>+</sup>, [EI-] 507 (M-H)<sup>+</sup>.

20

**Example 51**

**3-{4-[2-(1-Biphenyl-4-ylmethyl-3-methyl-2-oxo-imidazolidin-4-yl)-ethoxy]-phenyl}-2-ethoxy-2-methyl-propionic acid,**  
 and purified by chromatotron (CH<sub>3</sub>OH/CH<sub>2</sub>Cl<sub>2</sub> gradient) to  
 provide the title compound (0.033g, 55%). <sup>1</sup>H NMR (400MHz,  
 25 CDCl<sub>3</sub>): δ 7.58-7.52 (m, 4H), 7.43 (t, 2H,  $J = 7.8$  Hz), 7.35-7.31 (m, 3H), 7.05 (d, 2H,  $J = 7.8$  Hz), 6.72 (d, 2H,  $J = 8.5$

Hz), 4.45, 4.37 (AB<sub>q</sub>, 2H,  $J = 14.8$  Hz), 3.95 (t, 2H,  $J = 5.6$  Hz), 3.64-3.52 (m, 3H), 3.40 (t, 1H,  $J = 8.3$  Hz), 3.02-2.91 (m, 3H), 2.85 (s, 3H), 2.25-2.20 (m, 1H), 1.91-1.85 (m, 1H), 1.46 (s, 3H), 1.23 (t, 3H,  $J = 7.2$  Hz). MS [ES+]  $m/z$  exact mass calcd for C<sub>31</sub>H<sub>37</sub>N<sub>2</sub>O<sub>5</sub> 517.2702, found 517.2711.  $R_f=0.31$  in 10% CH<sub>3</sub>OH in CH<sub>2</sub>Cl<sub>2</sub>.

### Example 52

3-(4-{2-[1-(3,4-Difluoro-benzyl)-3-methyl-2-oxo-  
imidazolidin-4-yl]-ethoxy}-phenyl)-2-methyl-2-phenoxy-  
propionic acid,

and purified by flash chromatography to provide the title compound (0.42g, 63%). <sup>1</sup>H NMR (400MHz, CDCl<sub>3</sub>):  $\delta$  7.26-7.23 (m, 3H), 7.19 (d, 2H,  $J = 8.3$  Hz), 7.11-7.04 (m, 3H), 6.91 (d, 2H,  $J = 8.3$  Hz), 6.92-6.90 (m, 1H), 6.75 (d, 2H,  $J = 8.3$  Hz), 4.21, 4.15 (AB<sub>q</sub>, 2H,  $J = 14.8$  Hz), 3.87 (t, 2H,  $J = 6.6$  Hz), 3.59-3.53 (m, 1H), 3.30n (t, 1H,  $J = 8.2$  Hz), 3.23, 3.03 (AB<sub>q</sub>, 2H,  $J = 13.1$  Hz), 2.91 (t, 1H,  $J = 8.2$  Hz), 2.74 (s, 3H), 2.16-2.10 (m, 1H), 1.81-1.76 (m, 1H), 1.42 (s, 3H). MS [EI+] 525 (M+H)<sup>+</sup>, [EI-] 523 (M-H)<sup>+</sup>.

### Example 53

2-Ethoxy-2-methyl-3-(4-{2-[3-methyl-2-oxo-1-(4-trifluoromethyl-benzyl)-imidazolidin-4-yl]-ethoxy}-phenyl)-propionic acid

and concentrated *in vacuo* to provide the title compound (0.029g, 83%). <sup>1</sup>H NMR (400MHz, CDCl<sub>3</sub>):  $\delta$  7.55 (d, 2H,  $J = 8.6$  Hz), 7.36 (d, 2H,  $J = 8.6$  Hz), 7.07 (d, 2H,  $J = 8.6$  Hz), 6.70 (d, 2H,  $J = 8.6$  Hz), 4.45, 4.37 (AB<sub>q</sub>, 2H,  $J = 15.6$  Hz), 3.95 (t, 2H,  $J = 6.2$  Hz), 3.66-3.51 (m, 3H), 3.37 (t, 1H,  $J = 8.6$  Hz), 3.03-2.92 (m, 3H), 2.84 (s, 3H), 2.26-2.19 (m,

1H), 2.26-2.19 (m, 1H), 1.42 (s, 3H), 1.22 (t, 3H,  $J = 7.0$  Hz). MS [EI+] 509 (M+H)<sup>+</sup>, [EI-] 507 (M-H)<sup>+</sup>.

#### Example 54

5        **2-Ethoxy-2-methyl-3-(4-{2-[3-methyl-2-oxo-1-(4-trifluoromethyl-benzyl)-imidazolidin-4-yl]-ethoxy}-phenyl)-propionic acid**

The title compound is prepared using substantially the procedures described herein and concentrated *in vacuo* to provide the title compound (0.036g, 88%). <sup>1</sup>H NMR (400MHz, CDCl<sub>3</sub>):  $\delta$  7.55 (d, 2H,  $J = 8.6$  Hz), 7.36 (d, 2H,  $J = 8.6$  Hz), 7.07 (d, 2H,  $J = 8.6$  Hz), 6.70 (d, 2H,  $J = 8.6$  Hz), 4.45, 4.37 (AB<sub>q</sub>, 2H,  $J = 15.6$  Hz), 3.95 (t, 2H,  $J = 6.2$  Hz), 3.66-3.51 (m, 3H), 3.37 (t, 1H,  $J = 8.6$  Hz), 3.03-2.92 (s, 3H), 2.26-2.19 (m, 1H), 1.91-1.84 (m, 1H), 1.42 (s, 3H), 1.22 (t, 3H,  $J = 7.0$  Hz). MS [EI+] 509 (M+H)<sup>+</sup>, [EI-] 507 (M-H)<sup>+</sup>.

#### Example 55

20        **2-Methyl-3-(4-{2-[3-methyl-2-oxo-1-(4-trifluoromethyl-benzyl)-imidazolidin-4-yl]-ethoxy}-phenyl)-2-phenoxy-propionic acid**

#### Step A

25        **3-(4-{2-[3-(4-Methoxy-benzyl)-2-oxo-1-(4-trifluoromethyl-benzyl)-imidazolidin-4-yl]-ethoxy}-phenyl)-2-methyl-2-phenoxy-propionic acid ethyl ester:** Cesium carbonate (2.18g, 6.702mmol) is added to a solution of 3-(4-Hydroxy-phenyl)-2-methyl-2-phenoxy-propionic acid ethyl ester (1.55g, 5.155mmol) and Toluene-4-sulfonic acid 2-[3-methyl-2-oxo-1-(4-trifluoromethyl-benzyl)-imidazolidin-4-yl]-ethyl ester (3.19g, 5.67mmol) in DMF (30mL). The resultant mixture is



stirred at 55°C under an atmosphere of nitrogen for 18 h, then diluted with ethyl acetate. The organic layer is washed, then dried, concentrated, and purified by flash chromatography (17% acetone in hexanes) to provide the title compound (3.46g, 97%). <sup>1</sup>H NMR (400MHz, CDCl<sub>3</sub>): δ 7.57 (d, 2H, J = 7.01 Hz), 7.37 (d, 2H, J = 7.1 Hz), 7.23-7.19 (m, 4H), 7.13 (d, 2H, J = 8.7 Hz), 6.97 (t, 1H, J = 7.1 Hz), 6.87-6.80 (m, 4H), 6.67 (d, 2H, J = 8.7 Hz), 4.79, 4.07 (AB<sub>q</sub>, 2H, J = 15.0 Hz), 4.47, 4.43 (AB<sub>q</sub>, 2H, J = 15.0 Hz), 4.20 (q, 2H, J = 7.1 Hz), 3.92-3.87 (m, 2H), 3.80 (s, 3H), 3.61-3.55 (m, 1H), 3.31 (t, 2H, J = 8.7 Hz), 3.24, 3.10 (AB<sub>q</sub>, 2H, J = 14.2 Hz), 3.00 (t, 1H, J = 7.9 Hz), 2.21-2.14 (m, 1H), 1.87-1.78 (m, 1H), 1.38 (s, 3H), 1.21 (t, 3H, J = 7.1 Hz). MS [EI+] 691 (M+H)<sup>+</sup>. R<sub>f</sub>=0.49 in 50% acetone in hexanes.

#### Step B

**2-Methyl-3-(4-{2-[2-oxo-1-(4-trifluoromethyl-benzyl)-imidazolidin-4-yl]-ethoxy}-phenyl)-2-phenoxy-propionic acid ethyl ester:** Trifluoroacetic acid (60mL) is added dropwise to a solution of 3-(4-{2-[3-(4-Methoxy-benzyl)-2-oxo-1-(4-trifluoromethyl-benzyl)-imidazolidin-4-yl]-ethoxy}-phenyl)-2-methyl-2-phenoxy-propionic acid ethyl ester (3.46g, 5.01mmol) and triethylsilane (1.6mL, 10.0mmol, d=0.728). The reaction mixture is stirred at ambient temperature for 2h, concentrated, and diluted with ethyl acetate. The solution is washed, then dried, and concentrated in vacuo to provide the title compound. <sup>1</sup>H NMR (400MHz, CDCl<sub>3</sub>): δ 7.58 (d, 2H, J = 8.2 Hz), 7.38 (d, 2H, J = 8.2 Hz), 7.23-7.14 (m, 4H), 6.97 (t, 1H, J = 7.4 Hz), 6.82 (d, 2H, J = 7.4 Hz), 6.78 (d, 2H, J = 7.4 Hz), 4.44, 4.38 (AB<sub>q</sub>, 2H, J = 15.6 Hz), 4.20 (q, 2H, J = 7.4 Hz), 4.07-3.99 (m, 2H), 3.96-3.89 (m,

1H), 3.47 (t, 1H,  $J = 8.2$  Hz), 3.27, 3.09 (AB<sub>q</sub>, 2H,  $J = 14.1$  Hz), 3.04, 3.02 (AB<sub>q</sub>, 1H,  $J = 6.7$  Hz), 2.08-1.99 (m, 1H), 1.98-1.92 (m, 1H), 1.38 (s, 3H), 1.21 (t, 3H,  $J = 6.7$  Hz). MS [EI+] 571 (M+H)<sup>+</sup>, [EI-] 569 (M-H)<sup>+</sup>.  $R_f=0.07$  in 33%

5 acetone in hexanes.

#### Step C

**2-Methyl-3-(4-{2-[3-methyl-2-oxo-1-(4-trifluoromethyl-benzyl)-imidazolidin-4-yl]-ethoxy}-phenyl)-2-phenoxy-propionic acid ethyl ester:** A slurry of 2-Methyl-3-(4-{2-[2-oxo-1-(4-trifluoromethyl-benzyl)-imidazolidin-4-yl]-ethoxy}-phenyl)-2-phenoxy-propionic acid ethyl ester (0.544g, 10 0.953mmol) and sodium hydride (.042g, 1.05 mmol, 60% dispersion on mineral oil) in DMF (10mL) is stirred for 1h, then cooled to 0° C. Iodomethane (0.60mL, 9.53mmol, 15 d=2.28) is added, then the reaction mixture is stirred 18h at ambient temperature and diluted with ethyl acetate. The organic layer is washed, dried, concentrated, and purified by flash chromatography to provide the title compound. <sup>1</sup>H NMR (400MHz, CDCl<sub>3</sub>):  $\delta$  7.56 (d, 2H,  $J = 8.0$  Hz), 7.37 (d, 20 2H,  $J = 8.0$  Hz), 7.26-7.19 (m, 2H), 7.15 (d, 2H,  $J = 8.9$  Hz), 6.97 (t, 1H,  $J = 8.0$  Hz), 6.82 (d, 2H,  $J = 8.0$  Hz), 6.74 (d, 2H,  $J = 8.9$  Hz), 4.46, 4.38 (AB<sub>q</sub>, 2H,  $J = 15.1$  Hz), 4.20 (q, 2H,  $J = 7.1$  Hz), 3.97 (t, 2H,  $J = 7.1$  Hz), 3.67-3.59 (m, 1H), 3.78 (t, 1H,  $J = 8.9$  Hz), 3.26, 3.10 (AB<sub>q</sub>, 2H, 25  $J = 14.2$  Hz), 2.99 (t, 1H,  $J = 8.9$  Hz), 2.85 (s, 3H), 2.28-2.21 (m, 1H), 1.93-1.84 (m, 1H), 1.39 (s, 3H), 1.21 (t, 3H,  $J = 7.1$  Hz). MS [EI+]. 585 (M+H)<sup>+</sup>.  $R_f=0.35$  in 50% acetone in hexanes.

#### Step D

30 **2-Methyl-3-(4-{2-[3-methyl-2-oxo-1-(4-trifluoromethyl-benzyl)-imidazolidin-4-yl]-ethoxy}-phenyl)-2-phenoxy-propionic acid:** A solution of 2-methyl-3-(4-{2-[3-methyl-2-

oxo-1-(4-trifluoromethyl-benzyl)-imidazolidin-4-yl]-ethoxy}-phenyl)-2-phenoxy-propionic acid ethyl ester and 5N NaOH (0.7mL) in ethanol (6mL) is refluxed for 1h, cooled to ambient temperature, and concentrated. The residue is  
5 diluted with CH<sub>2</sub>Cl<sub>2</sub>, washed, dried, and concentrated to provide the title compound (0.186g, 100%). <sup>1</sup>H NMR (400MHz, CDCl<sub>3</sub>): δ 7.54 (d, 2H, J = 8.3 Hz), 7.35 (d, 2H, J = 8.3 Hz), 7.24-7.17 (m, 4H), 7.00 (t, 1H, J = 7.4 Hz), 6.89 (d, 2H, J = 7.4 Hz), 6.73 (d, 2H, J = 8.3 Hz), 4.45, 4.37 (AB<sub>q</sub>,  
10 2H, J = 15.7 Hz), 3.96 (t, 2H, J = 5.5 Hz), 3.68-3.62 (m, 1H), 3.38 (t, 1H, J = 8.3 Hz), 3.32, 3.10 (AB<sub>q</sub>, 2H, J = 12.9 Hz), 3.00 (t, 1H, J = 8.3 Hz), 2.84 (s, 3H), 2.27-2.19 (m, 1H), 1.92-1.85 (m, 1H), 1.39 (s, 3H). MS [EI+] 557 (M+H)<sup>+</sup>, [EI-] 555 (M-H)<sup>+</sup>.

15

### Example 56


3-(4-{2-[3-Ethyl-2-oxo-1-(4-trifluoromethyl-benzyl)-imidazolidin-4-yl]-ethoxy}-phenyl)-2-methyl-2-phenoxy-propionic acid  
20

#### Step A

3-(4-{2-[3-(4-Methoxy-benzyl)-2-oxo-1-(4-trifluoromethyl-benzyl)-imidazolidin-4-yl]-ethoxy}-phenyl)-2-methyl-2-phenoxy-propionic acid ethyl ester: Cesium carbonate (2.49g, 7.647mmol) is added to a solution of 3-(4-Hydroxy-phenyl)-2-methyl-2-phenoxy-propionic acid ethyl ester (1.77g, 5.88mmol) and toluene-4-sulfonic acid 2-[3-(4-methoxy-benzyl)-2-oxo-1-(4-trifluoromethyl-benzyl)-imidazolidin-4-yl]-ethyl ester (3.64g, 6.47mmol) in DMF (40mL). The  
25 resultant mixture is stirred at 55°C under an atmosphere of nitrogen for 18 h, then diluted with ethyl acetate. The organic layer is washed, then dried, concentrated, and  
30

purified by flash chromatography (acetone/hexanes gradient) to provide the title compound (3.80g, 94%).  $^1\text{H}$  NMR (400MHz,  $\text{CDCl}_3$ ):  $\delta$  7.57 (d, 2H,  $J = 8.4$  Hz), 7.37 (d, 2H,  $J = 7.8$  Hz), 7.23-7.19 (m, 4H), 7.13 (d, 2H,  $J = 8.4$  Hz), 6.97 (t, 1H,  $J = 7.8$  Hz), 6.86 (d, 2H,  $J = 8.4$  Hz), 6.81 (d, 2H,  $J = 8.4$  Hz), 6.67 (d, 2H,  $J = 7.8$  Hz), 4.66, 4.20 (AB<sub>q</sub>, 2H,  $J = 14.9$  Hz), 4.47, 4.43 (AB<sub>q</sub>, 2H,  $J = 15.5$  Hz), 4.20 (q, 2H,  $J = 7.1$  Hz), 3.89-3.85 (m, 2H), 3.80 (s, 3H), 3.61-3.55 (m, 1H), 3.31 (t, 1H,  $J = 9.1$  Hz), 3.26, 3.08 (AB<sub>q</sub>, 2H,  $J = 13.6$  Hz), 3.00 (t, 1H,  $J = 9.1$  Hz), 2.20-2.14 (m, 1H), 1.87-1.78 (m, 1H), 1.38 (s, 3H), 1.21 (t, 3H,  $J = 7.1$  Hz). MS [EI+] 691 (M+H)<sup>+</sup>.  $R_f=0.46$  in 50% acetone in hexanes.

#### Step B




15

**2-Methyl-3-(4-{2-[2-oxo-1-(4-trifluoromethyl-benzyl)-imidazolidin-4-yl]-ethoxy}-phenyl)-2-phenoxy-propionic acid ethyl ester:** Trifluoroacetic acid (66mL) is added dropwise to a solution of 3-(4-{2-[3-(4-Methoxy-benzyl)-2-oxo-1-(4-trifluoromethyl-benzyl)-imidazolidin-4-yl]-ethoxy}-phenyl)-2-methyl-2-phenoxy-propionic acid ethyl ester (3.80g, 5.50mmol) and triethylsilane (1.76mL, 11.0mmol,  $d=0.728$ ). The reaction mixture is stirred at ambient temperature for 4hours, concentrated *in vacuo*, and diluted with ethyl acetate. The solution is washed, then dried, and

25

concentrated to provide the title compound.  $^1\text{H}$  NMR (400MHz,  $\text{CDCl}_3$ ):  $\delta$  7.58 (d, 2H,  $J = 7.7$  Hz), 7.38 (d, 2H,  $J = 7.7$  Hz), 7.22 (t, 2H,  $J = 9.1$  Hz), 7.15 (d, 2H,  $J = 9.1$  Hz), 6.97 (t, 1H,  $J = 7.7$  Hz), 6.82 (d, 2H,  $J = 7.7$  Hz), 6.78 (d, 2H,  $J = 9.1$  Hz), 4.41 (q, 2H,  $J = 14.8$  Hz), 4.22, 4.18 (ABq, 2H,  $J = 7.0$  Hz), 4.04-4.01 (m, 2H), 3.97-3.90 (m, 1H), 3.43 (t, 1H,  $J = 8.4$  Hz), 3.26, 3.10 (ABq, 2H,  $J = 14.1$  Hz), 3.04, 3.02 (ABq, 1H,  $J = 7.0$  Hz), 2.08-1.99 (m, 1H), 1.98-1.90 (m, 1H), 1.38 (s, 3H), 1.21 (t, 3H,  $J = 7.0$  Hz). MS [EI+] 571 (M+H) $^+$ , [EI-] 569 (M-H) $^+$ .  $R_f=0.17$  in 33% acetone in hexanes.

### Step C



15

**3-(4-{2-[3-Ethyl-2-oxo-1-(4-trifluoromethyl-benzyl)-imidazolidin-4-yl]-ethoxy}-phenyl)-2-methyl-2-phenoxy-propionic acid ethyl ester:** A slurry of 2-Methyl-3-(4-{2-[2-oxo-1-(4-trifluoromethyl-benzyl)-imidazolidin-4-yl]-ethoxy}-phenyl)-2-phenoxy-propionic acid ethyl ester (0.500g, 0.875mmol) and sodium hydride (.039g, 0.96 mmol, 60% dispersion on mineral oil) in DMF (10mL) is stirred for 1hour, then cooled to 0° C. Iodoethane (0.70mL, 8.75mmol,  $d=1.975$ ) is added, then the reaction mixture is stirred

18hours at ambient temperature and diluted with ethyl acetate. The organic layer is washed, dried, and concentrated to provide the title compound. <sup>1</sup>H NMR (400MHz, CDCl<sub>3</sub>): δ 7.55 (d, 2h, J = 8.8 Hz), 7.37 (d, 2H, J = 8.0 Hz), 7.21 (t, 2H, J = 8.8 Hz), 7.15 (d, 2H, J = 8.0 Hz), 6.96 (t, 1H, J = 7.2 Hz), 6.82 (d, 2H, J = 7.2 Hz), 6.74 (d, 2H, J = 8.8 Hz), 4.43, 4.39 (AB<sub>q</sub>, 2H, J = 15.2 Hz), 4.22, 4.17 (AB<sub>q</sub>, 2H, J = 7.2 Hz), 3.96 (t, 1H, J = 6.4 Hz), 3.85-3.78 (m, 1H), 3.61-3.55 (m, 1H), 3.35 (t, 1H, J = 8.8 Hz), 3.26, 3.12 (AB<sub>q</sub>, 2H, J = 13.6 Hz), 3.16-3.07 (m, 2H), 2.99 (t, 1H, J = 8.0 Hz), 2.26-2.19 (m, 1H), 1.90-1.83 (m, 1H), 1.39(s, 3H), 1.21 (t, 3H, J = 7.2 Hz), 1.14 (t, 3H, J = 7.2 Hz). MS [EI+] 599 (M+H)<sup>+</sup>.

15 Step D

**3-(4-{2-[3-Ethyl-2-oxo-1-(4-trifluoromethyl-benzyl)-imidazolidin-4-yl]-ethoxy}-phenyl)-2-methyl-2-phenoxy-propionic acid:** A solution of 3-(4-{2-[3-Ethyl-2-oxo-1-(4-trifluoromethyl-benzyl)-imidazolidin-4-yl]-ethoxy}-phenyl)-2-methyl-2-phenoxy-propionic acid ethyl ester (0.31g, 0.518mmol) and 5N NaOH (1mL) in ethanol (9mL) is refluxed for 1hour, cooled to ambient temperature, and concentrated. The residue is diluted with CH<sub>2</sub>Cl<sub>2</sub>, washed, dried, and concentrated in vacuo to provide the title compound (0.256g, 87%). <sup>1</sup>H NMR (400MHz, CDCl<sub>3</sub>): δ 7.54 (d, 2H, J = 7.6 Hz), 7.35 (d, 2H, J = 8.4 Hz), 7.21-7.17 (m, 4H), 6.97 (t, 1H, J = 7.6 Hz), 6.88 (d, 2H, J = 7.6 Hz), 6.72 (d, 2H, J = 8.4 Hz), 4.42, 4.38 (AB<sub>q</sub>, 2H, J = 15.1 Hz), 3.94 (t, 2H, J = 5.9 Hz), 3.87-3.81 (m, 1H), 3.62-3.53 (m, 1H), 3.36 (t, 1H, J = 8.4 Hz), 3.31, 3.08 (AB<sub>q</sub>, 2H, J = 8.4 Hz), 3.15-3.08 (m, 1H), 3.00 (t, 1H, J = 8.4 Hz), 2.23-2.19 (m, 1H), 1.89-1.82

(m, 1H), 1.37 (s, 3H), 1.13 (t, 3H,  $J = 6.7$  Hz). MS [EI+] 571 (M+H)<sup>+</sup>, [EI-] 569 (M-H)<sup>+</sup>.

#### Example 57

5                    **3-(4-{2-[3-(2-Methoxy-ethyl)-2-oxo-1-(4-trifluoromethyl-benzyl)-imidazolidin-4-yl]-ethoxy}-phenyl)-2-methyl-2-phenoxy-propionic acid**

The title compound is prepared using substantially the procedures described herein, and concentrated *in vacuo* to provide the title compound (0.139g, 88%). <sup>1</sup>H NMR (400MHz, CDCl<sub>3</sub>): δ 7.54 (d, 2H,  $J = 7.5$  Hz), 7.35 (d, 2H,  $J = 8.3$  Hz), 7.22-7.17 (m, 4H), 6.97 (t, 1H,  $J = 7.5$  Hz), 6.88 (d, 2H,  $J = 8.3$  Hz), 6.72 (d, 2H,  $J = 8.3$  Hz), 4.42, 4.40 (AB<sub>q</sub>, 2H,  $J = 15.8$  Hz), 3.95-3.90 (m, 3H), 3.69-3.63 (m, 1H), 15 3.55-3.63 (m, 2H), 3.33 (s, 3H), 3.41-3.23 (m, 3H), 3.11-3.01 (m, 2H), 2.73-2.34 (m, 1H), 1.91-1.87 (m, 1H), 1.38 (s, 3H). MS [EI+] 601 (M+H)<sup>+</sup>, [EI-] 599 (M-H)<sup>+</sup>.

#### Example 58

20                    **2-Butoxy-2-methyl-3-(4-{2-[3-methyl-2-oxo-1-(3-trifluoromethyl-benzyl)-imidazolidin-4-yl]-ethoxy}-phenyl)-propionic acid**

The title compound is prepared using substantially the procedures described herein and purified by flash chromatography (25% acetone in hexanes) to provide the title compound (0.62g, 83%). <sup>1</sup>H NMR (400MHz, CDCl<sub>3</sub>): δ 7.51-7.38 (m, 4H), 7.04 (d, 2H,  $J = 8.8$  Hz), 6.69 (d, 2H,  $J = 8.8$  Hz), 4.46, 4.34 (AB<sub>q</sub>, 2H,  $J = 14.9$  Hz), 3.94 (t, 2H,  $J = 5.7$  Hz), 3.68-3.60 (m, 1H), 3.57-3.52 (m, 1H), 3.48-3.43 (m, 1H), 30 3.38 (t, 1H,  $J = 8.8$  Hz), 2.98, 2.92 (AB<sub>q</sub>, 2H,  $J = 14.4$  Hz), 2.83 (s, 3H), 2.25-2.18 (m, 1H), 1.91-1.83 (m, 1H), 1.59-

1.52 (m, 2H), 1.45 (s, 3H), 1.39-1.30 (m, 2H), 0.91 (t, 3H,  $J = 7.2$  Hz). MS [EI+] 537 (M+H)<sup>+</sup>, [EI-] 535 (M-H)<sup>+</sup>.

#### Example 59

5     **3-{4-[2-(1-Biphenyl-4-ylmethyl-3-methyl-2-oxo-imidazolidin-4-yl)-ethoxy]-phenyl}-2-butoxy-2-methyl-propionic acid**

The title compound is prepared using substantially the procedures described herein and purified by LCMS to provide the title compound (5.5mg, 14% over two steps). <sup>1</sup>H NMR

10    (400MHz, CDCl<sub>3</sub>): δ 7.56-7.51 (m, 4H), 7.41 (t, 2H,  $J = 7.8$  Hz), 7.33-7.28 (m, 3H), 7.03 (d, 2H,  $J = 8.4$  Hz), 6.70 (d, 2H,  $J = 8.4$  Hz), 4.43, 4.35 (AB<sub>q</sub>, 2H,  $J = 15.6$  Hz), 3.94 (t, 2H,  $J = 6.0$  Hz), 3.66-3.59 (m, 1H), 3.47-3.38 (m, 1H), 3.01 (t, 1H,  $J = 7.8$  Hz), 2.97, 2.91 (AB<sub>q</sub>, 2H,  $J = 13.8$  Hz), 2.83  
15    (s, 3H), 2.25-2.18 (m, 1H), 1.91-1.83 (m, 1H), 1.58-1.51 (m, 2H), 1.45 (s, 3H), 1.38-1.29 (m, 2H), 0.89 (t, 3H,  $J = 7.8$  Hz). MS [EI+] 545 (M+H)<sup>+</sup>, [EI-] 543 (M-H)<sup>+</sup>.

#### Example 60

20     **3-{4-[2-(1-Benzyl-3-methyl-2-oxo-imidazolidin-4-yl)-ethoxy]-phenyl}-2-butoxy-2-methyl-propionic acid**

The title compound is prepared using substantially the procedures described herein, and purified by LCMS to provide the title compound (5.5mg, 14% over two steps). <sup>1</sup>H NMR

25    (400MHz, CDCl<sub>3</sub>): δ 7.31-7.21 (m, 5H), 7.04 (d, 2H,  $J = 8.6$  Hz), 6.69 (d, 2H,  $J = 8.6$  Hz), 3.38, 3.32 (AB<sub>q</sub>, 2H,  $J = 15.7$  Hz), 3.92 (t, 2H,  $J = 6.3$  Hz), 3.62-3.53 (m, 2H), 3.48-3.43 (m, 1H), 3.35 (t, 1H,  $J = 8.6$  Hz), 2.99-2.93 (AB<sub>q</sub>, 2H,  $J = 14.1$  Hz), 2.99, 2.93 (AB<sub>q</sub>, 2H,  $J = 14.1$  Hz), 2.95 (t, 2H,  $J = 7.9$  Hz), 2.88 (s, 3H), 2.23-2.17 (m, 1H), 1.88-1.82 (m, 1H), 1.60-1.52 (m, 2H), 1.47 (s, 3H), 1.38-1.32 (m, 2H),  
30



1.23 (t, 1H,  $J = 7.1$  Hz), 0.91 (t, 3H,  $J = 7.1$  Hz). MS  
[EI+] 469 (M+H)<sup>+</sup>, [EI-] 467 (M-H)<sup>+</sup>.

#### Example 61

5           **2-Methoxy-2-methyl-3-(4-{2-[3-methyl-2-oxo-1-(4-  
trifluoromethyl-benzyl)-imidazolidin-4-yl]-ethoxy}-phenyl)-  
propionic acid**

The title compound is prepared using substantially the  
procedures described herein, and purified by flash  
10 chromatography to provide the title compound (48mg, 39% over  
two steps). <sup>1</sup>H NMR (400MHz, CDCl<sub>3</sub>): δ 7.53 (d, 2H,  $J = 8.2$   
Hz), 7.34 (d, 2H,  $J = 8.2$  Hz), 7.07 (d, 2H,  $J = 8.2$  Hz),  
6.64 (d, 2H,  $J = 8.2$  Hz), 4.40, 4.36 (AB<sub>q</sub>, 2H,  $J = 15.5$  Hz),  
3.90-3.87 (m, 2H), 3.62-3.56 (m, 1H), 3.35 (t, 1H,  $J = 8.2$   
15 Hz), 3.26 (s, 3H), 3.04-2.87 (m, 4H), 2.82 (s, 3H), 2.20-  
2.16 (m, 1H), 1.87-1.80 (m, 1H), 1.31 (s, 3H). MS [EI+] 495  
(M+H)<sup>+</sup>, [EI-] 493 (M-H)<sup>+</sup>.


#### Example 62

20           **2-Methoxy-2-methyl-3-(4-{2-[3-methyl-2-oxo-1-(4-  
trifluoromethyl-benzyl)-imidazolidin-4-yl]-ethoxy}-phenyl)-  
propionic acid**

The title compound is prepared using substantially the  
procedures described herein, and purified by flash  
25 chromatography to provide the title compound (88mg, 63% over  
two steps). <sup>1</sup>H NMR (400MHz, CDCl<sub>3</sub>): δ 7.53 (d, 2H,  $J = 8.2$   
Hz), 7.34 (d, 2H,  $J = 8.2$  Hz), 7.07 (d, 2H,  $J = 8.2$  Hz),  
6.64 (d, 2H,  $J = 8.2$  Hz), 4.40, 4.36 (AB<sub>q</sub>, 2H,  $J = 15.5$  Hz),  
3.90-3.87 (m, 2H), 3.62-3.56 (m, 1H), 3.35 (t, 1H,  $J = 8.2$   
30 Hz), 3.26 (s, 3H), 3.04-2.87 (m, 4H), 2.82 (s, 3H), 2.20-  
2.16 (m, 1H), 1.87-1.80 (m, 1H), 1.31 (s, 3H). MS [EI+] 495  
(M+H)<sup>+</sup>, [EI-] 493 (M-H)<sup>+</sup>.

**Example 63**

**2-Methyl-3-(4-{2-[3-methyl-2-oxo-1-(4-trifluoromethyl-benzyl)-imidazolidin-4-yl]-ethoxy}-phenyl)-2-phenoxy-propionic acid**

**Step A**

**3-(4-{2-[3-(4-Methoxy-benzyl)-2-oxo-1-(4-trifluoromethyl-benzyl)-imidazolidin-4-yl]-ethoxy}-phenyl)-2-methyl-2-phenoxy-propionic acid ethyl ester:** Cesium carbonate (2.18g, 6.702mmol) is added to a solution of 3-(4-Hydroxy-phenyl)-2-methyl-2-phenoxy-propionic acid ethyl ester (1.55g, 5.155mmol) and Toluene-4-sulfonic acid 2-[3-methyl-2-oxo-1-(4-trifluoromethyl-benzyl)-imidazolidin-4-yl]-ethyl ester (3.19g, 5.67mmol) in DMF (30mL). The resultant mixture is stirred at 55°C under an atmosphere of nitrogen for 18 h, then diluted with ethyl acetate. The organic layer is washed, then dried, concentrated, and purified by flash chromatography (17% acetone in hexanes) to provide the title compound (3.46g, 97%). <sup>1</sup>H NMR (400MHz, CDCl<sub>3</sub>): δ 7.57 (d, 2H, J = 7.01 Hz), 7.37 (d, 2H, J = 7.1 Hz), 7.23-7.19 (m, 4H), 7.13 (d, 2H, J = 8.7 Hz), 6.97 (t, 1H, J = 7.1 Hz), 6.87-6.80 (m, 4H), 6.67 (d, 2H, J = 8.7 Hz), 4.79, 4.07 (AB<sub>q</sub>, 2H, J = 15.0 Hz), 4.47, 4.43 (AB<sub>q</sub>, 2H, J = 15.0 Hz),

4.20 (q, 2H,  $J = 7.1$  Hz), 3.92-3.87 (m, 2H), 3.80 (s, 3H),  
3.61-3.55 (m, 1H), 3.31 (t, 2H,  $J = 8.7$  Hz), 3.24, 3.10  
(ABq, 2H,  $J = 14.2$  Hz), 3.00 (t, 1H,  $J = 7.9$  Hz), 2.21-2.14  
(m, 1H), 1.87-1.78 (m, 1H), 1.38 (s, 3H), 1.21 (t, 3H,  $J =$   
5 7.1 Hz). MS [EI+] 691 (M+H)<sup>+</sup>.  $R_f=0.49$  in 50% acetone in  
hexanes.

#### Step B

**2-Methyl-3-(4-{2-[2-oxo-1-(4-trifluoromethyl-benzyl)-  
imidazolidin-4-yl]-ethoxy}-phenyl)-2-phenoxy-propionic acid**  
10 **ethyl ester**: Trifluoroacetic acid (60mL) is added dropwise  
to a solution of 3-(4-{2-[3-(4-Methoxy-benzyl)-2-oxo-1-(4-  
trifluoromethyl-benzyl)-imidazolidin-4-yl]-ethoxy}-phenyl)-  
2-methyl-2-phenoxy-propionic acid ethyl ester (3.46g,  
5.01mmol) and triethylsilane (1.6mL, 10.0mmol,  $d=0.728$ ).  
15 The reaction mixture is stirred at ambient temperature for  
2hours, concentrated, and diluted with ethyl acetate. The  
solution is washed with a saturated solution of aqueous  
sodium bicarbonate, water, and brine, then dried over  
 $\text{Na}_2\text{SO}_4$ , and concentrated in vacuo to provide the title  
20 compound.  $^1\text{H}$  NMR (400MHz,  $\text{CDCl}_3$ ):  $\delta$  7.58 (d, 2H,  $J = 8.2$   
Hz), 7.38 (d, 2H,  $J = 8.2$  Hz), 7.23-7.14 (m, 4H), 6.97 (t,  
1H,  $J = 7.4$  Hz), 6.82 (d, 2H,  $J = 7.4$  Hz), 6.78 (d, 2H,  $J =$   
7.4 Hz), 4.44, 4.38 (ABq, 2H,  $J = 15.6$  Hz), 4.20 (q, 2H,  $J =$   
7.4 Hz), 4.07-3.99 (m, 2H), 3.96-3.89 (m, 1H), 3.47 (t, 1H,  
25  $J = 8.2$  Hz), 3.27, 3.09 (ABq, 2H,  $J = 14.1$  Hz), 3.04, 3.02  
(ABq, 1H,  $J = 6.7$  Hz), 2.08-1.99 (m, 1H), 1.98-1.92 (m, 1H),  
1.38 (s, 3H), 1.21 (t, 3H,  $J = 6.7$  Hz). MS [EI+] 571  
(M+H)<sup>+</sup>, [EI-] 569 (M-H)<sup>+</sup>.  $R_f=0.07$  in 33% acetone in hexanes.

30

#### Step C

**2-Methyl-3-(4-{2-[3-methyl-2-oxo-1-(4-trifluoromethyl-  
benzyl)-imidazolidin-4-yl]-ethoxy}-phenyl)-2-phenoxy-**

**propionic acid ethyl ester:** A slurry of 2-Methyl-3-(4-{2-[2-oxo-1-(4-trifluoromethyl-benzyl)-imidazolidin-4-yl]-ethoxy}-phenyl)-2-phenoxy-propionic acid ethyl ester (0.544g, 0.953mmol) and sodium hydride (.042g, 1.05 mmol, 60% dispersion on mineral oil) in DMF (10mL) was stirred for 1h, then cooled to 0° C. Iodomethane (0.60mL, 9.53mmol, d=2.28) was added, then the reaction mixture is stirred 18hours at ambient temperature and diluted with ethyl acetate. The organic layer is washed, dried, concentrated, and purified by flash chromatography to provide the title compound. <sup>1</sup>H NMR (400MHz, CDCl<sub>3</sub>): δ 7.56 (d, 2H, J = 8.0 Hz), 7.37 (d, 2H, J = 8.0 Hz), 7.26-7.19 (m, 2H), 7.15 (d, 2H, J = 8.9 Hz), 6.97 (t, 1H, J = 8.0 Hz), 6.82 (d, 2H, J = 8.0 Hz), 6.74 (d, 2H, J = 8.9 Hz), 4.46, 4.38 (AB<sub>q</sub>, 2H, J = 15.1 Hz), 4.20 (q, 2H, J = 7.1 Hz), 3.97 (t, 2H, J = 7.1 Hz), 3.67-3.59 (m, 1H), 3.78 (t, 1H, J = 8.9 Hz), 3.26, 3.10 (AB<sub>q</sub>, 2H, J = 14.2 Hz), 2.99 (t, 1H, J = 8.9 Hz), 2.85 (s, 3H), 2.28-2.21 (m, 1H), 1.93-1.84 (m, 1H), 1.39 (s, 3H), 1.21 (t, 3H, J = 7.1 Hz). MS [EI+] 585 (M+H)<sup>+</sup>. R<sub>f</sub>=0.35 in 50% acetone in hexanes.

#### Step D

**2-Methyl-3-(4-{2-[3-methyl-2-oxo-1-(4-trifluoromethyl-benzyl)-imidazolidin-4-yl]-ethoxy}-phenyl)-2-phenoxy-propionic acid:** A solution of 2-methyl-3-(4-{2-[3-methyl-2-oxo-1-(4-trifluoromethyl-benzyl)-imidazolidin-4-yl]-ethoxy}-phenyl)-2-phenoxy-propionic acid ethyl ester and 5N NaOH (0.7mL) in ethanol (6mL) is refluxed for 1h, cooled to ambient temperature, and concentrated *in vacuo*. The residue is diluted with CH<sub>2</sub>Cl<sub>2</sub>, washed, dried, and concentrated to provide the title compound (0.186g, 100%). <sup>1</sup>H NMR (400MHz, CDCl<sub>3</sub>): δ 7.54 (d, 2H, J = 8.3 Hz), 7.35 (d, 2H, J = 8.3

Hz), 7.24-7.17 (m, 4H), 7.00 (t, 1H,  $J = 7.4$  Hz), 6.89 (d, 2H,  $J = 7.4$  Hz), 6.73 (d, 2H,  $J = 8.3$  Hz), 4.45, 4.37 (AB<sub>q</sub>, 2H,  $J = 15.7$  Hz), 3.96 (t, 2H,  $J = 5.5$  Hz), 3.68-3.62 (m, 1H), 3.38 (t, 1H,  $J = 8.3$  Hz), 3.32, 3.10 (AB<sub>q</sub>, 2H,  $J = 12.9$  5 Hz), 3.00 (t, 1H,  $J = 8.3$  Hz), 2.84 (s, 3H), 2.27-2.19 (m, 1H), 1.92-1.85 (m, 1H), 1.39 (s, 3H). MS [EI+] 557 (M+H)<sup>+</sup>, [EI-] 555 (M-H)<sup>+</sup>.

#### Example 64

10 **2,2-Dimethyl-3-(4-{2-[3-methyl-2-oxo-1-(4-trifluoromethyl-benzyl)-imidazolidin-4-yl]-ethoxy}-phenyl)-propionic acid**


##### Step A

15 **2,2-Dimethyl-3-(4-{2-[3-methyl-2-oxo-1-(4-trifluoromethyl-benzyl)-imidazolidin-4-yl]-ethoxy}-phenyl)-propionic acid methyl ester**

To a solution of 3-(4-Hydroxy-phenyl)-2,2-dimethyl-propionic acid methyl ester (31 mg, 0.15 mmol) and Toluene-4-sulfonic acid 2-[3-methyl-2-oxo-1-(4-trifluoromethyl-benzyl)-imidazolidin-4-yl]-ethyl ester (75 mg, 0.16 mmol) in 20 DMF (1 ml) is added Cs<sub>2</sub>CO<sub>3</sub> (64 mg, 0.19 mmol), and the mixture heated at 60° C for 15 hours. The mixture is cooled, diluted with water (25 ml) and extracted with ethyl acetate (2 x 25 ml). The combined organic extracts were washed with 25 brine (25 ml), dried (Na<sub>2</sub>SO<sub>4</sub>), and concentrated to an oil. Purification by flash chromatography on a Biotage silica cartridge (gradient elution, 2:1 hexanes:ethyl acetate to 1:4 hexanes:ethyl acetate) produced an oil (69 mg, 94%). MS [EI+] 493 (M+H)<sup>+</sup>. Structure confirmed by 1H-NMR.

Step B**2,2-Dimethyl-3-(4-{2-[3-methyl-2-oxo-1-(4-trifluoromethyl-benzyl)-imidazolidin-4-yl]-ethoxy}-phenyl)-propionic acid**


To a solution of the ester obtained from Step A (66 mg, 0.13 mmol) in methanol (1.5 ml) is added 5N NaOH (0.3 ml, 1.5 mmol), and the mixture stirred for 20 hours at ambient temperature. The mixture is poured into 1N HCl (20 ml), then extracted with ethyl acetate (2 x 15 ml). The combined ethyl acetate extracts were washed with water, brine, dried (Na<sub>2</sub>SO<sub>4</sub>), and concentrated to a foam (62 mg, 97%). MS [EI+] 479 (M+H)<sup>+</sup>. Structure confirmed by 1H-NMR.

**Example 65****2-methyl-3-(4-{2-[1-(4-methyl-benzyl)-2-oxo-3-propyl-imidazolidin-4-yl]-ethoxy}-phenyl)-2-phenoxy-propionic acid**Step A

A solution of (2,5-dioxo-imidazolidin-4-yl)-acetic acid, methyl ester (2.35 g, 13.7 mmol) in DMF (100 mL) is treated with p-methyl benzyl bromide (2.90 g, 15.7 mmol), MgSO<sub>4</sub> (3.30 g, 27.4 mmol) and then K<sub>2</sub>CO<sub>3</sub> (3.77 g, 27.3 mmol). The resultant mixture is stirred under N<sub>2</sub> in an ice bath for 30 minutes and then warmed to room temperature for 16 h. The reaction mixture is then filtered, and then aqueous 1N HCl (150 mL) is added to the filtrate. The filtrate is extracted with EtOAc and the organic layer dried. The solvent is removed to give 4.11 g crude product which is purified by flash chromatography using 100% CH<sub>2</sub>Cl<sub>2</sub> then 97.5:2.5 CH<sub>2</sub>Cl<sub>2</sub>:MeOH to give 3.67 g (97%) [1-(4-methyl-benzyl)-2,5-dioxo-imidazolidin-4-yl]-acetic acid methyl


ester.  $^1\text{H}$  NMR. MS ( $\text{ES}^+$ ) Calc'd for  $\text{C}_{14}\text{H}_{17}\text{N}_2\text{O}_4$  ( $\text{M} + 1$ ) 277.  
Found  $m/z$  277 (100%).

Step B



5 A 0 °C solution of compound [1-(4-methyl-benzyl)-2,5-dioxo-imidazolidin-4-yl]-acetic acid methyl ester (1.60 g, 5.8 mmol) in DMF (28 mL) is treated with sodium hydride (60% dispersion, 0.25 g, 6.4 mmol) and warmed to room temperature and stirred under  $\text{N}_2$  for 15 minutes. The resultant mixture  
10 is cooled to 0 °C and then treated with 1-propyl iodide (1.08 g, 6.4 mmol) and then warmed to room temperature and stirred for 30 minutes. The reaction is quenched with aqueous 1 N HCl (28 mL) and then worked up extractively with EtOAc and water. The organic layer is dried and the  
15 solvent removed to give crude product that is purified by flash chromatography using a gradient of 10:1 to 1:1 hexanes:EtOAc to afford 1.40 g (76%) [1-(4-methyl-benzyl)-2,5-dioxo-3-propyl-imidazolidin-4-yl]-acetic acid methyl ester.  $^1\text{H}$  NMR. MS ( $\text{ES}^+$ ) Calc'd for  $\text{C}_{17}\text{H}_{23}\text{N}_2\text{O}_4$  ( $\text{M} + 1$ ) 319.  
20 Found  $m/z$  319 (100%).


Step C



A solution of [1-(4-methyl-benzyl)-2,5-dioxo-3-propyl-imidazolidin-4-yl]-acetic acid methyl ester (1.67 g, 5.20  
25 mmol) in methanol (40 mL) is treated with aqueous 5 N NaOH (4.4 mL) and heated to reflux 1.5 hour. The reaction


mixture is cooled, the solvent removed. The resultant residue is acidified and extracted with EtOAc. The organic layer is dried and dried to afford 1.56 g (98%) [1-(4-methyl-benzyl)-2,5-dioxo-3-propyl-imidazolidin-4-yl]-acetic acid. <sup>1</sup>H NMR. MS (ES<sup>-</sup>) Calc'd for C<sub>16</sub>H<sub>19</sub>N<sub>2</sub>O<sub>4</sub> (M - 1) 303. Found m/z 303 (100%).

#### Step D



10 A solution of [1-(4-methyl-benzyl)-2,5-dioxo-3-propyl-imidazolidin-4-yl]-acetic acid (3.23 g, 10.6 mmol) in THF (50 mL) is treated dropwise with 1 M solution of borane-THF complex in THF (53.1 mL, 53.1 mmol) and then stirred at room temperature under N<sub>2</sub> for 4 hours. The reaction is quenched  
15 with methanol (30 mL) and stirred at room temperature for 16 hour. The solvent is removed to give crude product that is purified by flash chromatography using 97.5/2.5 CH<sub>2</sub>Cl<sub>2</sub>:MeOH to afford 2.46 g (84%) 4-(2-hydroxy-ethyl)-1-(4-methyl-benzyl)-3-propyl-imidazolidin-2-one. <sup>1</sup>H NMR. MS (ES<sup>+</sup>)  
20 Calc'd for C<sub>16</sub>H<sub>25</sub>N<sub>2</sub>O<sub>2</sub> (M +1) 277. Found m/z 277 (100%).

#### Step E




A solution of 4-(2-hydroxy-ethyl)-1-(4-methyl-benzyl)-3-propyl-imidazolidin-2-one (2.46 g, 8.90 mmol), pyridine  
25 (2.46 g, 31.1 mmol) and 4-dimethyl amino pyridine (0.33 g, 2.70 mmol) in CH<sub>2</sub>Cl<sub>2</sub> is treated with *p*-toluenesulfonic anhydride (4.65 g, 14.2 mmol) and the reaction stirred at room temperature for under N<sub>2</sub> for 1.5 hours. The reaction




mixture is washed with aqueous 0.5 N HCl (100 mL), the organic layer is dried, and the solvent removed to afford crude product that is purified by flash chromatography using 100% CH<sub>2</sub>Cl<sub>2</sub> then 97.5:2.5 CH<sub>2</sub>Cl<sub>2</sub>:MeOH to afford 3.31 g (86%)  
5 toluene-4-sulfonic acid 2-[1-(4-methyl-benzyl)-2-oxo-3-propyl-imidazolidin-4-yl]-ethyl ester. <sup>1</sup>H NMR. MS (ES<sup>+</sup>) Calc'd for C<sub>23</sub>H<sub>31</sub>N<sub>2</sub>O<sub>4</sub> (M +1) 431. Found m/z 431 (100%).

Step F




10 A mixture of 3-(4-hydroxy-phenyl)-2-methyl-2-phenoxy-propionic acid ethyl ester (0.80 g, 2.66 mmol), toluene-4-sulfonic acid 2-[1-(4-methyl-benzyl)-2-oxo-3-propyl-imidazolidin-4-yl]-ethyl ester (1.26 g, 2.93 mmol) and Cs<sub>2</sub>CO<sub>3</sub> (1.04 g, 3.19 mmol) in DMF (40 mL) is heated to 55 °C  
15 under N<sub>2</sub> for 17 h. The reaction is cooled and quenched with 1 N HCl (20 mL) and worked up extractively with Et<sub>2</sub>O and water. The organic layer is dried and the solvent removed to afford crude product that is purified by flash chromatography using 7:1 hexanes:acetone to afford 1.02 g  
20 (68%) 2-methyl-3-(4-{2-[1-(4-methyl-benzyl)-2-oxo-3-propyl-imidazolidin-4-yl]-ethoxy}-phenyl)-2-phenoxy-propionic acid ethyl ester. The diastereomeric products were separated by chiral HPLC (90:10 heptane:IPA, 280 mL/min, 220 nm,) to give products in >99% de. R<sub>f</sub> = 0.49 (1:1 acetone:hexanes). <sup>1</sup>H  
25 NMR. MS (ES<sup>+</sup>) Calc'd for C<sub>34</sub>H<sub>43</sub>N<sub>2</sub>O<sub>5</sub> (M +1) 559. Found m/z 559 (100%).

Step G

A solution of 2-methyl-3-(4-{2-[1-(4-methyl-benzyl)-2-oxo-3-propyl-imidazolidin-4-yl]-ethoxy}-phenyl)-2-phenoxy-propionic acid ethyl ester (0.456 g, 0.816 mmol) in ethanol (20 mL) is treated with aqueous 5 N NaOH (2 mL) and heated to reflux 1 h. The reaction mixture is cooled, the solvent removed. The resultant residue is acidified and extracted with CH<sub>2</sub>Cl<sub>2</sub>. The organic layer is dried and the solvent removed to afford 0.419 g (97%) 2-methyl-3-(4-{2-[1-(4-methyl-benzyl)-2-oxo-3-propyl-imidazolidin-4-yl]-ethoxy}-phenyl)-2-phenoxy-propionic acid. <sup>1</sup>H NMR. HRMS (ES<sup>+</sup>) *m/z* exact mass calc'd for C<sub>32</sub>H<sub>39</sub>N<sub>2</sub>O<sub>5</sub> (M + 1) 531.2859. Found *m/z* 531.2866.

**Example 66**

**Preparation of 2-methyl-3-{4-[2-(3-methyl-1-naphthalen-2-yl)methyl-2-oxo-imidazolidin-4-yl]-ethoxy}-phenyl}-2-phenoxy-propionic acid**


Step A

A solution of (2,5-dioxo-imidazolidin-4-yl)-acetic acid methyl ester (6.48 g, 37.6 mmol) in DMF (250 mL) is treated with 2-(bromomethyl)-naphthalene (9.15 g, 41.4 mmol), MgSO<sub>4</sub> (4.53 g, 37.6 mmol) and then K<sub>2</sub>CO<sub>3</sub> (15.61 g, 0.113 mol) at 0 °C. The resultant mixture is warmed to room temperature and stirred under N<sub>2</sub> for 16 hours. The reaction mixture is


filtered, and then aqueous 1N HCl (300 mL) is added to the filtrate. The filtrate is extracted with Et<sub>2</sub>O and the organic layer dried. The solvent is removed to give a crude oil which is purified by flash chromatography using 2:1

- 5 hexanes:acetone to afford 6.27 g of (1-naphthalen-2-ylmethyl-2,5-dioxo-imidazolidin-4-yl)-acetic acid methyl ester (53%).  $R_f = 0.32$  (1:1 hexanes:acetone). <sup>1</sup>H NMR.


Step B



- 10 A 0 °C solution of compound (1-naphthalen-2-ylmethyl-2,5-dioxo-imidazolidin-4-yl)-acetic acid methyl ester (6.29 g, 20.1 mmol) in DMF (60 mL) is treated with sodium hydride (60% dispersion, 0.97 g, 24.3 mmol) and warmed to room temperature and stirred under N<sub>2</sub> for 20 minutes. The
- 15 resultant mixture is cooled to 0 °C and then treated with 4-methoxybenzyl chloride (6.35 g, 40.6 mmol) and then warmed to room temperature and stirred for 16 hours. The reaction is quenched with aqueous 1 N HCl (100 mL) and then worked up extractively with Et<sub>2</sub>O and water. The organic layer is
- 20 dried and the solvent removed to give crude product that is purified by flash chromatography using a gradient of 4:1 hexanes:acetone to afford 8.31 g (95%) [3-(4-methoxybenzyl)-1-naphthalen-2-ylmethyl-2,5-dioxo-imidazolidin-4-yl]-acetic acid methyl ester.  $R_f = 0.44$  (1:1
- 25 hexanes:acetone). <sup>1</sup>H NMR.


Step C

A solution of [3-(4-methoxy-benzyl)-1-naphthalen-2-ylmethyl-2,5-dioxo-imidazolidin-4-yl]-acetic acid methyl ester (8.31 g, 19.2 mmol) in methanol (100 mL) is treated with aqueous 5 N NaOH (40 mL) and heated to reflux 1 hour. The reaction mixture is cooled, the solvent removed in vacuo. The resultant residue is acidified with aqueous 1 N HCl (300 mL) and extracted with Et<sub>2</sub>O and water. The organic layer is dried and the solvent removed in vacuo to afford 8.08 g (100%) acid that is utilized without purification. A solution of crude acid (8.08 g, assume 19.2 mmol) in THF (100 mL) is treated dropwise with 1 M solution of borane-THF complex in THF (116.0 mL, 0.116 mol) and then stirred at room temperature under N<sub>2</sub> for 16 hours. The reaction is quenched with methanol (100 mL) and stirred at room temperature for 1 hour. The solvent is removed to give crude product that is purified by flash chromatography using a gradient of 3:1 to 2:1 hexanes:acetone to afford 5.10 g (68%) 4-(2-hydroxy-ethyl)-3-(4-methoxy-benzyl)-1-naphthalen-2-ylmethyl-imidazolidin-2-one. <sup>1</sup>H NMR. MS (ES<sup>+</sup>) Calc'd for C<sub>24</sub>H<sub>27</sub>N<sub>2</sub>O<sub>3</sub> (M +1) 391. Found m/z 391 (100%).

Step D

A solution of 4-(2-hydroxy-ethyl)-3-(4-methoxy-benzyl)-1-naphthalen-2-ylmethyl-imidazolidin-2-one (5.10 g, 13.1 mmol), pyridine (3.62 g, 45.7 mmol) and 4-dimethyl amino pyridine (0.48 g, 3.92 mmol) in  $\text{CH}_2\text{Cl}_2$  (100 mL) is treated with *p*-toluenesulfonic anhydride (6.82 g, 20.9 mmol) and the reaction stirred at room temperature for under  $\text{N}_2$  for 1.5 h. The reaction mixture is washed with aqueous 1 N HCl (100 mL), the organic layer was dried, and the solvent removed in vacuo to afford crude product that is purified by flash chromatography using 3:1 hexanes:acetone to afford 6.74 g (95%) toluene-4-sulfonic acid 2-[3-(4-methoxy-benzyl)-1-naphthalen-2-ylmethyl-2-oxo-imidazolidin-4-yl]-ethyl ester.  $^1\text{H}$  NMR. MS ( $\text{ES}^+$ ) Calc'd for  $\text{C}_{31}\text{H}_{33}\text{N}_2\text{O}_5$  ( $M + 1$ ) 545. Found  $m/z$  545 (100%).


#### Step E



A mixture of 3-(4-hydroxy-phenyl)-2-methyl-2-phenoxy-propionic acid ethyl ester (0.231 g, 0.735 mmol), 2-[3-(4-methoxy-benzyl)-1-naphthalen-2-ylmethyl-2-oxo-imidazolidin-4-yl]-ethyl ester (1.74 g, 3.19 mmol) and  $\text{Cs}_2\text{CO}_3$  (1.14 g, 3.5 mmol) in DMF (40 mL) is heated to 65 °C under  $\text{N}_2$  for 17 h. The reaction is cooled and quenched with 1 N HCl (10 mL) and worked up extractively with  $\text{Et}_2\text{O}$  and water. The organic layer is dried and the solvent removed to afford crude product that is purified by flash chromatography using 8:1 then 7:1 hexanes:acetone to afford 1.53 g (78%) 3-(4-{2-[3-(4-methoxy-benzyl)-1-naphthalen-2-ylmethyl-2-oxo-imidazolidin-4-yl]-ethoxy}-phenyl)-2-methyl-2-phenoxy-propionic acid ethyl ester.  $R_f = 0.43$  (1:1


acetone:hexanes).  $^1\text{H}$  NMR. MS ( $\text{ES}^+$ ) Calc'd for  $\text{C}_{42}\text{H}_{45}\text{N}_2\text{O}_6$  ( $\text{M} + 1$ ) 673. Found  $m/z$  673 (100%).

#### Step F



5 A solution of 3-(4-{2-[3-(4-methoxy-benzyl)-1-naphthalen-2-ylmethyl-2-oxo-imidazolidin-4-yl]-ethoxy}-phenyl)-2-methyl-2-phenoxy-propionic acid ethyl ester (1.54 g, 2.29 mmol) and triethylsilane (0.53 g, 4.56 mmol) in trifluoroacetic acid (40 mL) is stirred at room temperature  
10 under  $\text{N}_2$  for 4 hours. The reaction mixture is diluted with water and extracted with EtOAc. The organic layer is dried and the solvent removed *in vacuo* to afford crude product that is purified by flash chromatography using 3:1 then 1:1 hexanes:acetone to afford 1.25 g (100%) 2-methyl-3-{4-[2-(1-naphthalen-2-ylmethyl-2-oxo-imidazolidin-4-yl)-ethoxy]-phenyl}-2-phenoxy-propionic acid ethyl ester. The  
15 diastereomeric mixture could be separated by chiral HPLC (2:3 IPA:heptane mobile phase, 14 mL/min, 225 nm) to afford >99% de 2-methyl-3-{4-[2-(1-naphthalen-2-ylmethyl-2-oxo-imidazolidin-4-yl)-ethoxy]-phenyl}-2-phenoxy-propionic acid  
20 ethyl esters.  $R_f$  = 0.19 (1:1 acetone:hexanes).  $^1\text{H}$  NMR. MS ( $\text{ES}^+$ ) Calc'd for  $\text{C}_{34}\text{H}_{37}\text{N}_2\text{O}_5$  ( $\text{M} + 1$ ) 553. Found  $m/z$  553 (100%).

#### Step G




25

A solution of a diasteriomeric mixture of 2-methyl-3-{4-[2-(1-naphthalen-2-ylmethyl-2-oxo-imidazolidin-4-yl)-

ethoxy]-phenyl}-2-phenoxy-propionic acid ethyl ester (0.275 g, 0.498 mmol) in DMF (7 mL) is treated with NaH (60% oil suspension, 0.030 g, 0.750 mmol) and stirred at room temperature under N<sub>2</sub> for 30 minutes. The reaction is cooled to 0 °C and treated with iodomethane (0.141 g, 0.996 mmol) and then warmed to room temperature and stirred for 3 hours. The reaction is quenched with 1 N HCl (5 mL) and worked up extractively with Et<sub>2</sub>O and water. The organic layer is dried and the solvent removed *in vacuo* to afford crude product that is purified by flash chromatography using 4:1 hexanes:acetone to afford 0.268 g (95%) 2-methyl-3-{4-[2-(3-methyl-1-naphthalen-2-ylmethyl-2-oxo-imidazolidin-4-yl)-ethoxy]-phenyl}-2-phenoxy-propionic acid ethyl ester. The diastereomeric products are separated by chiral HPLC (8 x 32 cm Chiralpak AD, 100% IPA, 350 mL/min, 270 nm, to give products in >99% de <sup>1</sup>H NMR. R<sub>f</sub> = 0.35 (1:1 acetone:hexanes). MS (ES<sup>+</sup>) Calc'd for C<sub>35</sub>H<sub>39</sub>N<sub>2</sub>O<sub>5</sub> (M +1) 567. Found m/z 567 (100%).

#### Step H



20

A solution of 2-methyl-3-{4-[2-(3-methyl-1-naphthalen-2-ylmethyl-2-oxo-imidazolidin-4-yl)-ethoxy]-phenyl}-2-phenoxy-propionic acid ethyl ester (0.161 g, 0.284 mmol) in ethanol (8 mL) is treated with aqueous 5 N NaOH (0.75 mL) and heated to reflux 1 h. The reaction mixture is cooled, the solvent removed *in vacuo*. The resultant residue is acidified with aqueous 1 N HCl (10 mL) and extracted with CH<sub>2</sub>Cl<sub>2</sub>. The organic layer is dried (Na<sub>2</sub>SO<sub>4</sub>) and the solvent removed *in vacuo* to afford 0.113 g (74%) 2-methyl-3-{4-[2-


25

(3-methyl-1-naphthalen-2-ylmethyl-2-oxo-imidazolidin-4-yl)-ethoxy]-phenyl}-2-phenoxy-propionic acid.  $^1\text{H}$  NMR. HRMS ( $\text{ES}^+$ )  $m/z$  exact mass calc'd for  $\text{C}_{33}\text{H}_{35}\text{N}_2\text{O}_5$  ( $M + 1$ ) 539.2546. Found  $m/z$  539.2548.

5

**Example 67**

**2-methyl-3-{4-[2-(1-naphthalen-2-ylmethyl-2-oxo-imidazolidin-4-yl)-ethoxy]-phenyl}-2-phenoxy-propionic acid**



10 A solution of 2-methyl-3-{4-[2-(1-naphthalen-2-ylmethyl-2-oxo-imidazolidin-4-yl)-ethoxy]-phenyl}-2-phenoxy-propionic acid ethyl esters (0.021 g, 0.040 mmol) in ethanol (8 mL) was treated with aqueous 5 N NaOH (1 mL) and heated to reflux 1 h. The reaction mixture was cooled, the solvent removed *in vacuo*. The resultant residue was acidified with  
15 aqueous 1 N HCl (10 mL) and extracted with  $\text{CH}_2\text{Cl}_2$ . The organic layer was dried ( $\text{Na}_2\text{SO}_4$ ) and the solvent removed *in vacuo* to afford 0.021 g (100%) 2-methyl-3-{4-[2-(1-naphthalen-2-ylmethyl-2-oxo-imidazolidin-4-yl)-ethoxy]-  
20 phenyl}-2-phenoxy-propionic acid.  $^1\text{H}$  NMR. HRMS ( $\text{ES}^+$ )  $m/z$  exact mass calc'd for  $\text{C}_{32}\text{H}_{33}\text{N}_2\text{O}_5$  ( $M + 1$ ) 525.2389. Found  $m/z$  525.2382.

**Example 68**

25 **3-(4-{2-[1-(3-methoxy-benzyl)-3-methyl-2-oxo-imidazolidin-4-yl]-ethoxy}-phenyl)-2-methyl-2-phenoxy-propionic acid**



Step A

A solution of 2-methyl-3-(4-[2-(3-methyl-2-oxo-imidazolidin-4-yl)-ethoxy]-phenyl)-2-phenoxy-propionic acid ethyl ester (0.40 g, 0.94 mmol) in DMF (10 mL) is treated with NaH (60% oil suspension, 0.094 g, 2.35 mmol) and stirred at room temperature under N<sub>2</sub> for 30 minutes. The reaction is cooled to 0 °C and treated with tetrabutylammonium iodide (0.040 g, 0.11 mmol) and 3-methoxybenzyl bromide (0.29 g, 1.43 mmol) and then warmed to room temperature and stirred for 1.5 h. The reaction is quenched with 1 N HCl (15 mL) and worked up extractively with Et<sub>2</sub>O and water. The organic layer is dried (MgSO<sub>4</sub>) and the solvent removed *in vacuo* to afford crude product that is purified by flash chromatography using 2:1 hexanes:acetone to afford 0.28 g (55%) 3-(4-{2-[1-(3-methoxy-benzyl)-3-methyl-2-oxo-imidazolidin-4-yl]-ethoxy}-phenyl)-2-methyl-2-phenoxy-propionic acid ethyl ester. <sup>1</sup>H NMR. R<sub>f</sub> = 0.41 (1:1 acetone:hexanes). MS (ES<sup>+</sup>) Calc'd for C<sub>32</sub>H<sub>39</sub>N<sub>2</sub>O<sub>6</sub> (M +1) 547. Found *m/z* 547 (100%).

Step B


A solution of 3-(4-{2-[1-(3-methoxy-benzyl)-3-methyl-2-oxo-imidazolidin-4-yl]-ethoxy}-phenyl)-2-methyl-2-phenoxy-propionic acid ethyl ester (0.28 g, 0.51 mmol) in ethanol (20 mL) is treated with aqueous 5 N NaOH (2 mL) and heated to reflux 1 h. The reaction mixture is cooled, the solvent removed *in vacuo*. The resultant residue is acidified with aqueous 1 N HCl (20 mL) and extracted with CH<sub>2</sub>Cl<sub>2</sub>. The organic layer is dried (Na<sub>2</sub>SO<sub>4</sub>) and the solvent removed *in vacuo* to afford 0.257 g (97%) 3-(4-{2-[1-(3-methoxy-benzyl)-3-methyl-2-oxo-imidazolidin-4-yl]-ethoxy}-phenyl)-2-methyl-2-phenoxy-propionic acid <sup>1</sup>H NMR (500 MHz, CDCl<sub>3</sub>)  $\delta$  7.28-7.20 (m, 3H), 7.18 (d, 2H, *J* = 8.80 Hz), 7.04 (t, 1H, *J* = 7.34

Hz), 6.91 (d, 2H,  $J = 7.83$  Hz), 6.82 (t, 1H,  $J = 7.83$  Hz), 6.79-6.78 (m, 2H), 6.73 (d, 2H,  $J = 8.83$  Hz), 4.38, 4.28 (AB<sub>q</sub>, 2H,  $J = 14.92$  Hz), 3.95 (t, 2H,  $J = 5.87$  Hz), 3.77 (s, 3H), 3.60 (ddd, 1H,  $J = 12.2$  Hz,  $J = 8.40$  Hz,  $J = 3.42$  Hz), 3.36 (t, 1H,  $J = 7.83$  Hz), 3.30, 3.10 (AB<sub>q</sub>, 2H,  $J = 13.69$  Hz), 2.96 (t, 1H,  $J = 7.83$  Hz), 2.83 (s, 3H), 2.25-2.18 (m, 1H), 1.92-1.84 (m, 1H), 1.42 (s, 3H). HRMS (ES<sup>+</sup>)  $m/z$  exact mass calc'd for C<sub>30</sub>H<sub>35</sub>N<sub>2</sub>O<sub>6</sub> (M + 1) 519.2495. Found  $m/z$  519.2504.

10

**Example 69**

**2-methyl-3-(4-{2-[3-methyl-2-oxo-1-(2-trifluoromethyl-benzyl)-imidazolidin-4-yl]-ethoxy}-phenyl)-2-phenoxy-propionic acid**



15


A solution of 2-methyl-3-{4-[2-(3-methyl-2-oxo-imidazolidin-4-yl)-ethoxy]-phenyl}-2-phenoxy-propionic acid ethyl ester (0.109 g, 0.255 mmol) in DMF (7 mL) is treated with NaH (60% oil suspension, 0.021 g, 0.525 mmol) and stirred at room temperature under N<sub>2</sub> for 20 minutes. The reaction is treated with 2-trifluoromethyl benzyl bromide (0.092 g, 0.385 mmol) and then warmed to room temperature and stirred for 3 hours. The reaction is quenched with acid and worked up extractively with Et<sub>2</sub>O and water. The organic layer is dried and the solvent removed to afford crude product that is dissolved in ethanol (20 mL) is treated with aqueous 5 N NaOH (1.5 mL) and heated to reflux 1 hour. The reaction mixture is cooled, the solvent removed. The resultant residue is acidified with aqueous 1 N HCl (20 mL)

and extracted with CH<sub>2</sub>Cl<sub>2</sub>. The organic layer is dried and the solvent removed *in vacuo* to afford crude acid that is purified by preparative HPLC to give 0.096 g (68%) of 2-methyl-3-(4-{2-[3-methyl-2-oxo-1-(2-trifluoromethyl-benzyl)-imidazolidin-4-yl]-ethoxy}-phenyl)-2-phenoxy-propionic acid

<sup>1</sup>H NMR. HRMS (ES<sup>+</sup>) *m/z* exact mass calc'd for C<sub>30</sub>H<sub>32</sub>N<sub>2</sub>O<sub>5</sub>F<sub>3</sub> (M +1) 557.2263. Found *m/z* 557.2274.

#### Example 70


10 3-(4-{2-[1-(2-methoxy-benzyl)-3-methyl-2-oxo-imidazolidin-4-yl]-ethoxy}-phenyl)-2-methyl-2-phenoxy-propionic acid



A solution of 2-methyl-3-{4-[2-(3-methyl-2-oxo-imidazolidin-4-yl)-ethoxy]-phenyl}-2-phenoxy-propionic acid ethyl ester (0.12 g, 0.281 mmol) in DMF (7 mL) is treated with NaH (60% oil suspension, 0.023 g, 0.575 mmol) and stirred at room temperature under N<sub>2</sub> for 20 minutes. The reaction is treated with 2-methoxybenzyl chloride (0.066 g, 0.421 mmol) and then warmed to room temperature and stirred for 3 h. The reaction is quenched with 1 N HCl (10 mL) and worked up extractively with Et<sub>2</sub>O and water. The organic layer is dried (MgSO<sub>4</sub>) and the solvent removed *in vacuo* to afford crude product that is dissolved in ethanol (20 mL) is treated with aqueous 5 N NaOH (1.5 mL) and heated to reflux 1 h. The reaction mixture is cooled, the solvent removed *in vacuo*. The resultant residue is acidified with aqueous 1 N HCl (20 mL) and extracted with CH<sub>2</sub>Cl<sub>2</sub>. The organic layer is dried (Na<sub>2</sub>SO<sub>4</sub>) and the solvent removed *in vacuo* to afford crude acid that is purified by preparative HPLC to give

0.077 g (53%) of 3-(4-{2-[1-(2-methoxy-benzyl)-3-methyl-2-oxo-imidazolidin-4-yl]-ethoxy}-phenyl)-2-methyl-2-phenoxy-propionic acid.  $^1\text{H}$  NMR. HRMS ( $\text{ES}^+$ )  $m/z$  exact mass calc'd for  $\text{C}_{30}\text{H}_{35}\text{N}_2\text{O}_6$  ( $M + 1$ ) 519.2495. Found  $m/z$  519.2515.

5

**Example 71**

**2-Methyl-3-(4-{2-[3-methyl-2-oxo-1-(4-trifluoromethylbenzyl)-imidazolidin-4-yl]-ethoxy}-phenyl)-2-phenoxy-propionic acid**

10


To a mixture of 54 mg of 2-Methyl-3-(4-[2-(3-methyl-2-oxo-imidazolidin-4-yl)-ethoxy]-phenyl)-2-phenoxy-propionic acid ethyl ester in 2 mL of dry DMF at 0 °C under an atmosphere of nitrogen, 10 mg of NaH (0.253 mmol) is added. The resulting solution is allowed to stand at ambient temperature for 20 min. Then 61 mg of 4-trifluoromethylbenzyl bromide is added and resulting mixture is allowed to stand at ambient temperature for overnight. Reaction mixture is diluted with  $\text{Et}_2\text{O}$  and 1N HCl. Organic layer is then washed with 1N HCl (2 x 10 mL), brine (2 x 10 mL), dried over  $\text{Na}_2\text{SO}_4$ , filtered and concentrated under vacuum. The crude material is in 5 mL of EtOH and 0.3 mL of 5N NaOH. The mixture is heated to reflux for 1 h. The organic solvent is then removed under vacuum and residue is dissolved in  $\text{CH}_2\text{Cl}_2$  and 1N HCl. Aqueous layer is washed with  $\text{CH}_2\text{Cl}_2$  (2 x 10 mL). Combined organic layer is dried over  $\text{Na}_2\text{SO}_4$ , filtered and concentrated under vacuum. The crude

25

material is purified by MS/LC to give 35 mg (49%) of the title product.  $^1\text{H}$  NMR (400 MHz,  $\text{CDCl}_3$ ):  $\delta$  7.56 (d, 2H,  $J$  = 8.0 Hz), 7.35 (d, 2H,  $J$  = 8.0 Hz), 7.28-7.18 (m, 4H), 7.05 (t, 1H,  $J$  = 7.2 Hz), 6.91 (d, 2H,  $J$  = 8.0 Hz), 6.74 (d, 2H,  $J$  = 8.0 Hz), 5.49 (bs, 1H), 4.45, 4.39 (ABq, 2H,  $J$  = 15.4 Hz), 3.98 (t, 2H,  $J$  = 5.8 Hz), 3.77-3.65 (m, 1H), 3.42 (t, 1H,  $J$  = 8.5 Hz), 3.28, 3.14 (ABq, 2H,  $J$  = 14.0 Hz), 3.05 (t, 1H,  $J$  = 8.5 Hz), 2.86 (s, 3H), 2.28-2.20 (m, 1H), 1.95-1.85 (m, 1H), 1.43 (s, 3H). HRMS ( $\text{ES}^+$ )  $m/z$  exact mass calcd for  $\text{C}_{30}\text{H}_{32}\text{N}_2\text{O}_5\text{F}_3$  ( $m+1$ ) 557.2263, found 557.2257.

#### Example 72


3-(4-{2-[1-(4-tert-Butyl-benzyl)-3-methyl-2-oxo-imidazolidin-4-yl]-ethoxy}-phenyl)-2-(4-fluoro-phenoxy)-2-methyl-propionic acid



To a mixture of 80 mg of 2-(4-Fluoro-phenoxy)-2-methyl-3-{4-[2-(3-methyl-2-oxo-imidazolidin-4-yl)-ethoxy]-phenyl}-propionic acid ethyl ester in 5 mL of dry DMF at 0 °C under an atmosphere of nitrogen, 18 mg of NaH (0.45 mmol) is added. The resulting solution is allowed to stand at r.t. for 20 min. Then 0.08 mL of 4-tert-butylbenzyl bromide is added and resulting mixture is allowed to stand at r.t. for

overnight. Reaction mixture is diluted with Et<sub>2</sub>O and 1N HCl. Organic layer is then washed with 1N HCl (2 x 10 mL), brine (2 x 10 mL), dried over Na<sub>2</sub>SO<sub>4</sub>, filtered and concentrated under vacuum. The crude material is in 5 mL of EtOH and 0.4 mL of 5N NaOH. The mixture is heated to reflux for 1 h. The organic solvent is then removed under vacuum and residue is dissolved in CH<sub>2</sub>Cl<sub>2</sub> and 1N HCl. Aqueous layer is washed with CH<sub>2</sub>Cl<sub>2</sub> (2 x 10 mL). Combined organic layer is dried over Na<sub>2</sub>SO<sub>4</sub>, filtered and concentrated under vacuum. The crude material is purified by MS/LC to give 45 mg (45%) of the title product. <sup>1</sup>H NMR (400 MHz, CDCl<sub>3</sub>): δ 7.33 (d, 2H, J = 8.4 Hz), 7.18, 7.16 (ABq, 4H, J = 8.6 Hz), 6.96-6.92 (m, 2H), 6.88-6.85 (m, 2H), 6.77 (d, 2H, J = 8.8 Hz), 4.38, 4.29 (ABq, 2H, J = 14.8 Hz), 3.98 (t, 2H, J = 6.0 Hz), 3.69 (bs, 1H), 3.73-3.66 (m, 1H), 3.44 (t, 1H, J = 8.6 Hz), 3.25, 3.11 (ABq, 2H, J = 13.8 Hz), 3.06 (t, 1H, J = 8.6 Hz), 2.86 (s, 3H), 2.28-2.19 (m, 1H), 1.97-1.89 (m, 1H), 1.38 (s, 3H), 1.30 (s, 9H). HRMS (ES<sup>+</sup>) m/z exact mass calcd for C<sub>33</sub>H<sub>40</sub>N<sub>2</sub>O<sub>5</sub>F (m+1) 563.2921, found 563.2917.

20

**Example 73**

**2-(4-Fluoro-phenoxy)-2-methyl-3-{4-[2-(3-methyl-1-naphthalen-2-ylmethyl-2-oxo-imidazolidin-4-yl)-ethoxy]-phenyl}-propionic acid**

25

To a mixture of 80 mg of 2-(4-Fluoro-phenoxy)-2-methyl-3-{4-[2-(3-methyl-2-oxo-imidazolidin-4-yl)-ethoxy]-phenyl}-propionic acid ethyl ester in 5 mL of dry DMF at 0 °C under an atmosphere of nitrogen, 18 mg of NaH (0.45 mmol) is added. The resulting solution is allowed to stand at r.t. for 20 min. Then 99 mg of 2-bromomethyl naphthalene is added and resulting mixture is allowed to stand at r.t. for overnight. Reaction mixture is diluted with Et<sub>2</sub>O and 1N HCl. Organic layer is then washed with 1N HCl (2 x 10 mL), brine (2 x 10 mL), dried over Na<sub>2</sub>SO<sub>4</sub>, filtered and concentrated under vacuum. The crude material is in 5 mL of EtOH and 0.4 mL of 5N NaOH. The mixture is heated to reflux for 1 h. The organic solvent is then removed under vacuum and residue is dissolved in CH<sub>2</sub>Cl<sub>2</sub> and 1N HCl. Aqueous layer is washed with CH<sub>2</sub>Cl<sub>2</sub> (2 x 10 mL). Combined organic layer is dried over Na<sub>2</sub>SO<sub>4</sub>, filtered and concentrated under vacuum. The crude material is purified by MS/LC to give 51 mg (51%) of the title product. <sup>1</sup>H NMR (400 MHz, CDCl<sub>3</sub>): δ 7.82-7.79 (m, 3H), 7.67 (s, 1H), 7.50-7.44 (m, 2H), 7.35 (d, 1H, J = 8.4 Hz), 7.14 (d, 2H, J = 8.6 Hz), 6.93 (t, 2H, J = 8.4 Hz), 6.87-6.83 (m, 2H), 6.71 (d, 2H, J = 8.6 Hz), 5.11 (bs, 1H), 4.53 (s, 2H), 3.95-3.90 (m, 2H), 3.85-3.70 (m, 1H), 3.46 (t, 1H, J = 8.7 Hz), 3.23, 3.09 (ABq, 2H, J = 14.0 Hz), 3.07 (t, 1H, J = 8.7 Hz), 2.89 (s, 3H), 2.26-2.18 (m, 1H), 1.96-1.87 (m, 1H), 1.36 (s, 3H). HRMS (ES<sup>+</sup>) m/z exact mass calcd for C<sub>33</sub>H<sub>34</sub>N<sub>2</sub>O<sub>5</sub>F (m+1) 557.2452, found 557.2439.

**3-(4-{2-[1-(3,4-Dimethyl-benzyl)-3-methyl-2-oxo-imidazolidin-4-yl]-ethoxy}-phenyl)-2-(4-fluoro-phenoxy)-2-methyl-propionic acid**

5           To a mixture of 80 mg of 2-(4-Fluoro-phenoxy)-2-methyl-3-{4-[2-(3-methyl-2-oxo-imidazolidin-4-yl)-ethoxy]-phenyl}-propionic acid ethyl ester in 5 mL of dry DMF at 0 °C under an atmosphere of nitrogen, 18 mg of NaH (0.45 mmol) is added. The resulting solution is allowed to stand at r.t.

10   for 20 min. Then 0.07 mL of 3,4-dimethylbenzyl chloride is added and resulting mixture is allowed to stand at r.t. for overnight. Reaction mixture is diluted with Et<sub>2</sub>O and 1N HCl. Organic layer is then washed with 1N HCl (2 x 10 mL), brine (2 x 10 mL), dried over Na<sub>2</sub>SO<sub>4</sub>, filtered and

15   concentrated under vacuum. The crude material is in 5 mL of EtOH and 0.4 mL of 5N NaOH. The mixture is heated to reflux for 1 h. The organic solvent is then removed under vacuum and residue is dissolved in CH<sub>2</sub>Cl<sub>2</sub> and 1N HCl. Aqueous layer is washed with CH<sub>2</sub>Cl<sub>2</sub> (2 x 10 mL). Combined organic layer

20   is dried over Na<sub>2</sub>SO<sub>4</sub>, filtered and concentrated under vacuum. The crude material is purified by MS/LC to give 54 mg (56%) of the title product. <sup>1</sup>H NMR (400 MHz, CDCl<sub>3</sub>): δ

25   7.18 (d, 2H, *J* = 8.4 Hz), 7.06 (d, 1H, *J* = 7.6 Hz), 7.00 (s, 1H), 6.97-6.85 (m, 5H), 6.75 (d, 2H, *J* = 8.4 Hz), 4.34, 4.25 (ABq, 2H, *J* = 15.0 Hz), 4.31 (bs, 1H), 3.98-3.90 (m, 2H), 3.69-3.61 (m, 1H), 3.39 (t, 1H, *J* = 8.7 Hz), 3.25, 3.11 (ABq, 2H, *J* = 14.0 Hz), 3.00 (t, 1H, *J* = 8.7 Hz), 2.84 (s, 3H), 2.29-2.17 (m, 1H), 2.22 (s, 6H), 1.94-1.87 (m, 1H), 1.37 (s, 3H). HRMS (ES<sup>+</sup>) *m/z* exact mass calcd for C<sub>31</sub>H<sub>36</sub>N<sub>2</sub>O<sub>5</sub>F

30   (*m*+1) 535.2608, found 535.2614.



## Example 75

3-(4-{2-[1-(4-tert-Butyl-benzyl)-3-methyl-2-oxo-imidazolidin-4-yl]-ethoxy}-phenyl)-2-methyl-2-(4-trifluoromethyl-phenoxy)-propionic acid

5

To a mixture of 80 mg of 2-Methyl-3-{4-[2-(3-methyl-2-oxo-imidazolidin-4-yl)-ethoxy]-phenyl}-2-(4-trifluoromethyl-phenoxy)-propionic acid ethyl ester in 5 mL of dry DMF at 0 °C under an atmosphere of nitrogen, 16 mg of NaH (0.40 mmol) is added. The resulting solution is allowed to stand at r.t. for 20 minute. Then 0.07 mL of 4-tert-butylbenzyl bromide is added and resulting mixture is allowed to stand at r.t. for overnight. Reaction mixture is diluted with Et<sub>2</sub>O and 1N HCl. Organic layer is then washed with 1N HCl (2 x 10 mL), brine (2 x 10 mL), dried over Na<sub>2</sub>SO<sub>4</sub>, filtered and concentrated under vacuum. The crude material is in 5 mL of EtOH and 0.5 mL of 5N NaOH. The mixture is heated to reflux for 1 h. The organic solvent is then removed under vacuum and residue is dissolved in CH<sub>2</sub>Cl<sub>2</sub> and 1N HCl. Aqueous layer is washed with CH<sub>2</sub>Cl<sub>2</sub> (2 x 10 mL). Combined organic layer is dried over Na<sub>2</sub>SO<sub>4</sub>, filtered and concentrated under vacuum. The crude material is purified by MS/LC to give 40 mg (40%) of the title product. <sup>1</sup>H NMR (400 MHz, CDCl<sub>3</sub>): δ 7.51 (d, 2H, J = 8.8 Hz), 7.38-7.32 (m, 2H), 7.16 (t, 4H, J = 8.8 Hz), 6.95 (d, 2H, J = 8.8 Hz), 6.76 (d, 2H, J = 8.8 Hz), 4.37, 4.28 (ABq, 2H, J = 14.8 Hz), 4.24 (bs, 1H), 3.97 (t, 2H, J = 6.0 Hz), 3.74-3.67 (m, 1H), 3.44 (t, 1H, J = 9.2 Hz), 3.31, 3.16 (ABq, 2H, J = 13.8 Hz), 3.07 (t, 1H, J = 9.2 Hz), 2.85 (s, 3H), 2.27-2.18 (m, 1H), 1.98-1.89 (m, 1H), 1.50 (s, 3H), 1.29 (s, 9H). HRMS (ES<sup>+</sup>) m/z exact mass calcd for C<sub>34</sub>H<sub>40</sub>N<sub>2</sub>O<sub>5</sub>F<sub>3</sub> (m+1) 613.2889, found 613.2872.

**Example 76****2-Methyl-3-{4-[2-(3-methyl-1-naphthalen-2-ylmethyl-2-oxo-imidazolidin-4-yl)-ethoxy]-phenyl}-2-(4-trifluoromethyl-phenoxy)-propionic acid**

5

To a mixture of 80 mg of 2-Methyl-3-{4-[2-(3-methyl-2-oxo-imidazolidin-4-yl)-ethoxy]-phenyl}-2-(4-trifluoromethyl-phenoxy)-propionic acid ethyl ester in 5 mL of dry DMF at 0 °C under an atmosphere of nitrogen, 16 mg of NaH (0.40 mmol) is added. The resulting solution is allowed to stand at r.t. for 20 minute. Then 89 mg of 2-bromomethyl naphthalene is added and resulting mixture is allowed to stand at r.t. for overnight. Reaction mixture is diluted with Et<sub>2</sub>O and 1N HCl. Organic layer is then washed with 1N HCl (2 x 10 mL), brine (2 x 10 mL), dried over Na<sub>2</sub>SO<sub>4</sub>, filtered and concentrated under vacuum. The crude material is in 5 mL of EtOH and 0.5 mL of 5N NaOH. The mixture is heated to reflux for 1 h. The organic solvent is then removed under vacuum and residue is dissolved in CH<sub>2</sub>Cl<sub>2</sub> and 1N HCl. Aqueous layer is washed with CH<sub>2</sub>Cl<sub>2</sub> (2 x 10 mL). Combined organic layer is dried over Na<sub>2</sub>SO<sub>4</sub>, filtered and concentrated under vacuum. The crude material is purified by MS/LC to give 51 mg (51%) of the title product. <sup>1</sup>H NMR (400 MHz, CDCl<sub>3</sub>): δ 7.81-7.77 (m, 3H), 7.66 (s, 1H), 7.48-7.45 (m, 4H), 7.35 (dd, 1H, J = 8.8, 1.6 Hz), 7.12 (d, 2H, J = 8.8 Hz), 6.92 (d, 2H, J = 8.4 Hz), 6.69 (d, 2H, J = 8.4 Hz), 6.20 (bs, 1H), 4.51 (s, 2H), 3.92-3.90 (m, 2H), 3.72-3.64 (m, 1H), 3.41 (t, 1H, J = 8.8 Hz), 3.29, 3.13 (ABq, 2H, J = 13.8 Hz), 3.03 (t, 1H, J = 8.8 Hz), 2.86 (s, 3H), 2.22-2.15 (m, 1H), 1.93-1.84 (m, 1H), 1.46 (s, 3H). HRMS (ES<sup>+</sup>) m/z exact mass calcd for C<sub>34</sub>H<sub>34</sub>N<sub>2</sub>O<sub>5</sub>F<sub>3</sub> (m+1) 6072420, found 6072417.

30

## Example 77

3-(4-{2-[1-(3,4-Dimethyl-benzyl)-3-methyl-2-oxo-imidazolidin-4-yl]-ethoxy}-phenyl)-2-methyl-2-(4-trifluoromethyl-phenoxy)-propionic acid

5

To a mixture of 80 mg of 2-Methyl-3-{4-[2-(3-methyl-2-oxo-imidazolidin-4-yl)-ethoxy]-phenyl}-2-(4-trifluoromethyl-phenoxy)-propionic acid ethyl ester in 5 mL of dry DMF at 0 °C under an atmosphere of nitrogen, 16 mg of NaH (0.40 mmol) is added. The resulting solution is allowed to stand at r.t. for 20 min. Then 0.06 mL of 3,4-dimethylbenzyl chloride is added and resulting mixture is allowed to stand at r.t. for overnight. Reaction mixture is diluted with Et<sub>2</sub>O and 1N HCl. Organic layer is then washed with 1N HCl (2 x 10 mL), brine (2 x 10 mL), dried over Na<sub>2</sub>SO<sub>4</sub>, filtered and concentrated under vacuum. The crude material is in 5 mL of EtOH and 0.5 mL of 5N NaOH. The mixture is heated to reflux for 1 h. The organic solvent is then removed under vacuum and residue is dissolved in CH<sub>2</sub>Cl<sub>2</sub> and 1N HCl. Aqueous layer is washed with CH<sub>2</sub>Cl<sub>2</sub> (2 x 10 mL). Combined organic layer is dried over Na<sub>2</sub>SO<sub>4</sub>, filtered and concentrated under vacuum. The crude material is purified by MS/LC to give 54 mg (56%) of the title product. <sup>1</sup>H NMR (400 MHz, CDCl<sub>3</sub>): δ 7.51 (d, 2H, J = 8.6 Hz), 7.16 (d, 1H, J = 8.6 Hz), 7.10 (d, 1H, J = 8.0 Hz), 7.06 (d, 1H, J = 8.0 Hz), 6.99-6.94 (m, 3H), 6.75 (d, 2H, J = 8.4 Hz), 4.33, 4.25 (ABq, 2H, J = 14.8 Hz), 4.07 (bs, 1H), 3.97-3.93 (m, 2H), 3.70-3.65 (m, 1H), 3.41 (t, 1H, J = 8.8 Hz), 3.32, 3.16 (ABq, 2H, J = 13.8 Hz), 3.03 (t, 1H, J = 8.8 Hz), 2.85 (s, 3H), 2.28-2.20 (m, 1H), 2.22 (s, 3H), 1.96-1.88 (m, 1H),

1.50 (s, 3H). HRMS (ES<sup>+</sup>) *m/z* exact mass calcd for  
C<sub>32</sub>H<sub>36</sub>N<sub>2</sub>O<sub>5</sub>F<sub>3</sub> (*m*+1) 585.2576, found 585.2578.

#### Example 79

5 **2-Methyl-3-(4-{2-[3-methyl-2-oxo-1-(3-phenoxy-benzyl)-  
imidazolidin-4-yl]-ethoxy}-phenyl)-2-phenoxy-propionic acid**

The titled compound is prepared, according to the procedures  
substantially as described herein, yield (0.192g, 52%).

10 Mass [EI+] 581 (M+H)<sup>+</sup>, [EI-] 579 (M-H)<sup>-</sup>.

#### Example 80

**2-Methyl-3-(4-{2-[3-methyl-2-oxo-1-(3-trifluoromethoxy-  
benzyl)-imidazolidin-4-yl]-ethoxy}-phenyl)-2-phenoxy-  
15 propionic acid**

The titled compound is prepared, according to the procedures  
substantially as described herein, yield (0.127g, 35%).

Mass [EI+] 573 (M+H)<sup>+</sup>, [EI-] 571 (M-H)<sup>-</sup>.

20

#### Example 81

**2-Methyl-3-(4-{2-[3-methyl-2-oxo-1-(3-trifluoromethoxy-  
benzyl)-imidazolidin-4-yl]-ethoxy}-phenyl)-2-phenoxy-  
25 propionic acid**

The titled compound is prepared, according to the procedures  
substantially as described herein, yield (0.238g, 63%).


Mass [EI+] 573 (M+H)<sup>+</sup>, [EI-] 571 (M-H)<sup>-</sup>.

30

**Example 82**

**2-(2-Fluoro-phenoxy)-2-methyl-3-(4-{2-[3-methyl-2-oxo-1-(4-trifluoromethoxy-benzyl)-imidazolidin-4-yl]-ethoxy}-phenyl)-propionic acid**

5



The titled compound is prepared, substantially according to the procedures described herein to produce a white foamy solid (0.087g, 64%). Mass [EI+] 591 (M+H)<sup>+</sup>, [EI-] 589 (M-H)<sup>-</sup>.

10

**Example 83**

**2-(2-Fluoro-phenoxy)-2-methyl-3-(4-{2-[3-methyl-1-(4-methylbenzyl)-2-oxo-imidazolidin-4-yl]-ethoxy}-phenyl)-propionic acid**

15 The titled compound is prepared, substantially according to the procedures described herein to produce a white foamy solid (0.022g, 18%). Mass [EI+] 521 (M+H)<sup>+</sup>, [EI-] 519 (M-H)<sup>-</sup>.

**Example 84**

20 **3-(4-{2-[1-(3,4-Dichloro-benzyl)-3-methyl-2-oxo-imidazolidin-4-yl]-ethoxy}-phenyl)-2-(2-fluoro-phenoxy)-2-methyl-propionic acid**

The titled compound is prepared, substantially according to the procedures described herein to produce a white foamy solid (0.107g, 80%). Mass [EI+] 574, 577 (M+H)<sup>+</sup>, [EI-] 573, 575 (M-H)<sup>-</sup>.

25

**Example 85**

**3-(4-{2-[1-(3,4-Difluoro-benzyl)-3-methyl-2-oxo-imidazolidin-4-yl]-ethoxy}-phenyl)-2-(2-fluoro-phenoxy)-2-methyl-propionic acid**

5

The titled compound is prepared, substantially according to the procedures described herein to produce a white foamy solid (0.107g, 84%). Mass [EI+] 543 (M+H)<sup>+</sup>, [EI-] 541 (M-H)<sup>-</sup>.

10

**Example 86**

**3-(4-{2-[1-(3,5-Difluoro-benzyl)-3-methyl-2-oxo-imidazolidin-4-yl]-ethoxy}-phenyl)-2-(2-fluoro-phenoxy)-2-methyl-propionic acid**

15 The titled compound is prepared, substantially according to the procedures described herein to produce a white foamy solid (0.0824g, 65%). Mass [EI+] 543 (M+H)<sup>+</sup>, [EI-] 541 (M-H)<sup>-</sup>.

20

**Example 87**

**2-(2-Fluoro-phenoxy)-2-methyl-3-{4-[2-(3-methyl-1-naphthalen-2-ylmethyl-2-oxo-imidazolidin-4-yl)-ethoxy]-phenyl}-propionic acid**

25

The titled compound is prepared, substantially according to the procedures described herein to produce a white foamy solid (0.0753g, 58%). Mass [EI+] 543 (M+H)<sup>+</sup>, [EI-] 541 (M-H)<sup>-</sup>.

**Example 88**

30

**3-{4-[2-(1-Biphenyl-4-ylmethyl-3-methyl-2-oxo-imidazolidin-4-yl)-ethoxy]-phenyl}-2-methyl-2-phenoxy-propionic acid**

The title compound is prepared substantially according to the procedures described herein, and concentrated to an oil

which is purified by MS/LC to yield an oil (25 mg, 25%).  
MS [EI+] 565 (M+H)<sup>+</sup>. Structure confirmed by 1H-NMR.

#### Example 89

5        **2-Methyl-3-(4-{2-[3-methyl-2-oxo-1-(4-trifluoromethoxy-  
benzyl)-imidazolidin-4-yl]-ethoxy}-phenyl)-2-phenoxy-  
propionic acid**

NaH (8.8 mg, 0.22 mmol, 60% w/w in mineral oil) is added to  
a cooled (0° C) solution of 2-Methyl-3-{4-[2-(3-methyl-2-  
10    oxo-imidazolidin-4-yl)-ethoxy]-phenyl}-2-phenoxy-propionic  
acid ethyl ester (78.5 mg, 0.18 mmol) in DMF (1 ml). After 5  
minutes, the reaction mixture is removed from the cooling  
bath and stirred while warming to ambient temperature over  
one hour. After the solution is cooled again to 0° C, 4-  
15    (trifluoromethoxy)benzyl bromide (59 ul, 0.37 mmol) is added  
in a single portion, the cooling bath removed, and the  
mixture stirred for one hour. The reaction mixture is  
quenched with 1N HCl and extracted with ethyl acetate (2x).  
The organic extracts were washed with water, brine, dried  
20    (Na<sub>2</sub>SO<sub>4</sub>), and concentrated to an oil. The crude product is  
dissolved in methanol (5 ml), treated with 5N NaOH (0.5 ml),  
and stirred for 14 hours at ambient temperature. After  
concentration to remove methanol, the residue is partitioned  
between diethyl ether and water. The aqueous layer is  
25    adjusted to pH 2 with conc. HCl, then extracted with ethyl  
acetate (2 x 15 ml). The combined ethyl acetate extracts  
were washed with water, brine, dried (Na<sub>2</sub>SO<sub>4</sub>), and  
concentrated to an oil which is purified by MS/LC to yield  
an oil (33 mg, 31%).  
30    MS [EI+] 573 (M+H)<sup>+</sup>. Structure confirmed by 1H-NMR.

#### Example 90

**3-(4-{2-[1-(4-Fluoro-3-trifluoromethyl-benzyl)-3-methyl-2-**

**oxo-imidazolidin-4-yl]-ethoxy}-phenyl)-2-methyl-2-phenoxy-propionic acid**

NaH (9.2 mg, 0.23 mmol, 60% w/w in mineral oil) is added to a cooled (0° C) solution of 2-Methyl-3-{4-[2-(3-methyl-2-oxo-imidazolidin-4-yl)-ethoxy]-phenyl}-2-phenoxy-propionic acid ethyl ester (82.0 mg, 0.19 mmol) in DMF (1 ml). After 5 minutes, the reaction mixture is removed from the cooling bath and stirred while warming to ambient temperature over one hour. After the solution is cooled again to 0° C, 4-fluoro-3-(trifluoromethyl)benzyl bromide (148 mg, 0.57 mmol) is added in a single portion, the cooling bath removed, and the mixture stirred for one hour. The reaction mixture is quenched with 1N HCl and extracted with ethyl acetate (2x). The organic extracts were washed with water, brine, dried (Na<sub>2</sub>SO<sub>4</sub>), and concentrated to an oil. The crude product is dissolved in methanol (5 ml), treated with 5N NaOH (0.5 ml), and stirred for 18 hours at ambient temperature. After concentration to remove methanol, the residue is partitioned between diethyl ether and water. The aqueous layer is adjusted to pH 2 with conc. HCl, then extracted with ethyl acetate (2 x 15 ml). The combined ethyl acetate extracts were washed with water, brine, dried (Na<sub>2</sub>SO<sub>4</sub>), and concentrated to an oil which is purified by MS/LC to yield an oil (41 mg, 37%). MS [EI+] 575 (M+H)<sup>+</sup>. Structure confirmed by 1H-NMR.

**Example 91**

**3-(4-{2-[1-(3-Fluoro-4-trifluoromethyl-benzyl)-3-methyl-2-oxo-imidazolidin-4-yl]-ethoxy}-phenyl)-2-methyl-2-phenoxy-propionic acid**

The title compound is prepared substantially as described herein and concentrated to an oil which is purified by



MS/LC to yield an oil (45 mg, 41%).


MS [ EI+ ] 575 (M+H)<sup>+</sup>. Structure confirmed by 1H-NMR.

#### Example 92

5      **2-Methyl-3-(4-{2-[3-methyl-2-oxo-1-(3-trifluoromethyl-  
benzyl)-imidazolidin-4-yl]-ethoxy}-phenyl)-2-phenoxy-  
propionic acid**

##### Step A

10      **2-Methyl-3-(4-{2-[3-methyl-2-oxo-1-(3-trifluoromethyl-  
benzyl)-imidazolidin-4-yl]-ethoxy}-phenyl)-2-phenoxy-  
propionic acid ethyl ester**



NaH (34 mg, 0.84 mmol, 60% w/w in mineral oil) is added to a  
15      cooled (0° C) solution of 2-Methyl-3-{4-[2-(3-methyl-2-oxo-  
imidazolidin-4-yl)-ethoxy]-phenyl}-2-phenoxy-propionic acid  
ethyl ester (300 mg, 0.70 mmol) in DMF (1.5 ml). After 5  
minutes, the reaction mixture is removed from the cooling  
bath and stirred while warming to ambient temperature over  
20      1.5 hours. After the solution is cooled again to 0° C, 3-  
(trifluoromethyl)benzyl bromide ( 217 mg, 0.91 mmol) is  
added in a single portion, the cooling bath removed, and the  
mixture stirred for 2.5 hours. The reaction mixture is  
quenched with 1N HCl and extracted with ethyl acetate (2x).  
25      The organic extracts were washed with water, brine, dried  
(Na<sub>2</sub>SO<sub>4</sub>), and concentrated to an oil. The crude product is  
purified by flash chromatography on a Biotage silica  
cartridge (gradient elution, 5:1 hexanes:ethyl acetate to  
1:1 hexanes:ethyl acetate) to provide an oil (273 mg) which


is a mixture of the title compound and corresponding substituted-benzyl ester as two closely eluting compounds. MS [ EI+] 585 (M+H)<sup>+</sup>. Structure confirmed by 1H-NMR.

5

Step B

2-Methyl-3-(4-{2-[3-methyl-2-oxo-1-(3-trifluoromethyl-benzyl)-imidazolidin-4-yl]-ethoxy}-phenyl)-2-phenoxy-propionic acid

10




To a solution of the esters obtained from Step A (270 mg) in absolute ethanol (4.5 ml) is added 5N NaOH (1.9 ml, 9.5 mmol), and the mixture stirred for 20 hours at ambient temperature. The mixture is poured into 1N HCl, then extracted with ethyl acetate (2 x 20 ml). The combined ethyl acetate extracts were washed with water, brine, dried (Na<sub>2</sub>SO<sub>4</sub>), and concentrated to an oil which is purified is purified by flash chromatography on a Biotage silica cartridge (gradient elution, 3:1 hexanes:ethyl acetate to 1:2 hexanes:ethyl acetate to yield a foam (78 mg). MS [ EI+] 557 (M+H)<sup>+</sup>. Structure confirmed by 1H-NMR.

25

**Example 93**

3-(4-{2-[1-(3,4-Dichloro-benzyl)-3-methyl-2-oxo-

**imidazolidin-4-yl]-ethoxy}-phenyl)-2-methyl-2-phenoxy-  
propionic acid**



5

Step A

3-(4-{2-[1-(3,4-Dichloro-benzyl)-3-methyl-2-oxo-  
imidazolidin-4-yl]-ethoxy}-phenyl)-2-methyl-2-phenoxy-  
propionic acid ethyl ester

10

NaH (34 mg, 0.84 mmol, 60% w/w in mineral oil) is added to a cooled (0° C) solution of 2-Methyl-3-{4-[2-(3-methyl-2-oxo-imidazolidin-4-yl)-ethoxy]-phenyl}-2-phenoxy-propionic acid  
ethyl ester (300 mg, 0.70 mmol) in DMF (1.5 ml). After 5  
minutes, the reaction mixture is removed from the cooling  
bath and stirred while warming to ambient temperature over  
1.5 hours. After the solution is cooled again to 0° C, 3,4-  
dichlorobenzyl bromide ( 218 mg, 0.91 mmol) is added in a  
single portion, the cooling bath removed, and the mixture  
stirred for 2.5 hours. The reaction mixture is quenched with  
1N HCl and extracted with ethyl acetate (2x). The organic  
extracts were washed with water, brine, dried (Na<sub>2</sub>SO<sub>4</sub>), and  
concentrated to an oil. The crude product is purified by  
flash chromatography on a Biotage silica cartridge (gradient  
elution, 5:1 hexanes:ethyl acetate to 1:1 hexanes:ethyl  
acetate) to provide an oil (320 mg) which is a mixture of  
the title compound and corresponding substituted-benzyl

25

ester as two closely eluting compounds.

MS [ EI+] 586 (M+H)<sup>+</sup>. Structure confirmed by 1H-NMR.

#### Step B

5           3-(4-{2-[1-(3,4-Dichloro-benzyl)-3-methyl-2-oxo-  
imidazolidin-4-yl]-ethoxy}-phenyl)-2-methyl-2-phenoxy-  
propionic acid

To a solution of the esters obtained from Step A (320 mg) in  
10 absolute ethanol (5.5 ml) is added 5N NaOH (2.2 ml, 10.9  
mmol), and the mixture stirred for 20 hours at ambient  
temperature. The mixture is poured into 1N HCl, then  
extracted with ethyl acetate (2 x 20 ml). The combined ethyl  
acetate extracts were washed with water, brine, dried  
15 (Na<sub>2</sub>SO<sub>4</sub>), and concentrated to an oil which is purified is  
purified by flash chromatography on a Biotage silica  
cartridge (gradient elution, 3:1 hexanes:ethyl acetate to  
1:2 hexanes:ethyl acetate to yield a foam (180 mg.)  
MS [ EI+] 558 (M+H)<sup>+</sup>. Structure confirmed by 1H-NMR.

20

#### **Example 94**

3-(4-{2-[1-(3,5-Bis-trifluoromethyl-benzyl)-3-methyl-2-oxo-  
imidazolidin-4-yl]-ethoxy}-phenyl)-2-methyl-2-phenoxy-  
propionic acid

25 The title compound was prepared using substantially the  
procedures described herein.

#### **Example 95**

30 3-(4-{2-[1-(3,5-Bis-trifluoromethyl-benzyl)-3-methyl-2-oxo-  
imidazolidin-4-yl]-ethoxy}-phenyl)-2-methyl-2-phenoxy-  
propionic acid

Step A

3-(4-{2-[1-(3,5-Bis-trifluoromethyl-benzyl)-3-methyl-2-oxo-imidazolidin-4-yl]-ethoxy}-phenyl)-2-methyl-2-phenoxy-propionic acid ethyl ester

- 5 To a cooled (0° C) suspension of NaH (41 mg, 1.03 mmol, 60% w/w in mineral oil) in DMF (1.0 ml) is added a solution of 2-Methyl-3-{4-[2-(3-methyl-2-oxo-imidazolidin-4-yl)-ethoxy]-phenyl}-2-phenoxy-propionic acid ethyl ester (294 mg, 0.69 mmol) in DMF (1.5 ml). After 5 minutes, the reaction
- 10 mixture is removed from the cooling bath and stirred while warming to ambient temperature over 45 minutes. After the solution is cooled again to 0° C, 3,5-Bis(trifluoromethyl)benzyl bromide (423 mg, 1.38 mmol) is added in a single portion, the cooling bath removed, and the
- 15 mixture stirred for 1 hour. The reaction mixture is quenched with 1N HCl and extracted with ethyl acetate (2x). The organic extracts were washed with water, brine, dried (Na<sub>2</sub>SO<sub>4</sub>), and concentrated to an oil. The crude product is purified by flash chromatography on a Biotage silica
- 20 cartridge (gradient elution, 5:1 hexanes:ethyl acetate to 1:1 hexanes:ethyl acetate) to provide an oil (380mg) which is a mixture of the title compound and corresponding substituted-benzyl ester as two closely eluting compounds. MS [ EI+ ] 653 (M+H)<sup>+</sup>. Structure confirmed by <sup>1</sup>H-NMR.

25

Step B

3-(4-{2-[1-(3,5-Bis-trifluoromethyl-benzyl)-3-methyl-2-oxo-imidazolidin-4-yl]-ethoxy}-phenyl)-2-methyl-2-phenoxy-propionic acid

30

To a solution of the esters obtained from **Step A** (378 mg) in absolute ethanol (4 ml) is added 5N NaOH (1.5 ml, 7.5 mmol),

and the mixture stirred for 20 hours at ambient temperature. The mixture is poured into 1N HCl, then extracted with ethyl acetate (2 x 20 ml). The combined ethyl acetate extracts were washed with water, brine, dried (Na<sub>2</sub>SO<sub>4</sub>), and concentrated to an oil which is purified is purified by flash chromatography on a Biotage silica cartridge (gradient elution, 3:1 hexanes:ethyl acetate to 1:2 hexanes:ethyl acetate to yield a foam (266 mg.) MS [ EI+ ] 625 (M+H)<sup>+</sup>. Structure confirmed by <sup>1</sup>H-NMR.

10

**Example 96**

**3-(4-{2-[1-(4-Benzoyl-benzyl)-3-methyl-2-oxo-imidazolidin-4-yl]-ethoxy}-phenyl)-2-methyl-2-phenoxy-propionic acid**

15

Step A

3-(4-{2-[1-(4-Benzoyl-benzyl)-3-methyl-2-oxo-imidazolidin-4-yl]-ethoxy}-phenyl)-2-methyl-2-phenoxy-propionic acid ethyl ester

20

To a cooled (0° C) suspension of NaH (45 mg, 1.1 mmol, 60% w/w in mineral oil) in DMF (0.5 ml) is added a solution of 2-Methyl-3-{4-[2-(3-methyl-2-oxo-imidazolidin-4-yl)-ethoxy]-phenyl}-2-phenoxy-propionic acid ethyl ester (320 mg, 0.75 mmol) in DMF (1.5 ml). After 5 minutes, the reaction mixture is removed from the cooling bath and stirred while warming to ambient temperature over 30 minutes. The solution is cooled again to 0° C, and 4-(bromomethyl)benzophenone (188 mg, 0.68 mmol) is added in a single portion, the cooling bath removed, and the mixture stirred for one hour. The reaction mixture is quenched with 1N HCl and extracted with ethyl acetate (2x). The organic extracts were washed with water, brine, dried (Na<sub>2</sub>SO<sub>4</sub>), and concentrated to an oil. The crude product is purified by flash chromatography

30

on a Biotage silica cartridge (gradient elution, 2:1 hexanes:ethyl acetate to 1:2 hexanes:ethyl acetate) to provide an oil (46 mg, 10%).

MS [EI+] 621 (M+H)<sup>+</sup>. Structure confirmed by <sup>1</sup>H-NMR.

5

#### Step B

3-(4-{2-[1-(4-Benzoyl-benzyl)-3-methyl-2-oxo-imidazolidin-4-yl]-ethoxy}-phenyl)-2-methyl-2-phenoxy-propionic acid

- 10 To a solution of the ester obtained from **Step A** (46 mg, 0.1 mmol) in absolute ethanol (2 ml) is added 5N NaOH (0.5 ml, 2.5 mmol), and the mixture stirred for 14 hours at ambient temperature. The mixture is poured into 1N HCl, then extracted with ethyl acetate (2 x 20 ml). The combined ethyl
- 15 acetate extracts were washed with water, brine, dried (Na<sub>2</sub>SO<sub>4</sub>), and concentrated to an oil which is purified is purified by MS/LC to provide a oil (26 mg, 7%).
- MS [EI+] 593 (M+H)<sup>+</sup>. Structure confirmed by <sup>1</sup>H-NMR.

20

#### **Example 97**

3-(4-{2-[1-(4-Isopropyl-benzyl)-3-methyl-2-oxo-imidazolidin-4-yl]-ethoxy}-phenyl)-2-methyl-2-phenoxy-propionic acid

- To a cooled (0° C) suspension of NaH (24 mg, 0.60 mmol, 60% w/w in mineral oil) in DMF (1 ml) is added a solution of 2-
- 25 Methyl-3-{4-[2-(3-methyl-2-oxo-imidazolidin-4-yl)-ethoxy]-phenyl}-2-phenoxy-propionic acid ethyl ester (128 mg, 0.30 mmol) in DMF (1.5 ml). After 5 minutes, the reaction mixture is removed from the cooling bath and stirred while warming to ambient temperature over 45 minutes. After the
- 30 solution is cooled again to 0° C, 4-isopropylbenzyl chloride (126 mg, 0.75 mmol) is added in a single portion, the cooling bath removed, and the mixture stirred for 50

minutes. The reaction mixture is quenched with 1N HCl and extracted with ethyl acetate (2x). The organic extracts were washed with water, brine, dried (Na<sub>2</sub>SO<sub>4</sub>), and concentrated to an oil. The crude product is dissolved in ethanol (4 ml),  
5 treated with 5N NaOH (1.5 ml), and stirred for 24 hours at ambient temperature. The mixture is poured into 1N HCl, then extracted with ethyl acetate (2 x 20 ml). The combined ethyl acetate extracts were washed with water, brine, dried (Na<sub>2</sub>SO<sub>4</sub>), and concentrated to an oil which is purified by  
10 MS/LC to provide a foam ( 110 mg, 69%).  
MS [ EI+] 531 (M+H)<sup>+</sup>. Structure confirmed by 1H-NMR.

#### Example 98

**2-Methyl-3-(4-{2-[3-methyl-1-(6-methyl-naphthalen-2-ylmethyl)-2-oxo-imidazolidin-4-yl]-ethoxy}-phenyl)-2-phenoxy-propionic acid**  
15

The title compound was prepared substantially as described herein and concentrated to an oil which is purified by MS/LC to provide a white solid ( 83 mg, 50%).  
20 MS [ EI+] 553 (M+H)<sup>+</sup>. Structure confirmed by 1H-NMR.  
To a solution of the ester obtained from Step A (69 mg, 0.1 mmol ) in methanol (2 ml) is added 5N NaOH (0.2 ml, 1 mmol), and the mixture stirred for 16 hours at ambient temperature. The mixture is poured into 1N HCl (20 ml), then extracted  
25 with ethyl acetate (2 x 15 ml). The combined ethyl acetate extracts were washed with water, brine, dried (Na<sub>2</sub>SO<sub>4</sub>), and concentrated to a foam (62 mg, 100%).  
MS [ EI+] 625 (M+H)<sup>+</sup>. Structure confirmed by 1H-NMR.

30

#### Example 99

**2-Methyl-3-(3-{2-[3-methyl-2-oxo-1-(4-trifluoromethyl-benzyl)-imidazolidin-4-yl]-ethoxy}-phenyl)-2-phenoxy-**



**propionic acid (diastereomer 4)**Step A

2-Methyl-3-(3-{2-[3-methyl-2-oxo-1-(4-trifluoromethyl-  
5 benzyl)-imidazolidin-4-yl]-ethoxy}-phenyl)-2-phenoxy-  
propionic acid ethyl ester (diastereomer 4)

To a solution of 3-(3-Hydroxy-phenyl)-2-methyl-2-phenoxy-  
propionic acid ethyl ester (Isomer 2) (43 mg, 0.143 mmol,)   
10 and Toluene-4-sulfonic acid 2-[3-methyl-2-oxo-1-(4-  
trifluoromethyl-benzyl)-imidazolidin-4-yl]-ethyl ester  
(Isomer 2) (72 mg, 0.157 mmol) in DMF (1 ml) at 23° C is  
added Cs<sub>2</sub>CO<sub>3</sub> ( 61 mg, 0.186 mmol) and the suspension heated  
at 55° C for 7 hours. The mixture is cooled, filtered, and  
15 the filter cake washed with DMF (10 ml). The filtrate is  
poured into 1N HCl (35 ml) and extracted with ethyl acetate  
(2x20 ml). The combined organic extracts were washed with  
water (25 ml), brine (20 ml), dried (Na<sub>2</sub>SO<sub>4</sub>) and  
concentrated to an oil. Purification by flash chromatography  
20 on a Biotage silica cartridge (gradient elution, 4:1  
hexanes:acetone to 1:1 hexanes:acetone) provided an oil,  
(72 mg, 87%) (diastereomer 4).  
MS [ EI+ ] 585 (M+H)<sup>+</sup>. Structure confirmed by 1H-NMR.

25

Step B

2-Methyl-3-(3-{2-[3-methyl-2-oxo-1-(4-trifluoromethyl-  
benzyl)-imidazolidin-4-yl]-ethoxy}-phenyl)-2-phenoxy-  
propionic acid (diastereomer 4)

30

To a solution of the ester obtained from Step A (72 mg, 0.12  
mmol ) in methanol (2 ml) is added 5N NaOH (0.5 ml, 2.5  
mmol), and the mixture stirred for 20 hours at ambient

temperature. The mixture is poured into 1N HCl (20 ml), then extracted with ethyl acetate (2 x 15 ml). The combined ethyl acetate extracts were washed with water, brine, dried ( $\text{Na}_2\text{SO}_4$ ), and concentrated to an oil (64 mg, 94%)


5 (diastereomer 4).

MS [EI+] 557 (M+H)<sup>+</sup>. Structure confirmed by <sup>1</sup>H-NMR.

### Example 100

10 **2-methyl-3-(4-{3-[3-methyl-2-oxo-1-(4-trifluoromethyl-benzyl)-imidazolidin-4-yl]-propyl}-phenyl)-2-phenoxy-propionic acid**

#### Step A



15 A solution of crude 3-methyl-1-(4-trifluoromethyl-benzyl)-4-[2-(triphenyl- $\square^5$ -phosphanyl)-ethyl]-imidazolidin-2-one iodide (0.65 g, 0.964 mmol) in dry THF (4 mL) is treated with NaH (60% oil suspension, 0.081 g, 2.02 mmol) and stirred at room temperature under  $\text{N}_2$  for 20 minutes to form the ylide. A solution of 3-(4-formyl-phenyl)-2-methyl-2-phenoxy-propionic acid ethyl ester (0.21 g, 0.672 mmol) in THF (4 mL) is then added to the ylide mixture dropwise at room temperature and the reaction is stirred for 16 h. The reaction is diluted with  $\text{Et}_2\text{O}$  and washed with 1 N HCl and water. The organic layer is dried ( $\text{Na}_2\text{SO}_4$ ) and the solvent removed in vacuo to afford crude product that is purified by flash chromatography using 6:1 hexanes:acetone to afford 0.070 g (18%) 2-methyl-3-(4-{3-[3-methyl-2-oxo-1-(4-trifluoromethyl-benzyl)-imidazolidin-4-yl]-propenyl}-phenyl)-2-phenoxy-propionic acid ethyl ester.  $R_f = 0.22$


20

25

30


(1:1 acetone:hexanes).  $^1\text{H}$  NMR. MS ( $\text{ES}^+$ ) Calc'd for  $\text{C}_{33}\text{H}_{36}\text{N}_2\text{O}_4\text{F}_3$  ( $M + 1$ ) 581. Found  $m/z$  581 (100%).

### Step B



5 A mixture of 2-methyl-3-(4-{3-[3-methyl-2-oxo-1-(4-trifluoromethyl-benzyl)-imidazolidin-4-yl]-propenyl}-phenyl)-2-phenoxy-propionic acid ethyl ester (0.069 g, 0.119 mmol) and 10% Pd/C (70 mg) in EtOAc (50 mL) is purged with  
10  $\text{N}_2$  then  $\text{H}_2$  and then stirred under a  $\text{H}_2$  balloon at room temperature for 2 h. The reaction mixture is filtered through hyflo and the solvent removed *in vacuo* to afford 0.069 g (100%) 2-methyl-3-(4-{3-[3-methyl-2-oxo-1-(4-trifluoromethyl-benzyl)-imidazolidin-4-yl]-propyl}-phenyl)-  
15 2-phenoxy-propionic acid ethyl ester.  $^1\text{H}$  NMR. MS ( $\text{ES}^+$ ) Calc'd for  $\text{C}_{33}\text{H}_{38}\text{N}_2\text{O}_4\text{F}_3$  ( $M + 1$ ) 583. Found  $m/z$  583 (100%).

### Step C




20 A solution of 2-methyl-3-(4-{3-[3-methyl-2-oxo-1-(4-trifluoromethyl-benzyl)-imidazolidin-4-yl]-propyl}-phenyl)-2-phenoxy-propionic acid ethyl ester (0.062 g, 0.106 mmol) in ethanol (8 mL) is treated with aqueous 5 N NaOH (1 mL) and heated to reflux 1 h. The reaction mixture is cooled,  
25 the solvent removed *in vacuo*. The resultant residue is acidified with aqueous 1 N HCl (20 mL) and extracted with

CH<sub>2</sub>Cl<sub>2</sub>. The organic layer is dried (Na<sub>2</sub>SO<sub>4</sub>) and the solvent removed *in vacuo* to afford 0.062 g (100%) 2-methyl-3-(4-{3-[3-methyl-2-oxo-1-(4-trifluoromethyl-benzyl)-imidazolidin-4-yl]-propyl}-phenyl)-2-phenoxy-propionic acid. <sup>1</sup>H NMR. HRMS (ES<sup>+</sup>) *m/z* exact mass calc'd for C<sub>31</sub>H<sub>34</sub>N<sub>2</sub>O<sub>4</sub>F<sub>3</sub> (M +1) 555.2471. Found *m/z* 555.2459.

#### Example 101

3-(2-(1,3-Dioxo-1,3-dihydro-isoindol-2-ylmethyl)-4-{2-[3-methyl-2-oxo-1-(4-trifluoromethyl-benzyl)-imidazolidin-4-yl]-ethoxy}-phenyl)-propionic acid



#### Step A

3-(2-(1,3-Dioxo-1,3-dihydro-isoindol-2-ylmethyl)-4-{2-[3-methyl-2-oxo-1-(4-trifluoromethyl-benzyl)-imidazolidin-4-yl]-ethoxy}-phenyl)-propionic acid tert-butyl ester

To a solution of 3-[2-(1,3-Dioxo-1,3-dihydro-isoindol-2-ylmethyl)-4-hydroxy-phenyl]-propionic acid tert-butyl ester (57 mg, 0.15 mmol) and Toluene-4-sulfonic acid 2-[3-methyl-2-oxo-1-(4-trifluoromethyl-benzyl)-imidazolidin-4-yl]-ethyl ester (75 mg, 0.16 mmol) in DMF (1 ml) is added Cs<sub>2</sub>CO<sub>3</sub> (64 mg, 0.19 mmol), and the mixture heated at 60° C for 14 hours. The mixture is cooled, diluted with water (25 ml) and extracted with ethyl acetate (2 x 25 ml). The combined

organic extracts were washed with brine (30 ml), dried (Na<sub>2</sub>SO<sub>4</sub>), and concentrated to an oil. Purification by flash chromatography on a Biotage silica cartridge (gradient elution, 5:1 to 1:1 hexanes:acetone) produced an oil (72 mg, 73%).

MS [EI+] 666 (M+H)<sup>+</sup>. Structure confirmed by 1H-NMR.

#### Step B


3-(2-(1,3-Dioxo-1,3-dihydro-isoindol-2-ylmethyl)-4-{2-[3-methyl-2-oxo-1-(4-trifluoromethyl-benzyl)-imidazolidin-4-yl]-ethoxy}-phenyl)-propionic acid

A solution of the ester obtained in Step A (70mg, 0.10 mmol) in trifluoroacetic (1 ml) and methylene chloride (2 ml) is stirred at 23° C for 7 hours. Concentration of the reaction mixture gave an oil which is partitioned between ethyl acetate (20 ml) and water (25 ml). The ethyl acetate layer is washed with water (15 ml), brine (20 ml), dried (Na<sub>2</sub>SO<sub>4</sub>), and concentrated to an oil which is purified by chromatography on a Biotage silica cartridge (gradient elution, 100% methylene chloride to 10:1 methylene chloride:methanol) to produce an oil (31 mg, 48%). MS [EI+] 610 (M+H)<sup>+</sup>. Structure confirmed by 1H-NMR.

25

#### Example 102

2,2-Dimethyl-3-(4-{2-[3-methyl-2-oxo-1-(4-trifluoromethyl-benzyl)-imidazolidin-4-yl]-ethoxy}-phenyl)-propionic acid



### Step A

2,2-Dimethyl-3-(4-{2-[3-methyl-2-oxo-1-(4-trifluoromethyl-benzyl)-imidazolidin-4-yl]-ethoxy}-phenyl)-propionic acid methyl ester

To a solution of 3-(4-Hydroxy-phenyl)-2,2-dimethyl-propionic acid methyl ester (31 mg, 0.15 mmol) and Toluene-4-sulfonic acid 2-[3-methyl-2-oxo-1-(4-trifluoromethyl-benzyl)-imidazolidin-4-yl]-ethyl ester (75 mg, 0.16 mmol) in DMF (1 ml) is added  $\text{Cs}_2\text{CO}_3$  (64 mg, 0.19 mmol), and the mixture heated at 60° C for 15 hours. The mixture is cooled, diluted with water (25 ml) and extracted with ethyl acetate (2 x 25 ml). The combined organic extracts were washed with brine (25 ml), dried ( $\text{Na}_2\text{SO}_4$ ), and concentrated to an oil. Purification by flash chromatography on a Biotage silica cartridge (gradient elution, 2:1 hexanes:ethyl acetate to 1:4 hexanes:ethyl acetate) produced an oil (64 mg, 88%). MS [EI+] 493 (M+H)<sup>+</sup>. Structure confirmed by 1H-NMR.



### Step B

2,2-Dimethyl-3-(4-{2-[3-methyl-2-oxo-1-(4-trifluoromethyl-benzyl)-imidazolidin-4-yl]-ethoxy}-phenyl)-propionic acid

To a solution of the ester obtained from Step A (61 mg, 0.12 mmol) in methanol (1.5 ml) is added 5N NaOH (0.3 ml, 1.5 mmol), and the mixture stirred for 20 hours at ambient temperature. The mixture is poured into 1N HCl (20 ml),


then extracted with ethyl acetate (2 x 15 ml). The combined ethyl acetate extracts were washed with water, brine, dried ( $\text{Na}_2\text{SO}_4$ ), and concentrated to a foam (56 mg, 97%. MS [EI+] 479 (M+H)<sup>+</sup>. Structure confirmed by <sup>1</sup>H-NMR.

5

**Example 103**Step A

- 10 The hydantoin from **Example 13** (4.0g, 16.1mmol) and 3,4-dimethylbenzyl chloride (2.5ml, 17.7mmol) were stirred together in DMF (100ml). Potassium carbonate (8.9g, 64.4mmol) and magnesium sulfate (3.0g, 25.0mmol) were added and the resulting mixture was heated to 45 °C for 16 hrs
- 15 under a drying tube. The reaction was cooled, added slowly to 1 N hydrochloric acid (300ml) and extracted twice with ethyl acetate. The organic layers were combined, washed with aqueous brine and dried over sodium sulfate. Evaporation of the solvent and subsequent purification by
- 20 flash chromatography (hexanes:ethyl acetate) gave the desired alkylated hydantoin (1.6g).

$\text{C}_{22}\text{H}_{26}\text{N}_2\text{O}_3$  (MW = 366.5); MS (M<sup>+</sup>, 367.3, M-365.4)




### Step B

Sodium hydride (60% dispersion in mineral oil, 0.1g, 2.4mmol) was suspended in DMF (25ml) and cooled to 0 °C under a drying tube. The product from **Example 103, Step A** (0.8g, 2.2mmol), dissolved in DMF (5ml), was added slowly and allowed to stir at 0 °C for 45 min. Iodopropane (0.23ml, 2.4mmol) added to the reaction and allowed to stir for 2.5 hrs. Hydrochloric acid (1N) added slowly, the reaction was added to water and extracted three times with ethyl acetate. The organic layers were combined, washed with aqueous brine and dried over sodium sulfate. Evaporation of the solvent and subsequent purification by flash chromatography (hexanes:ethyl acetate) gave the desired product (0.7g).

$C_{25}H_{32}N_2O_3$  (MW = 408.6); MS ( $M^+$ , 409.3)

### Step C




Lithium aluminum hydride (0.084g, 2.2mmol) was suspended in THF (10ml) and cooled to 0 °C under a drying tube. The product from **Example 103, Step B** (0.7g, 1.7mmol), dissolved in THF (5ml), was added slowly and stirred for 20 min. Hydrochloric acid (5N, 5ml) added slowly and stirred for 30 min. The reaction mixture was added to water (40ml) and



extracted twice with ethyl acetate. The organic layers were combined, washed with aqueous brine then dried over sodium sulfate. Evaporation of the solvent and subsequent purification by flash chromatography (hexanes:ethyl acetate) gave the desired product (0.53g).

$C_{25}H_{32}N_2O_2$  (MW = 392.6); MS ( $M^+$ , 393.2)


Step D



Boron tribromide was stirred in methylene chloride and cooled to 0 °C. The product from **Example 103, Step C** (0.5g, 1.3mmol) was added and stirred for 30 min. Methanol (2ml) slowly added and the mixture added to water which was extracted twice with methylene chloride. The organic layers were combined, washed with aqueous brine then dried over sodium sulfate. Evaporation of the solvent gave the desired crude product (0.5g).

$C_{24}H_{30}N_2O_2$  (MW = 378.5); MS ( $M^+$ , 379.3)

Step E




The product from **Example 103, Step D** (0.5g, 1.3 mmol) and ethyl 2-bromoisobutyrate (0.57ml, 3.9 mmol) stirred together in ethanol (15ml). Potassium carbonate (powdered, Aldrich, 0.72g, 5.2 mmol) and magnesium sulfate (0.16g, 1.3 mmol)

added and the mixture was stirred at 60 °C overnight. The reaction mixture was cooled and carefully added to 5 N hydrochloric acid (30ml) and the resulting solution was extracted twice with methylene chloride. The organic layers were combined, washed with aqueous brine then dried over sodium sulfate. Evaporation of the solvent and subsequent purification by flash chromatography (hexanes:ethyl acetate) gave the desired product (0.48g).



$C_{30}H_{40}N_2O_4$  (MW = 492.7); MS ( $M^+$ , 493.3)

Step F



The product from **Example 103, Step E** (0.45g, 0.91mmol) was dissolved in dioxane (8ml); lithium hydroxide (0.04g, 1.8mmol, in 2ml water) added and stirred at ambient temperature overnight. The reaction was added to water and washed with ether. The aqueous layer was made acidic with the addition of 5 N hydrochloric acid which was then extracted twice with methylene chloride. These organic layers were combined, washed with aqueous brine and dried over sodium sulfate. Evaporation of the solvent and placement under a vacuum gave the desired product (0.37g)


$C_{28}H_{36}N_2O_4$  (MW = 464.6); MS ( $M^+$ , 465.3,  $M^-$ , 463.4)

**Example 104**Step A

5 The hydantoin from **Example 13** (3.0g, 12.1mmol) and p-tert-butylbenzyl bromide (Aldrich, 2.4ml, 13.3mmol) were stirred together in DMF (75ml). Potassium carbonate (powdered, Aldrich, 6.7g, 48.4mmol) and magnesium sulfate (1.8g, 15.0mmol) were added and the resulting mixture was heated to  
10 50 °C for 4 hrs under a drying tube. The reaction was cooled, added slowly to 5 N hydrochloric acid (100ml) and extracted three times with ethyl acetate. The organic layers were combined, washed with aqueous brine and dried over sodium sulfate. Evaporation of the solvent and  
15 subsequent purification by flash chromatography (hexanes:ethyl acetate) gave the desired alkylated hydantoin (2.8g).

$C_{24}H_{30}N_2O_3$  (MW = 394.5); MS ( $M^+$ , 395.2,  $M^-$ , 393.3)

20


Step B

Sodium hydride (60% dispersion in mineral oil, 0.1g, 2.5mmol) was suspended in DMF (25ml) and cooled to 0 °C

under a drying tube. The product from **Example 104, Step A** (0.9g, 2.3mmol), dissolved in DMF (5ml), was added slowly and allowed to stir at 0 °C for 90 min. Iodomethane (0.15ml, 2.5mmol) added to the reaction and allowed to stir  
5 for 30 min. Hydrochloric acid (5N, 1ml) added slowly to quench the reaction which was then added to water (75ml) and extracted twice with ethyl acetate. The organic layers were combined, washed with aqueous brine then dried over sodium sulfate. Evaporation of the solvent and subsequent  
10 purification by flash chromatography (hexanes:ethyl acetate) gave the desired product (0.82g).

$C_{25}H_{32}N_2O_3$  (MW = 408.6); MS ( $M^+$ , 409.2)


Step C



15 Following the procedure of **Example 103, Step C**, the product from **Example 104, Step B** (0.82g, 2.0mmol) and lithium aluminum hydride (0.114g, 3.00mmol) were used to yield the desired imidazolone as a crude product (0.61g).

20  $C_{25}H_{32}N_2O_2$  (MW = 392.6); MS ( $M^+$ , 393.1)

Step D




The product from **Example 104, Step C** (0.6g, 1.5mmol) was  
25 stirred in methylene chloride (10ml) and boron tribromide (0.28ml, 3.0mmol) added via syringe. The mixture was

stirred at ambient temperature for 30 min. then quenched by the slow addition of methanol. The mixture was added to water (50ml) and extracted twice with methylene chloride. The organic layers were combined, washed with aqueous brine and dried over sodium sulfate. Evaporation of the solvent

gave the desired crude product (0.51g).  
 $C_{24}H_{30}N_2O_2$  (MW = 378.5); MS ( $M^+$ , 379.2)


#### Step E



Following the procedure of **Example 103, Step E**, the product was from **Example 104, Step D** (0.5g, 1.3 mmol), ethyl 2-bromoisobutyrate (0.57ml, 3.9 mmol), potassium carbonate (powdered, Aldrich, 0.72g, 5.2 mmol) and magnesium sulfate (0.16g, 1.3 mmol) were used to yield the desired product


(0.13g).  
 $C_{30}H_{40}N_2O_4$  (MW = 492.7); MS ( $M^+$ , 493.3)

#### Step F




Following the procedure of **Example 103, Step F**, the product from **Example 104, Step E** (0.12g, 0.24mmol) and lithium hydroxide (0.02g, 0.72mmol) were used to yield the desired product after evaporation of the solvent and placement under vacuum (0.089g).

$C_{28}H_{36}N_2O_4$  (MW = 464.6); MS ( $M^+$ , 465.1,  $M^-$ , 463.2)


**Example 105**

5

Step A

Following the procedure of **Example 104, Step B**, the product from **Example 104, Step A** (0.9g, 2.3mmol), sodium hydride (60% dispersion in mineral oil, 0.1g, 2.5mmol) and iodoethane (0.2ml, 2.5mmol) were used to yield the desired product (0.68g).


$C_{26}H_{34}N_2O_3$  (MW = 422.6); MS ( $M^+$ , 423.2)

Step B

15


Following the procedure of **Example 103, Step C**, the product from **Example 105, Step A** (0.68g, 1.6mmol) and lithium aluminum hydride (0.091g, 2.4mmol) were used to yield the desired imidazolone as a crude product (0.6g).

20  $C_{26}H_{34}N_2O_2$  (MW = 406.6); MS ( $M^+$ , 407.1)

Step C


Following the procedure of **Example 104, Step D**, the product from **Example 105, Step B** (0.6g, 1.5mmol) and boron tribromide (0.28ml, 3.0mmol) were used to yield the crude product (0.52g).

$C_{25}H_{32}N_2O_2$  (MW = 392.6); MS ( $M^+$ , 393.2)

Step D

Following the procedure of **Example 103, Step E**, the product from **Example 105, Step C** (0.5g, 1.3 mmol), ethyl 2-bromoisobutyrate (0.57ml, 3.9 mmol), potassium carbonate (powdered, Aldrich, 0.72g, 5.2 mmol) and magnesium sulfate (0.16g, 1.3 mmol) were used to yield the product (0.31g).

$C_{31}H_{42}N_2O_4$  (MW = 506.7); MS ( $M^+$ , 507.1)



Step E

Following the procedure of **Example 103, Step F**, the product from **Example 105, Step D** (0.30g, 0.59mmol) and lithium hydroxide (0.04g, 1.7mmol) were used to yield the desired

product as a foam after evaporation of the solvent and placement under vacuum (0.244g).

$C_{29}H_{38}N_2O_4$  (MW = 478.6); MS ( $M^+$ , 479.1,  $M^-$ , 477.2)


5

**Example 106**Step A

Following the procedure of **Example 104, Step B**, the product from **Example 104, Step A** (0.9g, 2.3mmol), sodium hydride dispersion in mineral oil, 0.1g, 2.5mmol) and iodopropane (0.2ml, 2.5mmol) were used to yield the desired product (0.56g).

$C_{27}H_{36}N_2O_3$  (MW = 436.6); MS ( $M^+$ , 437.2)

15


Step B

Following the procedure of **Example 103, Step C**, the product from **Example 106, Step A** (0.56g, 1.3mmol) and lithium



aluminum hydride (Aldrich, 0.076g, 2.0mmol) were used to yield the desired imidazolone as a crude product (0.51g).  
 $C_{27}H_{36}N_2O_2$  (MW = 420.6); MS ( $M^+$ , 421.1)


5

Step C

Following the procedure of **Example 104, Step D**, the product from **Example 106, Step B** (0.5g, 1.2mmol) and boron tribromide (0.23ml, 2.4mmol) were used to yield the crude product (0.45g).


10

$C_{26}H_{34}N_2O_2$  (MW = 406.6); MS ( $M^+$ , 407.3)

Step D

Following the procedure of **Example 103, Step E**, the product from **Example 106, Step C** (0.45g, 1.1mmol), ethyl 2-bromoisobutyrate (0.57ml, 3.9 mmol), potassium carbonate (0.72g, 5.2 mmol) and magnesium sulfate (0.16g, 1.3 mmol) were used to yield the product (0.26g).



20  $C_{32}H_{44}N_2O_4$  (MW = 520.7); MS ( $M^+$ , 521.3)

Step E

Following the procedure of **Example 103, Step F**, the product from **Example 106, Step D** (0.26g, 0.50mmol) and lithium  
5 hydroxide (0.036g, 1.5mmol) were used to yield the desired product after evaporation of the solvent and placement under vacuum (0.184g).


$C_{30}H_{40}N_2O_4$  (MW = 492.7); MS ( $M^+$ , 493.3,  $M^-$ , 491.4)

10

**Example 107**Step A


The hydantoin from **Example 17, Step A** (490.6 mg, 0.988 mmol) was dissolved in DMF (10 mL) and cooled to 0 °C under an atmosphere of N<sub>2</sub>. The mixture was treated with NaH (60% dispersion in oil, 45.7 mg, 1.14 mmol) and, after 30 minutes, n-propyl iodide (112 µL, 1.15 mmol). The reaction mixture was warmed to room temperature, stirred for 1 hour, then poured into 1 N HCl (50 mL). The resulting solution was extracted with ethyl acetate (2x50 mL). The combined organic extracts were dried over Na<sub>2</sub>SO<sub>4</sub> and concentrated. Purification by chromatography (4:1 hexanes:ethyl acetate) gave the desired product as a colorless oil (416.1 mg, 78%). C<sub>31</sub>H<sub>42</sub>N<sub>2</sub>O<sub>6</sub> (MW = 538.69); mass spectroscopy: (M+NH<sub>4</sub><sup>+</sup>) = 556.3

### Step B




15 A CH<sub>2</sub>Cl<sub>2</sub> solution (15 mL) of the ester from **Step A** (0.411 g, 0.764 mmol) was cooled to 0 °C and treated with TFA (2.0 mL, 26 mmol). The mixture was warmed to room temperature and stirred for 3 hours. The solvent was concentrated to give  
20 the crude acid, which was used in the subsequent reaction without further purification.


C<sub>27</sub>H<sub>34</sub>N<sub>2</sub>O<sub>6</sub> (MW = 482.58); mass spectroscopy: (MH<sup>+</sup>) = 483.2, (MH<sup>-</sup>) = 481.3

Step C


The hydantoin from **Step B** was dissolved in ethanol (15 mL) and treated with  $\text{NaBH}_4$  (331 mg, 8.8 mmol). Additional  $\text{NaBH}_4$  was added to the reaction mixture after 5 hr (150 mg, 4.0 mmol). The resulting mixture was stirred at room temperature overnight then quenched by the careful addition of 5 N HCl (5 mL). The mixture was stirred for 30 minutes then diluted with additional 5 N HCl (50 mL) and extracted with ethyl acetate (3x50 mL). The combined organic extracts were washed with brine, dried over  $\text{Na}_2\text{SO}_4$  and concentrated. Purification by chromatography (gradient: 2:1 ethyl acetate:hexane to 100% ethyl acetate to 9:1 ethyl acetate:methanol) gave the desired imidazolone as a white solid (237 mg, 66% for two steps).

$\text{C}_{27}\text{H}_{34}\text{N}_2\text{O}_5$  (MW = 466.58); mass spectroscopy:  $(\text{MH}^+) = 467.3$ ,  $(\text{MH}^-) = 465.3$

**Example 108**

Step A

The ester from **Example 14, Step F** (1.88 g, 0.0050 mol) was dissolved in DMF and treated with  $\alpha$ -chloro-p-xylene (0.773 g, 0.0055 mol) and powdered  $K_2CO_3$ . The resulting mixture was stirred at room temperature overnight. The reaction mixture was poured into 1N HCl and combined with ethyl acetate. The organic layer was extracted and washed with brine then dried and concentrated. Purification of the crude material by flash chromatography (4:1 hexanes: ethyl acetate) yielded the desired hydantoin (2.10 g, 88 %).  $C_{28}H_{36}N_2O_5$  (MW = 480.26); mass spectroscopy ( $MH^+$ ) = 480.1

Step B


15

Sodium hydride (0.050 g, 0.0011 mol) was dissolved in DMF (5 ml) and cooled to 0°C. The hydantoin from **Step A** (0.50 g, 0.0010 mol) was added as a solution in DMF (10 ml). The mixture was stirred for ninety minutes. Iodomethane (0.068 ml, 0.0011 mol) was added and the reaction was stirred for thirty minutes. Hydrochloric acid was added to quench the reaction. The aqueous layer was extracted with ethyl

20

acetate. The organic layer was concentrated. Purification of the resulting material by flash chromatography (10:1 hexanes: ethyl acetate) gave the desired hydantoin (0.247 g, 0.0010 mol).  $C_{29}H_{38}N_2O_5$  (MW = 494.28); mass spectroscopy  
5  $(MH^+) = 494.0$


### Step C



The hydantoin from **Step B** (0.247 g, 0.00050 mol) was  
10 dissolved in methylene chloride (5 ml) and treated with trifluoro acetic acid (0.269 ml, 0.0035 mol) and stirred overnight. The solvent was concentrated and the product was vacuum dried.

$C_{25}H_{30}N_2O_5$  (MW = 438.22); mass spectroscopy  $(MH^+) = 439$   
15


### Step D




The acid from **Step C** (0.289, 0.00066 mol) was dissolved in ethanol (10 ml) and treated with sodium borohydride (0.249 g, 0.0066 mol). One hour later, additional sodium  
20 borohydride (0.249 g) was added and the reaction was stirred overnight. On the next day, more sodium borohydride (0.249

g) was added to drive the reaction. The reaction was left to stir. On the fifth day, 5N HCl (50 ml) was added to the reaction mixture followed by water (50 ml). The aqueous solution was extracted with ethyl acetate (2X, 50 ml). The organic layer was washed with brine, dried over Na<sub>2</sub>SO<sub>4</sub>, and concentrated. Purification by flash chromatography (100% ethyl acetate) gave the desired acid as a white solid (0.062 g, 23 %) C<sub>25</sub>H<sub>30</sub>N<sub>2</sub>O<sub>4</sub> (MW = 422.53); mass spectroscopy (MH<sup>+</sup>) = 423.2

#### Example 109



#### Step A




The hydantoin from **Example 13** (1.0g, 4.0mmol), 2-(chloromethyl)quinoline hydrochloride (0.9g, 4.4mmol) and triethylamine (0.4g, 4.0mmol) were combined in DMF (25ml). Potassium carbonate (2.2g, 16.0mmol) and magnesium sulfate (0.6g, 5.0mmol) added and the mixture was heated to 50 °C, under a drying tube, overnight. The reaction mixture was added to hydrochloric acid (1N, 50ml) and extracted three times with methylene chloride. The organic layers were combined, washed with aqueous brine then dried over sodium sulfate. Evaporation of the solvent and washing with 7:3 hexane/ethyl acetate yielded the desired product as a solid

(0.8g)

$C_{23}H_{23}N_3O_3$  (MW = 389.5); MS ( $M^+$ , 390.2)

Step B




5

Following the procedure of **Example 104, Step B**, the product from **Example 109, Step A** (0.5g, 1.3mmol), sodium hydride (Aldrich, 60% dispersion in mineral oil, 0.06g, 1.4mmol) and iodopropane (Aldrich, 0.14ml, 1.4mmol) were used to yield the desired product (0.41g).

10

$C_{26}H_{29}N_3O_3$  (MW = 431.5); MS ( $M^+$ , 432.1)

Step C




15 Following the procedure of **Example 103, Step C**, the product from **Example 109, Step B** (0.4g, 0.93mmol) and lithium aluminum hydride (0.05g, 1.4mmol) were used to yield the desired imidazolone as a crude product (0.45g).

$C_{26}H_{29}N_3O_2$  (MW = 415.5); MS ( $M^+$ , 416.2)


20



Step D


Following the procedure of **Example 104, Step D**, the product from **Example 109, Step C** (0.39g, 0.93mmol) and boron tribromide (0.19ml, 2.0mmol) were used to yield the crude product (0.47g).

$C_{25}H_{27}N_3O_2$  (MW = 401.5); MS ( $M^+$ , 402.2)

Step E

Following the procedure of **Example 103, Step E**, the product from **Example 109, Step D** (0.37g, 0.93mmol), ethyl 2-bromoisobutyrate (0.48ml, 3.3 mmol), potassium carbonate (powdered, Aldrich, 0.58g, 4.2 mmol) and magnesium sulfate (0.18g, 1.5 mmol) were used to yield the product (0.05g).

$C_{31}H_{37}N_3O_4$  (MW = 515.7); MS ( $M^+$ , 516.3)



Step F

Following the procedure of **Example 103, Step F**, the product from **Example 109, Step E** (0.05g, 0.1mmol) and lithium


hydroxide (0.07g, 0.3mmol) were used to yield the desired product (0.022g).

$C_{29}H_{33}N_3O_4$  (MW = 487.6); MS ( $M^+$ , 488.3,  $M^-$ , 486.4)


5

**Example 110**Step A

- 10 The methoxy ether from **Example 13** (3.0 g, 0.012 mol) was dissolved in DMF and treated with *p*-methyl chlorobenzene (1.87 g, 0.013 mol) and powdered  $K_2CO_3$  (6.62 g, 0.048 mol). The resulting mixture was stirred at room temperature overnight. The reaction mixture was poured into 1N HCl and
- 15 combined with ethyl acetate. The organic layer was extracted and washed with brine then dried and concentrated. Purification of the crude material by flash chromatography (2:1 hexanes: ethyl acetate) yielded the desired hydantoin (3.15 g, 75 %).  $C_{21}H_{24}N_2O_3$  (MW =352.18); mass spectroscopy
- 20 ( $MH^+$ ) = 353

Step B


The hydantoin from **Step A** (0.750 g, 0.0021 mol) was dissolved in DMF (5 ml) and treated with NaH (0.091 g, 0.0023 mol) followed by 1-iodo-propane (0.228 ml, 0.0023 mol). The reaction was stirred overnight under nitrogen. The reaction mixture was poured into 1N HCl and combined with ethyl acetate. The organic layer was extracted and washed with brine then dried and concentrated. Purification of the crude material by flash chromatography (4:1 hexanes: ethyl acetate) yielded the desired product (0.454 g, 55 %).  
 $C_{24}H_{30}N_2O_3$  (MW = 394.52); mass spectroscopy (MH<sup>+</sup>) = 395.2

Step C


Lithium aluminum hydride (0.0651 g, 0.0017 mol) was dissolved in THF (5 ml). A THF solution of the hydantoin from **Step B** was added. The reaction was stirred overnight at room temperature. The reaction was quenched by the addition of 5N HCl. After stirring for thirty minutes, water was

added and the solution was extracted with ethyl acetate. The organic layer was extracted and washed with brine then dried and concentrated. The crude product was carried forth without further purification (0.400 g, 96 %).  $C_{24}H_{30}N_2O_2$  (MW = 378.23); mass spectroscopy (MH<sup>+</sup>) = 379.2

Step D




The methoxy ether from **Step C** (0.400 g, 0.001 mol) was dissolved in methylene chloride (5 ml) and cooled to 0°C. To this solution was added, dropwise, a solution of BBr<sub>3</sub> (0.200 ml, 0.002 mol) in methylene chloride (5 ml). After stirring for about thirty minutes, the reaction mixture was cooled to 0°C and quenched by the dropwise addition of methanol/methylene chloride. The solvent was concentrated and the resulting material was dissolved in methylene chloride. The organic layer was extracted with water followed by brine. Upon evaporation of the solvent, the phenol (0.334 g, 92%) was obtained and was carried forth without further purification.  $C_{23}H_{28}N_2O_2$  (MW = 364.22); mass spectroscopy (MH<sup>+</sup>) = 365.2

Step E

The phenol from **Step D** (0.334 g, 0.00092 mol) was dissolved in EtOH (5 ml) and treated with ethyl 2-bromoisobutyrate  
5 (0.404 ml, 0.0028 mol), powdered  $K_2CO_3$  (0.508 g, 0.0037 mol), and  $MgSO_4$  (0.110 g, 0.00092 mol). The reaction was stirred overnight at 55°C. Upon cooling, the reaction mixture was poured into 5N HCl and combined with EtOAc. The organic layer was extracted with water followed by brine then  
10 concentrated to dryness. Purification by flashed chromatography (1:1 hexanes: ethyl acetate) gave the ester (0.205 g, 46 %).  $C_{29}H_{38}N_2O_4$  (MW = 478.28); mass spectroscopy ( $MH^+$ ) = 479.3


15

Step F


The ester from **Step E** (0.205 g, 0.00043 mol) was dissolved in methanol (4 ml) and treated with a solution of LiOH in water (1 ml). The reaction was stirred overnight. The  
20 reaction was cooled and water (20 ml) was added to the

solution. The solution was then acidified using 1N HCl to pH = 3 then extracted with ethyl acetate. The organic layer was concentrated to afford the desired carboxylic acid (0.128 g, 66 %).  $C_{27}H_{34}N_2O_4$  (MW = 450.58); mass spectroscopy (MH<sup>+</sup>) = 451.2

### Example 111



### Step A



10


The hydantoin from **Example 110; Step A** (0.748 g, 0.0021 mol) was dissolved in DMF (5 ml) and treated with NaH (0.091 g, 0.0023 mol) followed by 1-iodo-hexane (0.339 ml, 0.0023 mol). The reaction was stirred overnight under nitrogen. The reaction mixture was poured into 1N HCl and combined with ethyl acetate. The organic layer was extracted and washed with brine then dried and concentrated. Purification of the

15


crude material by flash chromatography (5:1 hexanes: ethyl acetate) yielded the desired product (0.622 g, 68 %).

$C_{27}H_{32}F_2N_2O_5$  (MW =502.23); mass spectroscopy (MH+) = 447.1

5

Step B

Lithium aluminum hydride (0.622 g, 0.0014 mol) was dissolved in THF (10 ml). A THF solution of the hydantoin from **Step A** was added. The reaction was stirred for thirty minutes at room temperature. The reaction was quenched by the addition of 5N HCl. After stirring for thirty minutes, water was added and the solution was extracted with ethyl acetate. The organic layer was extracted and washed with brine then dried and concentrated. The crude product was carried forth without further purification (0.556 g, 95 %).  $C_{27}H_{32}F_2N_2O_5$  (MW =502.23); mass spectroscopy (MH+) = 447.1

Step C


The methoxy ether from **Step B** (0.556 g, 0.0013 mol) was dissolved in methylene chloride (7 ml) and cooled to 0°C.

5 To this solution was added, dropwise, a solution of BBr<sub>3</sub> (0.250 ml, 0.0026 mol) in methylene chloride (5 ml). After stirring for about twenty minutes, the reaction mixture was cooled to 0°C and quenched by the dropwise addition of methanol/methylene chloride. The solvent was concentrated


10 and the resulting material was dissolved in methylene chloride. The organic layer was extracted with water followed by brine. Upon evaporation of the solvent, the phenol (0.465 g, 88%) was obtained and was carried forth without further purification. C<sub>22</sub>H<sub>27</sub>N<sub>3</sub>O<sub>2</sub> (MW = 365.48); mass

15 spectroscopy (MH<sup>+</sup>) = 366.3



Step D



The phenol from **Step C** (0.465 g, 0.0011 mol) was dissolved in EtOH (10 ml) and treated with ethyl 2-bromoisobutyrate (0.504 ml, 0.0034 mol), powdered  $K_2CO_3$  (0.607 g, 0.0044 mol), and  $MgSO_4$  (0.132 g, 0.0011 mol). The reaction was stirred overnight at 55°C. Upon cooling, the reaction mixture was poured into 5N HCl and combined with EtOAc. The organic layer was extracted with water followed by brine then concentrated to dryness. Purification by flashed chromatography (3:1 hexanes: ethyl acetate; 2:1 hexanes: ethyl acetate ) gave the ester (0.241 g, 42 %).  $C_{28}H_{37}N_3O_4$  (MW = 479.62); mass spectroscopy ( $MH^+$ ) = 480.3

Step E

The ester from **Step D** (0.241 g, 0.00046 mol) was dissolved

in methanol (4 ml) and treated with a solution of LiOH in water (1 ml). The reaction was stirred overnight. The reaction was cooled and water (20 ml) was added to the solution. The solution was then acidified using 1N HCl to pH = 3 then extracted with ethyl acetate. The organic layer was concentrated to afford the desired carboxylic acid (0.089 g, 40 %).  $C_{26}H_{33}N_3O_4$  (MW = 451.57); mass spectroscopy ( $MH^+$ ) = 452.3

10

**Example 112**Step A


A DMF solution (55 mL) of the hydantoin from **Example 13** (1.92 g, 7.73 mmol) was treated sequentially with 2-bromomethyl-1-naphthalene (1.87 g, 8.46 mmol),  $K_2CO_3$  (2.2 g, 16 mmol), and  $MgSO_4$  (2.3 g, 19 mmol). The resulting mixture was stirred at room temperature overnight. The reaction was quenched by the slow addition of 1 N HCl (150 mL). The resulting mixture was extracted with ethyl acetate (2x100 mL). The combined organic extracts were washed with brine then dried over  $Na_2SO_4$  and concentrated. Purification by

chromatography (gradient: 3:1 to 1:1 hexanes:ethyl acetate) gave the desired alkylated hydantoin as a white solid (1.47 g, 49%).

$C_{24}H_{24}N_2O_3$  (MW = 388.47); mass spectroscopy:  $(MH^+) = 389.2$ ,


5  $(MH^-) = 387.2$

Step B




The hydantoin from **Step A** (637 mg, 1.64 mmol) was dissolved  
10 in DMF (20 mL) and cooled to 0 °C under an atmosphere of  $N_2$ .  
The mixture was treated with NaH (60% dispersion in oil,  
76.3 mg, 1.91 mmol) and, after 15 minutes, n-propyl iodide  
(192  $\mu$ L, 1.97 mmol). The reaction mixture was warmed to  
room temperature, stirred for 1 hour, then poured into 1 N  
15 HCl (100 mL). The resulting solution was extracted with  
ethyl acetate (2x75 mL). The combined organic extracts were  
dried over  $Na_2SO_4$  and concentrated. Purification by  
chromatography (4:1 hexanes:ethyl acetate) gave the desired  
product as a yellowish oil (693 mg, 98%).

20  $C_{27}H_{30}N_2O_3$  (MW = 430.55); mass spectroscopy:  $(MH^+) = 431.2$

Step C

To a slurry of LAH (99.3 mg, 2.6 mmol) in THF (10 mL) at 0 °C under N<sub>2</sub> was added the hydantoin from **Step B** (0.693 g, 1.61 mmol) as a solution in THF (10 mL). After 15 minutes the reaction was quenched by the addition of 5 N HCl (5 mL) in THF (5 mL), stirred for 30 minutes, then diluted with H<sub>2</sub>O (75 mL). The resulting mixture was extracted with ethyl acetate (3x50 mL). The combined organic extracts were washed with brine, dried over Na<sub>2</sub>SO<sub>4</sub> and concentrated to give the desired imidazolone. The crude product was carried forward without further purification.

C<sub>27</sub>H<sub>30</sub>N<sub>2</sub>O<sub>2</sub> (MW = 414.55); mass spectroscopy: (MH<sup>+</sup>) = 415.2


Step D

The imidazolone from **Step C** was dissolved in CH<sub>2</sub>Cl<sub>2</sub> (15 mL) and cooled to 0 °C under an atmosphere of N<sub>2</sub>. BBr<sub>3</sub> (475 µL, 5.0 mmol) was added dropwise, then the reaction mixture was warmed to room temperature. After 1 hr, the solution was again cooled to 0 °C and quenched by the slow addition of a methanol (4 mL) solution in CH<sub>2</sub>Cl<sub>2</sub> (12 mL). The resulting

mixture was extracted with H<sub>2</sub>O (50 mL). The aqueous extract was washed with CH<sub>2</sub>Cl<sub>2</sub> (50 mL) then the combined organic extracts were washed with brine, dried over Na<sub>2</sub>SO<sub>4</sub>, and concentrated to give the desired phenol as an oil. The crude product was carried forward without further purification.

C<sub>26</sub>H<sub>28</sub>N<sub>2</sub>O<sub>2</sub> (MW = 400.53); mass spectroscopy: (MH<sup>+</sup>) = 401.2, (MH<sup>-</sup>) = 399.3

Step E




10

An ethanol solution (15 mL) of the phenol from **Step D** was treated sequentially with ethyl 2-bromoisobutyrate (900  $\mu$ L, 6.6 mmol), K<sub>2</sub>CO<sub>3</sub> (1.0 g, 7.2 mmol), and MgSO<sub>4</sub> (1.0 g, 8.3 mmol). The resulting mixture was heated to 55-65 °C overnight. The reaction was quenched by the slow addition of 1 N HCl (5 mL) then poured into additional 1 N HCl (50 mL). The resulting mixture was extracted with CH<sub>2</sub>Cl<sub>2</sub> (2x50 mL). The combined organic extracts were washed with brine then dried over Na<sub>2</sub>SO<sub>4</sub> and concentrated. Purification by chromatography (gradient: 2:1 to 1:1 hexanes:ethyl acetate) gave the desired ester as a slightly yellow oil (536.4 mg, 65% for 3 steps).


20

C<sub>32</sub>H<sub>38</sub>N<sub>2</sub>O<sub>4</sub> (MW = 514.67); mass spectroscopy: (MH<sup>+</sup>) = 515.4


Step F

A dioxane solution (10 mL) of the ester from **Step E** (516.5 mg, 1.0 mmol) was treated with an aqueous solution (5 mL) of  
5 LiOH (88 mg, 3.7 mmol). The mixture was heated to 50 °C for 2 hr then the solvent was concentrated and the resulting oil diluted with H<sub>2</sub>O (50 mL) and extracted with Et<sub>2</sub>O (50 mL). The aqueous extract was acidified with 1 N HCl and extracted with ethyl acetate (2x50 mL). The combined organic extracts  
10 were washed with brine, dried over Na<sub>2</sub>SO<sub>4</sub>, and concentrated to give the desired carboxylic acid as a slightly yellow foam-like solid (392.8 mg, 81%)  
C<sub>30</sub>H<sub>34</sub>N<sub>2</sub>O<sub>4</sub> (MW = 486.62); mass spectroscopy: (MH<sup>+</sup>) = 487.2, (MH<sup>-</sup>) = 485.4

15


**Example 113**

20

Step A

The methoxy ether from **Example 13** (1.0 g, 0.0040 mol) was dissolved in DMF and treated with 3, 5 difluoro  
5 bromobenzene (0.574 ml, 0.0044 mol) and powdered  $K_2CO_3$  (2.20 g, 0.0080 mol). The resulting mixture was stirred at room temperature overnight. The reaction mixture was poured into 1N HCl and combined with ethyl acetate. The organic layer was extracted and washed with brine then dried and  
10 concentrated. Purification of the crude material by flash chromatography (4:1 hexanes: ethyl acetate; 4:1 hexanes: ethyl acetate) yielded the desired hydantoin (0.710 g, 46 %).  $C_{20}H_{20}F_2N_2O_3$  (MW = 366.414); mass spectroscopy (MH<sup>+</sup>) = 375.1

15


Step B

The hydantoin from **Step A** (0.710 g, 0.0019 mol) was dissolved in DMF (10 ml) and treated with NaH (0.084 g, 0.0021 mol) followed by 1-iodo-propane (0.200 ml, 0.0021  
20 mol). The reaction was stirred overnight under nitrogen. The reaction mixture was poured into 1N HCl and combined with

ethyl acetate. The organic layer was extracted and washed with brine then dried and concentrated. Purification of the crude material by flash chromatography (2:1 hexanes: ethyl acetate) yielded the desired product (0.742 g, 94 %).


5  $C_{23}H_{26}F_2N_2O_3$  (MW =416.19); mass spectroscopy (MH+) = 417

Step C




Lithium aluminum hydride (0.742 g, 0.0018 mol) was dissolved  
10 in THF (7 ml). A THF solution of the hydantoin from **Step B**  
was added. The reaction was stirred for ninety minutes at  
room temperature. The reaction was quenched by the addition  
of 5N HCl. After stirring for thirty minutes, water was  
15 added and the solution was extracted with ethyl acetate. The  
organic layer was extracted and washed with brine then dried  
and concentrated. The crude product was carried forth  
without further purification (0.690 g, 96 %).  $C_{23}H_{26}F_2N_2O_2$  (MW  
=400.20); mass spectroscopy (MH+) = 401



Step D

The methoxy ether from **Step C** (0.690 g, 0.0017 mol) was dissolved in methylene chloride and cooled to 0°C. To this solution was added, dropwise, a solution of BBr<sub>3</sub> (0.326 ml, 0.0035 mol) in methylene chloride. After stirring for about thirty minutes, the reaction mixture was cooled to 0°C and quenched by the dropwise addition of methanol/methylene chloride. The solvent was concentrated and the resulting material was dissolved in methylene chloride. The organic layer was extracted with water followed by brine. Upon evaporation of the solvent, the phenol (0.550 g, 84%) was obtained and was carried forth without further purification. C<sub>22</sub>H<sub>24</sub>F<sub>2</sub>N<sub>2</sub>O<sub>2</sub> (MW = 386.18); mass spectroscopy (MH<sup>+</sup>) = 387.2

15


Step E

The phenol from **Step D** (0.055 g, 0.0014 mol) was dissolved in EtOH (7 ml) and treated with ethyl 2-bromoisobutyrate (0.627 ml, 0.0043 mol), powdered K<sub>2</sub>CO<sub>3</sub> (0.773 g, 0.0056 mol),

20

and  $\text{MgSO}_4$  (0.168 g, 0.0014 mol). The reaction was stirred overnight at  $77^\circ\text{C}$ . Upon cooling, the reaction mixture was poured into 5N HCl and combined with EtOAc. The organic layer was extracted with water followed by brine then concentrated to dryness. Purification by flashed chromatography (1:1 hexanes: ethyl acetate) gave the ester (0.135 g, 19 %).  $\text{C}_{28}\text{H}_{34}\text{F}_2\text{N}_2\text{O}_4$  (MW = 500.25); mass spectroscopy ( $\text{MH}^+$ ) = 501.2

#### Step F





10

The ester from **Step D** (0.135 g, 0.00027 mol) was dissolved in methanol (4 ml) and treated with a solution of LiOH in water (1 ml). The reaction was stirred overnight. The reaction was cooled and water (20 ml) was added to the solution. The solution was then acidified using 1N HCl to pH = 3 then extracted with ethyl acetate. The organic layer was concentrated to afford the desired carboxylic acid.  $\text{C}_{26}\text{H}_{30}\text{F}_2\text{N}_2\text{O}_4$  (MW = 472.22); mass spectroscopy ( $\text{MH}^+$ ) = 473.1

20

#### Example 114




Step A

Following the procedure of **Example 104, Step B**, the product from **Example 103, Step A** (1.0g, 2.7mmol), sodium hydride (Aldrich, 60% dispersion in mineral oil, 0.12g, 3.0mmol) and (bromomethyl)cyclopropane (Aldrich, 0.29ml, 3.0mmol) were used to yield the desired product (1.05g).


$C_{26}H_{32}N_2O_3$  (MW = 420.6); MS ( $M^+$ , 421.2)

10

Step B

Following the procedure of **Example 103, Step C**, the product from **Example 114, Step A** (1.0g, 2.4mmol) and lithium aluminum hydride (Aldrich, 0.14g, 3.6mmol) were used to yield the desired imidazolone as a crude product (0.94g).

$C_{26}H_{32}N_2O_2$  (MW = 404.6); MS ( $M^+$ , 405.2)


Step C

20

Following the procedure of **Example 104, Step D**, the product from **Example 114, Step B** (0.9g, 2.2mmol) and boron tribromide (Aldrich, 0.47ml, 5.0mmol) were used to yield the crude product (0.9g).

5  $C_{25}H_{30}N_2O_2$  (MW = 390.5); MS ( $M^+$ , 391.2,  $M^-$ , 389.4)


Step D



10 Following the procedure of **Example 103, Step E**, the product from **Example 114, Step C** (0.8g, 2.0 mmol), ethyl 2-bromoisobutyrate (Aldrich, 0.9ml, 6.0 mmol), potassium carbonate (powdered, Aldrich, 1.1g, 8.0 mmol) and magnesium sulfate (Mallinkrodt, 0.3g, 2.5 mmol) were used to yield the product (0.7g).



15  $C_{31}H_{40}N_2O_4$  (MW = 504.7); MS ( $M^+$ , 505.4)

Step E




20 Following the procedure of **Example 103, Step F**, the product from **Example 114, Step D** (0.7g, 1.4mmol), lithium hydroxide (0.07g, 2.8mmol), methanol (8ml) and water (2ml) were used to yield the desired product as a foam after evaporation of the solvent and placement under vacuum (0.295g).


$C_{29}H_{36}N_2O_4$  (MW = 476.6); MS ( $M^+$ , 477.3,  $M^-$ , 475.2)

**Example 115**Step A

- 5 A DMF solution (20 mL) of the hydantoin from **Example 13**  
(960.6 mg, 3.87 mmol) was treated sequentially with benzyl  
bromide (507  $\mu$ L, 4.26 mmol),  $K_2CO_3$  (1.2 g, 8.7 mmol), and  
 $MgSO_4$  (1.2 g, 10.0 mmol). The resulting mixture was stirred  
at room temperature overnight. The reaction was quenched by  
10 the slow addition of 1 N HCl (55 mL). The resulting mixture  
was extracted with ethyl acetate (2x50 mL). The combined  
organic extracts were washed with brine then dried over  
 $Na_2SO_4$  and concentrated. Purification by chromatography  
(gradient: 3:1 to 2:1 hexanes:ethyl acetate) gave the  
15 desired alkylated hydantoin as a white solid (932.3 mg,  
71%).  
 $C_{20}H_{22}N_2O_3$  (MW = 338.41); mass spectroscopy:  $(MH^+) = 339.1$ ,  
 $(MH^-) = 337.2$

Step B

The hydantoin from **Step A** (897.8 mg, 2.65 mmol) was dissolved in DMF (15 mL) and cooled to 0 °C under an atmosphere of N<sub>2</sub>. The mixture was treated with NaH (60% dispersion in oil, 125.5 mg, 3.0 mmol) and, after 10 minutes, n-propyl iodide (297 µL, 3.0 mmol). The reaction mixture was warmed to room temperature, stirred for 45 minutes, then quenched by the addition of H<sub>2</sub>O (0.5 mL). After stirring overnight, the mixture was poured into 1 N HCl (50 mL). The resulting solution was extracted with ethyl acetate (2x50 mL). The combined organic extracts were dried over Na<sub>2</sub>SO<sub>4</sub> and concentrated. Purification by chromatography (gradient: 5:1 to 3:1 hexanes:ethyl acetate) gave the desired product as a yellowish oil (951.7 mg, 94%). C<sub>23</sub>H<sub>28</sub>N<sub>2</sub>O<sub>3</sub> (MW = 380.49); mass spectroscopy: (MH<sup>+</sup>) = 381.1

Step C

The hydantoin from **Step B** (938.4 mg, 2.47 mmol) was dissolved in THF (20 mL) and cooled to 0 °C under an atmosphere of N<sub>2</sub>. LAH (145.7 mg, 3.8 mmol) was added in one


portion and, after 30 minutes, the reaction was quenched by the addition of 5 N HCl (5 mL). The resulting mixture was stirred at room temperature for 30 minutes then diluted with H<sub>2</sub>O (150 mL) and extracted with ethyl acetate (3x50 mL).

- 5 The combined organic extracts were washed with brine, dried over Na<sub>2</sub>SO<sub>4</sub> and concentrated to give the desired imidazolone (871.3 mg, 97%). The crude product was carried forward without further purification.

C<sub>23</sub>H<sub>28</sub>N<sub>2</sub>O<sub>2</sub> (MW = 364.49); mass spectroscopy: (MH<sup>+</sup>) = 365.3

10


Step D



- The imidazolone from **Step C** (845.6 mg, 2.32 mmol) was dissolved in CH<sub>2</sub>Cl<sub>2</sub> (20 mL) and cooled to 0 °C under an atmosphere of N<sub>2</sub>. BBr<sub>3</sub> (1.0 mL, 10.6 mmol) was added dropwise, then the reaction mixture was warmed to room temperature. After 30 min, the solution was again cooled to 0 °C and quenched by the slow addition of a methanol (1 mL) solution in CH<sub>2</sub>Cl<sub>2</sub> (10 mL). Additional methanol:CH<sub>2</sub>Cl<sub>2</sub> (1:1. 15 10 mL) was added and the reaction was warmed to room temperature. The resulting mixture was extracted with H<sub>2</sub>O (50 mL). The aqueous extract was washed with CH<sub>2</sub>Cl<sub>2</sub> (50 mL) then the combined organic extracts were washed with brine, dried over Na<sub>2</sub>SO<sub>4</sub>, and concentrated to give the desired 20 phenol as a foam (782.9 mg, 96%). The crude product was carried forward without further purification. 25


$C_{22}H_{26}N_2O_2$  (MW = 350.46); mass spectroscopy:  $(MH^+)$  = 351.3,  
 $(MH^-)$  = 349.0

Step E



5 An ethanol solution (20 mL) of the phenol from **Step D** (775 mg, 2.21 mmol) was treated sequentially with ethyl 2-bromoisobutyrate (900  $\mu$ L, 6.6 mmol),  $K_2CO_3$  (1.5 g, 10.9 mmol), and  $MgSO_4$  (1.6 g, 13.3 mmol). The resulting mixture was heated to 55-65  $^{\circ}C$  overnight. The reaction was quenched  
10 by the slow addition of 1 N HCl (10 mL) then poured into additional 1 N HCl (50 mL). The resulting mixture was extracted with ethyl acetate (2x50 mL). The combined organic extracts were washed with brine then dried over  $Na_2SO_4$  and concentrated. Purification by chromatography  
15 (gradient: 10:1 to 1:1 hexanes:ethyl acetate) gave the desired ester as a slightly yellow oil (659.0 mg, 64%).  
 $C_{28}H_{36}N_2O_4$  (MW = 464.61); mass spectroscopy:  $(MH^+)$  = 465.2


Step F






A dioxane solution (15 mL) of the ester from **Step E** (633.5 mg, 1.4 mmol) was treated with an aqueous solution (5 mL) of LiOH (100 mg, 4.2 mmol). The mixture was stirred at room temperature overnight. The solvent was concentrated and the resulting oil diluted with H<sub>2</sub>O (50 mL) and extracted with Et<sub>2</sub>O (50 mL). The aqueous extract was acidified with 1 N HCl and extracted with ethyl acetate (2x50 mL). The combined organic extracts were washed with brine, dried over Na<sub>2</sub>SO<sub>4</sub>, and concentrated to give the desired carboxylic acid as a slightly yellow foam-like solid (539.9 mg, 91%)  
C<sub>26</sub>H<sub>32</sub>N<sub>2</sub>O<sub>4</sub> (MW = 436.56); mass spectroscopy: (MH<sup>+</sup>) = 437.3, (MH<sup>-</sup>) = 435.1

#### Example 116



#### Step A




Following the procedure of **Example 104, Step B**, the product from **Example 16, Step A** (0.5g, 1.0mmol), sodium hydride (Aldrich, 60% dispersion in mineral oil, 0.05g, 1.1mmol) and

bromoethylmethyl ether (Aldrich, 0.1ml, 1.1mmol) were used to yield the desired product (0.41g).

$C_{32}H_{44}N_2O_6$  (MW = 552.7);  $^1H$  NMR

5

Step B




Following the procedure of **Example 25, Step B**, the product from **Example 116, Step A** (0.4g, 0.72mmol) and trifluoroacetic acid (2ml) were used to yield the desired product (0.36g).

10


$C_{28}H_{36}N_2O_6$  (MW = 496.6); MS ( $M^+$ , 497.3)

Step C




Following the procedure of **Example 15, Step B**, product from **Example 116, Step B** (0.36g, 0.72mmol) and sodium borohydride (0.54g, 15.4mmol) were used to yield the desired product after flash chromatography (methylene chloride:methanol) (0.032g, 22%).

20  $C_{28}H_{36}N_2O_5$  (MW = 480.6); MS ( $M^+$ , 481.1,  $M^-$ , 479.0)


**Example 117**Step A

- 5 Following the procedure of **Example 104, Step B**, the product from **Example 103, Step A** (1.0g, 2.7mmol), sodium hydride (60% dispersion in mineral oil, 0.12g, 3.0mmol) and iodomethane (0.19ml, 3.0mmol) were used to yield the desired product (1.1g).
- 10  $C_{23}H_{28}N_2O_3$  (MW = 380.5); MS ( $M^+$ , 381.2)

Step B

- 15 Following the procedure of **Example 103, Step C**, the product from **Example 117, Step A** (1.0g, 2.7mmol) and lithium aluminum hydride (0.16g, 4.1mmol) were used to yield the desired imidazolone as a crude product (0.92g).
- $C_{23}H_{28}N_2O_2$  (MW = 364.5); MS ( $M^+$ , 365.2)


20

Step C

Following the procedure of **Example 104, Step D**, the product from **Example 117, Step B** (0.92g, 2.5mmol) and boron tribromide (0.47ml, 5.0mmol) were used to yield the crude product (0.86g).

5  $C_{22}H_{26}N_2O_2$  (MW = 350.5); MS ( $M^+$ , 351.2,  $M^-$ , 349.3)

Step D




Following the procedure of **Example 103, Step E**, the product from **Example 117, Step C** (0.85g, 2.4 mmol), ethyl 2-bromoisobutyrate (1.1ml, 7.2 mmol), potassium carbonate (1.3g, 9.6 mmol) and magnesium sulfate (0.4g, 3.0 mmol) were used to yield the product (0.75g).

10  $C_{28}H_{36}N_2O_4$  (MW = 464.6); MS ( $M^+$ , 465.3)



15

Step E




Following the procedure of **Example 103, Step F**, the product from **Example 117, Step D** (0.04g, 0.04mmol), sodium hydroxide (2N, 1ml) and methanol (4ml) were used to yield the desired product after evaporation of the solvent and placement under vacuum (0.0095g).


20  $C_{26}H_{32}N_2O_4$  (MW = 436.6); MS ( $M^+$ , 437.3,  $M^-$ , 435.1)

**Example 118**Step A

- 5 Following the procedure of **Example 15, Step A**, the hydantoin from **Example 13** (1.0g, 4.0mmol), 3-chlorobenzylbromide 0.67ml, 4.4mmol), potassium carbonate (2.2g, 16.0mmol) and magnesium sulfate (0.6g, 5.0mmol) were used to yield the desired product (0.8g).
- 10  $C_{20}H_{21}N_2O_3Cl$  (MW = 372.9); MS ( $M^+$ , 373.2).


Step B

- Following the procedure of **Example 104, Step B**, the product from **Example 118, Step A** (0.75g, 2.0mmol), sodium hydride (0.12g, 3.0 mmol) and iodopropane (0.29ml, 3.0mmol) were used to yield the desired product (0.55g).
- $C_{23}H_{27}N_2O_3Cl$  (MW = 414.9); MS ( $M^+$ , 415.1)

Step C


Following the procedure of **Example 103, Step C**, the product from **Example 118, Step B** (0.5g, 1.2mmol) and lithium  
5 aluminum hydride (0.07g, 1.8mmol) were used to yield the desired imidazolone as a crude product (0.5g).

$C_{23}H_{27}N_2O_2Cl$  (MW = 389.9); MS ( $M^+$ , 390.2)

Step D

10 Following the procedure of **Example 104, Step D**, the product from **Example 118, Step C** (0.5g, 1.3mmol) and boron tribromide (0.34ml, 3.5mmol) were used to yield the crude product (0.5g).

15  $C_{22}H_{25}N_2O_2Cl$  (MW = 384.9); MS ( $M^+$ , 385.2)


Step E

20 Following the procedure of **Example 103, Step E**, the product from **Example 118, Step D** (0.5g, 1.3mmol), ethyl 2-bromoisobutyrate (0.57ml, 3.9 mmol), potassium carbonate

(0.7g, 5.2 mmol) and magnesium sulfate (0.1g, 1.5 mmol) were used to yield the product (0.21g).

$C_{28}H_{35}N_2O_4Cl$  (MW = 499.1); MS ( $M^+$ , 499.2)


5

Step F


Following the procedure of **Example 103, Step F**, the product from **Example 118, Step E** (0.2g, 0.4mmol), sodium hydroxide (2N, 4ml) and methanol (10ml) were used to yield the desired product after evaporation of the solvent and placement under vacuum (0.163g).

10

$C_{26}H_{31}N_2O_4Cl$  (MW = 471.0);  $^1H$  NMR


**Example 119**

15

Step A


The methoxy ether from **Example 13** (1.0 g, 0.004 mol) was dissolved in DMF and treated with 3,4 difluoro bromobenzene (0.568 ml, 0.0044 mol) and powdered  $K_2CO_3$  (2.20 g, 0.0080 mol). The resulting mixture was stirred at room temperature overnight. The reaction mixture was poured into 1N HCl and combined with ethyl acetate. The organic layer was extracted and washed with brine then dried and concentrated. Purification of the crude material by flash chromatography (4:1 hexanes: ethyl acetate; 1:1 hexanes: ethyl acetate) yielded the desired hydantoin (0.530 g, 35 %).  $C_{20}H_{20}F_2N_2O_3$  (MW = 374.14); mass spectroscopy (MH+) = 375.2

Step B




The hydantoin from **Step A** (0.530 g, 0.00141 mol) was dissolved in DMF (5 ml) and treated with NaH (0.062 g, 0.00156 mol) followed by 1-iodo-propane (0.152 ml, 0.00156 mol). The reaction was stirred overnight under nitrogen. The reaction mixture was poured into 1N HCl and combined with ethyl acetate. The organic layer was extracted and washed with brine then dried and concentrated. Purification of the crude material by flash chromatography (2:1 hexanes: ethyl acetate) yielded the desired product (0.505 g, 86 %).  $C_{23}H_{26}F_2N_2O_3$  (MW = 416.19); mass spectroscopy (MH+) = 417.2



Step C

Lithium aluminum hydride (0.0609 g, 0.0018 mol) was dissolved in THF (5 ml). A THF solution of the hydantoin  
5 (0.505 g, 0.0012 mol) from **Step B** was added. The reaction was stirred overnight at room temperature. The reaction was quenched by the addition of 5N HCl. After stirring for thirty minutes, water was added and the solution was extracted with ethyl acetate. The organic layer was  
10 extracted and washed with brine then dried and concentrated. The crude product was carried forth without further purification (0.483 g, 100 %).  $C_{23}H_{26}F_2N_2O_2$  (MW = 400.20); mass spectroscopy (MH<sup>+</sup>) = 401


15

Step D

The methoxy ether from **Step C** (0.483 g, 0.0012 mol) was dissolved in methylene chloride (5 ml) and cooled to 0°C. To this solution was added, dropwise, a solution of BBr<sub>3</sub>  
20 (0.228 ml, 0.0024 mol) in methylene chloride (5 ml). After


stirring for about thirty minutes, the reaction mixture was cooled to 0°C and quenched by the dropwise addition of methanol/methylene chloride. The solvent was concentrated and the resulting material was dissolved in methylene chloride. The organic layer was extracted with water followed by brine. Upon evaporation of the solvent, the phenol (0.438 g, 95%) was obtained and was carried forth without further purification.  $C_{22}H_{24}F_2N_2O_2$  (MW = 386.18); mass spectroscopy (MH<sup>+</sup>) = 387.2

10



Step E

The phenol from **Step D** (0.438 g, 0.00110 mol) was dissolved in EtOH (7 ml) and treated with ethyl 2-bromoisobutyrate (0.500 ml, 0.0034 mol), powdered  $K_2CO_3$  (0.607 g, 0.0044 mol), and  $MgSO_4$  (0.132 g, 0.00110 mol). The reaction was stirred overnight at 77.7°C. Upon cooling, the reaction mixture was poured into 5N HCl and combined with EtOAc. The organic layer was extracted with water followed by brine then concentrated to dryness. Purification by flashed chromatography (1:1 hexanes: ethyl acetate) gave the ester (0.267 g, 49 %).  $C_{28}H_{34}F_2N_2O_4$  (MW = 500.59); mass spectroscopy (MH<sup>+</sup>) = 501.3

20

Step F

The ester from **Step E** (0.253 g, 0.00050 mol) was dissolved in methanol (4 ml) and treated with a solution of LiOH in water (1 ml). The reaction was stirred overnight. The reaction was cooled and water (20 ml) was added to the solution. The solution was then acidified using 1N HCl to pH = 3 then extracted with ethyl acetate. The organic layer was concentrated to afford the desired carboxylic acid (0.112 g, 47 %).  $C_{26}H_{30}F_2N_2O_4$  (MW = 472.22); mass spectroscopy ( $MH^+$ ) = 473.2


**Example 120**Step A

Following the procedure of **Example 15, Step A**, the hydantoin from **Example 13** (1.0g, 4.0mmol), 3,4-dichlorobenzylbromide (0.8ml, 4.4mmol), potassium carbonate (2.2g, 16.0mmol) and magnesium sulfate (0.6g, 5.0mmol) were used to yield the

desired product (1.03g).

$C_{20}H_{20}N_2O_3Cl_2$  (MW = 407.3);  $^1H$  NMR

Step B




5

Following the procedure of **Example 104, Step B**, the product from **Example 120, Step A** (1.0g, 2.46mmol), sodium hydride (0.11g, 2.7 mmol) and iodopropane (0.26ml, 2.71mmol) were used to yield the desired product (0.91g).

10  $C_{23}H_{26}N_2O_3Cl_2$  (MW = 449.4); MS ( $M^+$ , 449.1, 451.1)

Step C




15 Following the procedure of **Example 103, Step C**, the product from **Example 120, Step B** (0.9g, 2.0mmol) and lithium aluminum hydride (0.11g, 3.0mmol) were used to yield the desired imidazolone as a crude product (0.85g).

$C_{23}H_{26}N_2O_2Cl_2$  (MW = 433.4); MS ( $M^+$ , 433.1, 435.1)

20


Step D



Following the procedure of **Example 104, Step D**, the product from **Example 120, Step C** (0.85g, 2.0mmol) and boron tribromide (0.6ml, 6.0mmol) were used to yield the crude product.

5  $C_{22}H_{24}N_2O_2Cl_2$  (MW = 419.4); MS ( $M^+$ , 419.1, 421.1)

Step E




10 Following the procedure of **Example 103, Step E**, the product from **Example 120, Step D** (0.8g, 1.9mmol), ethyl 2-bromoisobutyrate (0.83ml, 5.7 mmol), potassium carbonate (1.0g, 7.6 mmol) and magnesium sulfate (0.3g, 2.5 mmol) were used to yield the product (0.48g).

$C_{28}H_{34}N_2O_4Cl_2$  (MW = 533.5); MS ( $M^+$ , 533.2, 535.2)

15



Step F




20 Following the procedure of **Example 103, Step F**, the product from **Example 120, Step E** (0.48g, 0.9mmol), lithium hydroxide (0.04g, 1.8mmol), methanol (8ml) and water (2ml) were used to yield the desired product after evaporation of the solvent and placement under vacuum (0.45g).

$C_{26}H_{30}N_2O_4Cl_2$  (MW = 505.5); MS ( $M^+$ , 505.2, 507.2)


25

**Example 121**Step A

- 5 Following the procedure of **Example 15, Step A**, the hydantoin from **Example 13** (1.0g, 4.0mmol), 4-fluorobenzylbromide (0.52ml, 4.4mmol), potassium carbonate (2.2g, 16.0mmol) and magnesium sulfate (0.6g, 5.0mmol) were used to yield the desired product (1.0g).
- 10  $C_{20}H_{21}N_2O_3F$  (MW = 356.4); MS ( $M^+$ , 357.2)


Step B

- 15 Following the procedure of **Example 104, Step B**, the product from **Example 121, Step A** (1.0g, 2.8mmol), sodium hydride (0.12g, 3.1 mmol) and iodopropane (0.3ml, 3.1mmol) were used to yield the desired product (0.85g).
- $C_{23}H_{27}N_2O_3F$  (MW = 398.5); MS ( $M^+$ , 399.2)

Step C

Following the procedure of **Example 103, Step C**, the product from **Example 121, Step B** (0.85g, 2.1mmol) and lithium aluminum hydride (0.12g, 3.2mmol) were used to yield the  
5 desired imidazolone as a crude product (0.64g).


$C_{23}H_{27}N_2O_2F$  (MW = 382.5); MS ( $M^+$ , 383.2)

Step D

10

Following the procedure of **Example 104, Step D**, the product from **Example 121, Step C** (0.64g, 1.7mmol) and boron tribromide (0.49ml, 5.1mmol) were used to yield the crude product.

15  $C_{22}H_{25}N_2O_2F$  (MW = 368.5); MS ( $M^+$ , 369.2)


Step E

Following the procedure of **Example 103, Step E**, the product from **Example 121, Step D** (0.62g, 1.7mmol), ethyl 2-bromoisobutyrate (0.74ml, 5.1 mmol), potassium carbonate  
20

(0.9g, 6.8 mmol) and magnesium sulfate (0.3g, 2.5 mmol) were used to yield the product (0.49g).

$C_{28}H_{35}N_2O_4F$  (MW = 482.6); MS ( $M^+$ , 483.3)

5



Step F

Following the procedure of **Example 103, Step F**, the product from **Example 121, Step E** (0.49g, 1.0mmol), sodium hydroxide (2N, 2ml) and methanol (8ml) were used to yield the desired product after evaporation of the solvent and placement under vacuum (0.4g).

$C_{26}H_{31}N_2O_4F$  (MW = 454.6); MS ( $M^+$ , 455.3,  $M^-$ , 453.1)

**Example 122**

15


Step A

20 Following the procedure of **Example 15, Step A**, the hydantoin




- from **Example 13** (1.0g, 4.0mmol), 3,5-bistrifluoromethylbenzyl bromide (0.78ml, 4.4mmol), potassium carbonate (2.2g, 16.0mmol) and magnesium sulfate (0.6g, 5.0mmol) were used to yield the desired product (0.85g).
- $C_{22}H_{20}N_2O_3F_6$  (MW = 474.4); MS ( $M^+$ , 475.1)

Step B




- Following the procedure of **Example 104, Step B**, the product from **Example 122, Step A** (1.5g, 3.2mmol), sodium hydride (0.13g, 3.5 mmol) and iodopropane (0.34ml, 3.5mmol) were used to yield the desired product (1.0g).
- $C_{25}H_{26}N_2O_3F_6$  (MW = 516.5); MS ( $M^+$ , 517.2)

Step C




- Following the procedure of **Example 103, Step C**, the product from **Example 122, Step B** (1.0g, 1.9mmol) and lithium aluminum hydride (0.11g, 2.9mmol) were used to yield the desired imidazolone as a crude product (0.84g).
- $C_{25}H_{26}N_2O_2F_6$  (MW = 500.5); MS ( $M^+$ , 501.2)

Step D


Following the procedure of **Example 104, Step D**, the product from **Example 122, Step C** (0.84g, 1.7mmol) and boron tribromide (0.49ml, 5.1mmol) were used to yield the crude product.

$C_{24}H_{24}N_2O_2F_6$  (MW = 486.5); MS ( $M^+$ , 487.2)

Step E

Following the procedure of **Example 103, Step E**, the product from **Example 122, Step D** (0.62g, 1.7mmol), ethyl 2-bromoisobutyrate (0.74ml, 5.1 mmol), potassium carbonate (0.9g, 6.8 mmol) and magnesium sulfate (0.3g, 2.5 mmol) were used to yield the product (0.57g).


$C_{30}H_{34}N_2O_4F_6$  (MW = 600.6); MS ( $M^+$ , 601.3)

Step F


Following the procedure of **Example 103, Step F**, the product from **Example 122, Step E** (0.57g, 0.9mmol), sodium hydroxide (2N, 2ml) and methanol (8ml) were used to yield the desired product after evaporation of the solvent and placement under vacuum (0.54g).

$C_{28}H_{30}N_2O_4F_6$  (MW = 572.6); MS ( $M^+$ , 573.3,  $M^-$ , 571.1)

### Example 123



#### Step A




A DMF solution (50 mL) of the hydantoin from **Example 13** (1.22 g, 4.2 mmol) was treated sequentially with phenethyl bromide (574  $\mu$ L, 4.2 mmol),  $K_2CO_3$  (1.3 g, 9.4 mmol), and  $MgSO_4$  (1.8 g, 15 mmol). The resulting mixture was stirred at room temperature overnight. The reaction was quenched with 1 N HCl (200 mL) extracted with diethyl ether (1x200 mL). The organic extract was washed with brine then dried over  $Na_2SO_4$  and concentrated. Purification by

chromatography (gradient: 2:1 to 1:1 hexanes:ethyl acetate) gave the desired alkylated hydantoin as a white solid (766.6 mg, 52%).

$C_{21}H_{24}N_2O_3$  (MW = 352.44); mass spectroscopy:  $(MH^+) = 353.2$ ,


5  $(MH^-) = 351.3$

Step B




The hydantoin from **Step A** (761.5 mg, 2.16 mmol) was dissolved in DMF (15 mL) and cooled to 0 °C under an atmosphere of  $N_2$ . The mixture was treated with NaH (60% dispersion in oil, 101 mg, 2.52 mmol) and, after 15 minutes, n-propyl iodide (243  $\mu$ L, 2.49 mmol). The reaction mixture was warmed to room temperature, stirred overnight, then poured into 1 N HCl (100 mL). The resulting solution was extracted with diethyl ether and the organic extract was dried over  $Na_2SO_4$  and concentrated. Purification by chromatography (gradient: 5:1 to 3:1 hexanes:ethyl acetate) gave the desired product as a colorless oil (720.8 mg, 85%).

15  $C_{24}H_{30}N_2O_3$  (MW = 394.52); mass spectroscopy:  $(MH^+) = 395.3$

Step C

The hydantoin from **Step B** (704.9 mg, 1.79 mmol) was dissolved in THF (20 mL) and cooled to 0 °C. LAH (68 mg, 1.79 mmol) was added in one portion and, after 45 minutes, the reaction was quenched by the addition of 5 N HCl (5 mL). The resulting mixture was stirred at room temperature for 30 minutes then diluted with H<sub>2</sub>O (100 mL) and extracted with ethyl acetate (2x50 mL). The combined organic extracts were washed with brine, dried over Na<sub>2</sub>SO<sub>4</sub> and concentrated to give the desired imidazolone (667.6 mg, 99%). The crude product was carried forward without further purification. C<sub>24</sub>H<sub>30</sub>N<sub>2</sub>O<sub>2</sub> (MW = 378.52); mass spectroscopy: (MH<sup>+</sup>) = 379.2


Step D

The imidazolone from **Step C** (659.4 mg, 1.74 mmol) was dissolved in CH<sub>2</sub>Cl<sub>2</sub> (20 mL) and cooled to 0 °C. BBr<sub>3</sub> (823 μL, 8.70 mmol) was added dropwise, then the reaction mixture was warmed to room temperature. After 1 hour, the solution

was again cooled to 0 °C and quenched by the slow addition of a methanol:CH<sub>2</sub>Cl<sub>2</sub> (1:4) solution. The resulting mixture was extracted with H<sub>2</sub>O (50 mL). The aqueous extract was washed with CH<sub>2</sub>Cl<sub>2</sub> (50 mL) then the combined organic  
5 extracts were washed with brine, dried over Na<sub>2</sub>SO<sub>4</sub>, and concentrated to give the desired phenol as a foam. The crude product was carried forward without further purification.


C<sub>23</sub>H<sub>28</sub>N<sub>2</sub>O<sub>2</sub> (MW = 364.49); mass spectroscopy: (MH<sup>+</sup>) = 365.3,  
10 (MH<sup>+</sup>) = 363.3

Step E





A DMF solution (20 mL) of the phenol from **Step D** was treated sequentially with ethyl 2-bromoisobutyrate (725 µL, 5.35  
15 mmol), K<sub>2</sub>CO<sub>3</sub> (1.2 g, 8.7 mmol), and MgSO<sub>4</sub> (1.2 g, 10 mmol). The resulting mixture was heated to 55-65 °C for two days. The reaction was quenched with 1 N HCl (100 mL) then extracted with diethyl ether (2x70 mL). The combined organic extracts were washed with brine then dried over  
20 Na<sub>2</sub>SO<sub>4</sub> and concentrated. Purification by chromatography (gradient: 2:1 to 1:1 hexanes:ethyl acetate) gave the desired ester as a slightly yellow oil (325.6 mg, 39% for two steps).

C<sub>29</sub>H<sub>38</sub>N<sub>2</sub>O<sub>4</sub> (MW = 478.64); mass spectroscopy: (MH<sup>+</sup>) = 479.4


Step F

A dioxane solution (10 mL) of the ester from **Step E** (317.6 mg, 0.66 mmol) was treated with an aqueous solution (5 mL) of LiOH (53 mg, 2.2 mmol). The mixture was stirred at room temperature for two hours. The solvent was concentrated and the resulting oil diluted with H<sub>2</sub>O (50 mL) and 1 N NaOH (10 mL), then extracted with Et<sub>2</sub>O (50 mL). The aqueous extract was acidified with 1 N HCl and extracted with diethyl ether (2x50 mL). The combined organic extracts were washed with brine, dried over Na<sub>2</sub>SO<sub>4</sub>, and concentrated to give the desired carboxylic acid as a foam-like solid (253.6 mg, 85%) C<sub>27</sub>H<sub>34</sub>N<sub>2</sub>O<sub>4</sub> (MW = 450.58); mass spectroscopy: (MH<sup>+</sup>) = 451.3, (MH<sup>-</sup>) = 449.1

**Example 124**

Step A

A DMF solution (50 mL) of the hydantoin from **Example 13** (1.02 g, 4.1 mmol) was treated with a DMF solution (10 mL)  
5 4-isopropylbenzyl bromide (0.693 g, 4.1 mmol) followed by  $K_2CO_3$  (1.2 g, 8.7 mmol), and  $MgSO_4$  (1.2 g, 10 mmol). The resulting mixture was stirred at room temperature overnight. The reaction was poured into 1 N HCl (150 mL) then extracted with ethyl acetate (2x100 mL). The combined organic  
10 extracts were washed with brine then dried over  $Na_2SO_4$  and concentrated. Purification by chromatography (gradient: 3:1 to 1:1 hexanes:ethyl acetate) gave the desired alkylated hydantoin as a white solid (842.5 mg, 54%).  
 $C_{23}H_{28}N_2O_3$  (MW = 380.49); mass spectroscopy:  $(MH^+) = 381.3$ ,  
15  $(MH^-) = 379.1$

Step B


The hydantoin from **Step A** (827.3 mg, 2.17 mmol) was dissolved in DMF (15 mL) and cooled to 0 °C. The mixture  
20 was treated with NaH (60% dispersion in oil, 106 mg, 2.65 mmol) and, after 15 minutes, n-propyl iodide (244  $\mu$ L, 2.50



mmol). The reaction mixture was warmed to room temperature, stirred overnight, then poured into 1 N HCl (100 mL). The resulting solution was extracted with diethyl ether and the organic extract was washed with brine, dried over Na<sub>2</sub>SO<sub>4</sub> and concentrated. Purification by chromatography (gradient: 5:1 to 3:1 hexanes:ethyl acetate) gave the desired product as a colorless oil (705.3 mg, 77%).


C<sub>26</sub>H<sub>34</sub>N<sub>2</sub>O<sub>3</sub> (MW = 422.57); mass spectroscopy: (MH<sup>+</sup>) = 423.3

10

Step C


The hydantoin from **Step B** (687.0 mg, 1.63 mmol) was dissolved in THF (20 mL) and cooled to 0 °C. LAH (63 mg, 1.66 mmol) was added in one portion and, after 45 minutes, the reaction was quenched by the addition of 5 N HCl (5 mL) in THF (10 mL). The resulting mixture was stirred at room temperature for 30 minutes then diluted with H<sub>2</sub>O (100 mL) and extracted with ethyl acetate (2x50 mL). The combined organic extracts were washed with brine, dried over Na<sub>2</sub>SO<sub>4</sub> and concentrated to give the desired imidazolone (640.1 mg, 97%). The crude product was carried forward without further purification.

C<sub>26</sub>H<sub>34</sub>N<sub>2</sub>O<sub>2</sub> (MW = 406.57); mass spectroscopy: (MH<sup>+</sup>) = 407.2

Step D

The imidazolone from **Step C** (630.8 mg, 1.55 mmol) was dissolved in  $\text{CH}_2\text{Cl}_2$  (20 mL) and cooled to 0 °C.  $\text{BBr}_3$  (733  
5  $\mu\text{L}$ , 7.75 mmol) was added dropwise, then the reaction mixture was warmed to room temperature. After 1 hour, the solution was again cooled to 0 °C and quenched by the slow addition of a methanol: $\text{CH}_2\text{Cl}_2$  (1:4) solution. The resulting mixture was extracted with  $\text{H}_2\text{O}$  (50 mL). The aqueous extract was  
10 washed with  $\text{CH}_2\text{Cl}_2$  (50 mL) then the combined organic extracts were washed with brine, dried over  $\text{Na}_2\text{SO}_4$ , and concentrated to give the desired phenol as a foam. The crude product was carried forward without further purification.

15  $\text{C}_{25}\text{H}_{32}\text{N}_2\text{O}_2$  (MW = 392.55); mass spectroscopy:  $(\text{MH}^+) = 393.3$ ,  $(\text{MH}^-) = 391.4$


Step E

A DMF solution (20 mL) of the phenol from **Step D** was treated sequentially with ethyl 2-bromoisobutyrate (650  $\mu$ L, 4.8 mmol),  $K_2CO_3$  (1.1 g, 8.0 mmol), and  $MgSO_4$  (1.1 g, 9.2 mmol). The resulting mixture was heated to 55-65  $^{\circ}C$  for two days.

5 The reaction was quenched with 1 N HCl (100 mL) then extracted with diethyl ether (2x75 mL). The combined organic extracts were washed with brine then dried over  $Na_2SO_4$  and concentrated. Purification by chromatography (gradient: 2:1 to 1:1 hexanes:ethyl acetate) gave the  
10 desired ester as a slightly yellow oil (202.1 mg, 26% for two steps).

$C_{27}H_{42}N_2O_4$  (MW = 506.69); mass spectroscopy:  $(MH^+) = 507.4$


Step F




15

A dioxane solution (10 mL) of the ester from **Step E** (194.5 mg, 0.43 mmol) was treated with an aqueous solution (5 mL) of LiOH (45 mg, 1.9 mmol). The mixture was stirred at room temperature for two hours. The solvent was concentrated and  
20 the resulting oil diluted with  $H_2O$  (50 mL) and 1 N NaOH (10 mL), then extracted with  $Et_2O$  (50 mL). The aqueous extract was acidified with 1 N HCl and extracted with diethyl ether (2x50 mL). The combined organic extracts were washed with brine, dried over  $Na_2SO_4$ , and concentrated to give the  
25 desired carboxylic acid as a foam-like solid (163.0 mg, 79%)

$C_{29}H_{38}N_2O_4$  (MW = 478.64); mass spectroscopy:  $(MH^+)$  = 479.3,  
 $(MH^-)$  = 477.1

**Example 125**


5

Step A

Following the procedure of **Example 15, Step A**, the hydantoin  
from **Example 13** (1.0g, 4.0mmol), 4-ethylbenzyl chloride  
10 (0.7ml, 4.4mmol), potassium carbonate (2.2g, 16.0mmol) and  
magnesium sulfate (0.6g, 5.0mmol) were used to yield the  
desired product (0.9g).


$C_{22}H_{26}N_2O_3$  (MW = 366.5); MS  $(M^-)$ , 365.1

15

Step B


Following the procedure of **Example 104, Step B**, the product  
from **Example 125, Step A** (0.9g, 2.5mmol), sodium hydride  
(0.12g, 2.8 mmol) and iodopropane (0.3ml, 2.8mmol) were used  
20 to yield the desired product (0.77g).

$C_{25}H_{32}N_2O_3$  (MW = 408.6); MS  $(M^+)$ , 409.3

Step C


Following the procedure of **Example 103, Step C**, the product from **Example 125, Step B** (0.76g, 1.9mmol) and lithium aluminum hydride (0.11g, 2.9mmol) were used to yield the desired imidazolone as a crude product (0.72g).

$C_{25}H_{32}N_2O_2$  (MW = 392.6); MS ( $M^+$ , 393.3)

Step D

Following the procedure of **Example 104, Step D**, the product from **Example 125, Step C** (0.72g, 1.8mmol) and boron tribromide (Aldrich, 0.52ml, 5.4mmol) were used to yield the crude product.

$C_{24}H_{30}N_2O_2$  (MW = 378.5); MS ( $M^+$ , 379.2)


Step E

Following the procedure of **Example 103, Step E**, the product from **Example 125, Step D** (0.68g, 1.8mmol), ethyl 2-bromoisobutyrate (0.83ml, 5.4 mmol), potassium carbonate

(1.0g, 7.2 mmol) and magnesium sulfate (0.36g, 3.0 mmol) were used to yield the product (0.29g).

$C_{30}H_{40}N_2O_4$  (MW = 492.7); MS ( $M^+$ , 493.3)


5

Step F


Following the procedure of **Example 103, Step F**, the product from **Example 125, Step E** (0.49g, 1.0mmol), sodium hydroxide (2N, 1ml) and methanol (4ml) were used to yield the desired product after evaporation of the solvent and placement under vacuum.

10

$C_{28}H_{36}N_2O_4$  (MW = 464.6); MS ( $M^+$ , 465.2,  $M^-$ , 463.4)

**Example 126**

15


Step A

The product from **Example 16, Step A** (0.6g, 1.2mmol) and 4-picolyl chloride hydrochloride (0.2ml, 1.3 mmol) and triethylamine (0.2ml) stirred together in DMF (20ml). Potassium carbonate (0.7g, 4.8 mmol) and magnesium sulfate

20


(0.2g, 1.5mmol) added and the mixture stirred at ambient temperature overnight. The reaction mixture was carefully added to 1 N hydrochloric acid (50ml) and the resulting solution was extracted twice with ethyl acetate. The organic layers were combined, washed with aqueous brine then dried over sodium sulfate. Evaporation of the solvent and subsequent purification by flash chromatography (hexanes:ethyl acetate) gave the desired product (0.42g).  
 $C_{35}H_{43}N_3O_5$  (MW = 585.8); MS ( $M^+$ , 586.3)

10

Step B


Following the procedure of **Example 25, Step B**, the product from **Example 126, Step A** (0.4g, 0.7mmol) and trifluoroacetic acid (2ml) were used to yield the desired product.  
 $C_{31}H_{35}N_3O_5$  (MW = 529.6); MS ( $M^+$ , 530.4)

15


Step C

Following the procedure of **Example 15, Step B**, product from **Example 126, Step B** (0.35g, 0.7mmol) and sodium borohydride (0.28g, 7.5mmol) were used to yield the desired product after flash chromatography (methylene chloride:methanol).  
 $C_{31}H_{35}N_3O_4$  (MW = 513.6); MS ( $M^+$ , 514.4,  $M^-$ , 512.2)

20


**Example 127**

5

Step A

Following the procedure of **Example 15, Step A**, the hydantoin from **Example 13** (1.0g, 4.0mmol), iodopropane (0.43ml, 4.4mmol), potassium carbonate (2.2g, 16.0mmol) and magnesium sulfate (0.6g, 5.0mmol) were used to yield the desired product (0.95g).

$C_{16}H_{22}N_2O_3$  (MW = 290.4); MS ( $M^+$ , 291.2)


Step B

15

Following the procedure of **Example 104, Step B**, the product from **Example 127, Step A** (0.9g, 3.1mmol), sodium hydride (0.13g, 3.4 mmol) and 3,4-dimethylbenzyl chloride (0.5ml, 3.4mmol) were used to yield the desired product (1.1g).


20  $C_{25}H_{32}N_2O_3$  (MW = 408.6); MS ( $M^+$ , 409.3)



Step C

Following the procedure of **Example 103, Step C**, the product from **Example 127, Step B** (0.8g, 1.9mmol) and lithium aluminum hydride (0.16g, 4.1mmol) were used to yield the  
5 desired imidazolone as a crude product (0.8g).


$C_{25}H_{32}N_2O_2$  (MW = 392.6); MS ( $M^+$ , 393.3)

Step D

10


Following the procedure of **Example 104, Step D**, the product from **Example 127, Step C** (0.8g, 2.1mmol) and boron tribromide (0.6ml, 6.3mmol) were used to yield the crude product (0.7g).

15  $C_{24}H_{30}N_2O_2$  (MW = 378.5); MS ( $M^+$ , 379.2)

Step E


Following the procedure of **Example 103, Step E**, the product from **Example 127, Step D** (0.7g, 1.8mmol), ethyl 2-bromoisobutyrate (0.8ml, 5.4 mmol), potassium carbonate (1.0g, 7.2 mmol) and magnesium sulfate (0.4g, 3.0 mmol) were used to yield the product (0.42g).  
C<sub>30</sub>H<sub>40</sub>N<sub>2</sub>O<sub>4</sub> (MW = 492.7); MS (M<sup>+</sup>, 493.3)

Step F




- 10 Following the procedure of **Example 103, Step F**, the product from **Example 127, Step E** (0.4g, 1.0mmol), sodium hydroxide (2N, 1.5ml) and methanol (5ml) were used to yield the desired product after evaporation of the solvent and placement under vacuum.
- 15 C<sub>28</sub>H<sub>36</sub>N<sub>2</sub>O<sub>4</sub> (MW = 464.6); MS (M<sup>+</sup>, 465.3, M<sup>-</sup>, 463.1)

**Example 128**





20. A CH<sub>2</sub>Cl<sub>2</sub> solution (15 mL) of the ester from **Example 16, Step A** (279.9 mg, 0.57 mmol) was cooled to 0 °C and treated with

- TFA (1 mL, 13 mmol). The mixture was warmed to room temperature and stirred for 1.5 hours. The solvent was concentrated to give the crude acid, which was dissolved in 1 N NaOH (25 mL) and washed with diethyl ether (25 mL). The aqueous solution was then acidified with 5 N HCl and extracted with diethyl ether (1x25 mL) and ethyl acetate (1x25 mL). The combined organic extracts were dried over Na<sub>2</sub>SO<sub>4</sub> and concentrated to give the desired caboxylic acid as a white foam (257.3 mg).
- 10 C<sub>25</sub>H<sub>30</sub>N<sub>2</sub>O<sub>5</sub> (MW = 438.53); mass spectroscopy: (MH<sup>+</sup>) = 439.1, (MH<sup>-</sup>) = 437.3


**Example 129**

- 15 see **Example 22, Step B**

**Example 130**

Step A


The hydantoin from **Example 16, Step A** (234.0 mg, 0.47 mmol) was dissolved in DMF (20 mL) and cooled to 0 °C under an atmosphere of N<sub>2</sub>. The mixture was treated with NaH (60% dispersion in oil, 25 mg, 0.63 mmol) and, after 10 minutes, n-propyl iodide (55 µL, 0.56 mmol). The reaction mixture was warmed to room temperature, stirred overnight, then poured into 1 N HCl (100 mL). The resulting solution was extracted with diethyl ether (2x70 mL). The combined organic extracts were dried over Na<sub>2</sub>SO<sub>4</sub> and concentrated. Purification by chromatography (5:1 hexanes:ethyl acetate) gave the desired product as a colorless oil (70.8 mg, 28%). C<sub>32</sub>H<sub>44</sub>N<sub>2</sub>O<sub>5</sub> (MW = 536.72); mass spectroscopy: (MH<sup>+</sup>-t-butyl) = 481.4, (M+NH<sub>4</sub><sup>+</sup>) = 554.5

Step B


A  $\text{CH}_2\text{Cl}_2$  solution (10 mL) of the ester from **Step A** (65 mg, 0.12 mmol) was cooled to 0 °C and treated with TFA (0.5 mL, 6.5 mmol). The mixture was warmed to room temperature and stirred for 2 hours. The solvent was concentrated to give the crude acid, which was dissolved in 1 N NaOH (50 mL) and washed with diethyl ether (50 mL). The aqueous solution was then acidified with 5 N HCl and extracted with ethyl acetate (2x25 mL). The combined organic extracts were dried over  $\text{Na}_2\text{SO}_4$  and concentrated to give the desired caboxylic acid as a colorless oil (55.1 mg, 95%).

$\text{C}_{28}\text{H}_{36}\text{N}_2\text{O}_5$  (MW = 480.61); mass spectroscopy:  $(\text{MH}^+) = 481.2$ ,  $(\text{MH}^-) = 479.3$

#### Example 131



#### Step A




The hydantoin from **Example 22, Step A** (1.06 g, 2.05 mmol) was dissolved in DMF (25 mL) and cooled to 0 °C under an atmosphere of  $\text{N}_2$ . The mixture was treated with NaH (60%

dispersion in oil, 101 mg, 2.5 mmol) and, after 5 minutes, a slight excess of 4-methoxybenzyl chloride. The reaction mixture was warmed to room temperature, stirred for 2 hours, then poured into 1 N HCl (100 mL). The resulting solution was extracted with diethyl ether (2x100 mL). The combined organic extracts were washed with brine, dried over Na<sub>2</sub>SO<sub>4</sub> and concentrated. Purification by chromatography (2:1 hexanes:ethyl acetate) gave the desired product as a colorless oil (70.8 mg, 28%).


10 C<sub>39</sub>H<sub>44</sub>N<sub>2</sub>O<sub>6</sub> (MW = 636.80); mass spectroscopy: (M+NH<sub>4</sub><sup>+</sup>) = 654.4

Step B




A CH<sub>2</sub>Cl<sub>2</sub> solution (10 mL) of the ester from **Step A** (55 mg, 0.086 mmol) was cooled to 0 °C and treated with TFA (0.5 mL, 6.5 mmol). The mixture was warmed to room temperature and stirred for 2 hours. The solvent was concentrated to give the crude acid, which was dissolved in 1 N NaOH (50 mL) and washed with diethyl ether (50 mL). The aqueous solution was then acidified with 5 N HCl and extracted with ethyl acetate (50 mL). The organic extract was dried over Na<sub>2</sub>SO<sub>4</sub> and concentrated to give the desired caboxylic acid (55.4 mg). C<sub>35</sub>H<sub>36</sub>N<sub>2</sub>O<sub>6</sub> (MW = 580.69); mass spectroscopy: (MH<sup>+</sup>) = 581.2, (MH<sup>-</sup>) = 579.4

## Example 132




## Step A



- 5 The 4-(4-methoxyphenyl) butyric acid (10.04 g, 0.052 mol) and hydroxylamine (5.04 g, 0.052 mol) were combined in THF and treated with EDAC (11.86 g, 0.062 mol), HOBt (7.0 g, 0.052 mol) and diisopropyl ethyl amine (16.9 ml, 0.124 mol). The reaction was stirred overnight. The reaction was
- 10 concentrated to one-half its original volume. Ethyl acetate (100 ml) was added and the solution was extracted with 1N HCl. The organic layer was washed with 2N NaOH then concentrated. Purification of the crude material by flash chromatography (2: 1 hexanes: ethyl acetate) gave the
- 15 Weinreb amide (9.77 g, 78 %).  $C_{13}H_{19}NO_3$  (MW = 237.14); mass spectroscopy (MH<sup>+</sup>) = 238.0

## Step B




- 20 The Weinreb amide from **Step A** (4.7 g, 0.0198 mol) was dissolved in ether (164 ml) and cooled to 0°C. Methyl magnesium bromide (3.0 M in ether, 33 ml, 99 mmol) was added slowly to the reaction mixture. The reaction was stirred

overnight. Hydrochloric acid (1N, 50 ml) was and the mixture was extracted with ether. The organic layer was stripped to afford the methyl ketone as a clear oil (3.66 g, 96%).

$C_{12}H_{16}O_2$  (MW = 192.12); mass spectroscopy (MH+) = 192.1

5


Step C



The methyl ketone from **Step B** (3.66 g, 0.0190 mol) was combined with potassium cyanide (2.73 g, 0.042 mol) and ammonium carbonate (9.13 g, 0.095 mol) in 1:1 methanol: water (35:35 ml) and stirred overnight at 50°C. The reaction mixture was poured into water and extracted with ethyl acetate. The organic layer was washed with brine then concentrated. Purification of the crude material by flash chromatography (2:1 hexanes: ethyl acetate) yielded the desired hydantoin as a thick, clear oil.  $C_{14}H_{18}N_2O_3$  (MW = 262.13); mass spectroscopy (MH+) = 263.1

15

Step D



20

The hydantoin from **Step C** (1 g, 0.0038 mol) was dissolved in DMF (20 ml) and treated with  $K_2CO_3$  (2.09 g, 0.0152 mol) and magnesium sulfate (0.689 g, 0.0057 mol) followed by 3, 5 difluoro-bromobenzene (0.543 ml, 0.0042 mol). The reaction was stirred overnight. Hydrochloric acid was added and the mixture was extracted with ethyl acetate. The organic layer

25




was concentrated. Purification of the crude material by flash chromatography (4: 1 ethyl acetate: hexanes) give the alkylated hydantoin (1.12 g, 76%).

$C_{21}H_{22}F_2N_2O_3$  (MW = 388.16); mass spectroscopy (MH+) = 389.2

5


#### Step E



Sodium hydride (0.121 g, 0.0030 mol) was suspended in DMF (5 ml) and cooled to 0°C. The hydantoin from **Step D** (1.07 g, 0.00276 mol) was added as a solution in DMF (5 ml).  
Iodopropane (0.295 ml, 0.0030 mol) was syringed into solution and the reaction was stirred overnight at room temperature. The reaction mixture was poured into 1N HCl and extracted with ethyl acetate. Purification of the crude material by flash chromatography (4:1 hexanes: ethyl acetate) gave the desired product as a clear oil (0.951 g, 80%).  $C_{24}H_{28}F_2N_2O_3$  (MW = 430.21); mass spectroscopy (MH+) = 431.2


20

#### Step F



The methoxy ether from **Step E** (0.951 g, 0.0022 mol) was dissolved in methylene chloride (5 ml) and cooled to 0°C. To this solution was added, dropwise, a solution of BBr<sub>3</sub> (0.418 ml, 0.0044 mol) in methylene chloride (5 ml). After stirring for about twenty minutes, the reaction mixture was cooled to 0°C and quenched by the dropwise addition of methanol/methylene chloride. The solvent was concentrated and the resulting material was dissolved in methylene chloride. The organic layer was extracted with water followed by brine. Upon evaporation of the solvent, the phenol (0.910 g, 99%) was obtained and was carried forth without further purification. C<sub>23</sub>H<sub>26</sub>F<sub>2</sub>N<sub>2</sub>O<sub>3</sub> (MW = 416.19); mass spectroscopy (MH<sup>+</sup>) = 417.2


15

Step G

The phenol from **Step F** (0.910 g, 0.00220 mol) was dissolved in EtOH and treated with ethyl 2-bromoisobutyrate (0.963 ml, 0.0066 mol), powdered K<sub>2</sub>CO<sub>3</sub> (1.210 g, 0.0088 mol), and MgSO<sub>4</sub> (0.264 g, 0.0022 mol). The reaction was stirred overnight at 77.7°C. Upon cooling, the reaction mixture was poured into 5N HCl and combined with EtOAc. The organic layer was extracted with water followed by brine then concentrated to dryness. Purification by flashed chromatography (5:1 hexanes: ethyl acetate) gave the ester (0.850 g, 69 %). C<sub>29</sub>H<sub>36</sub>F<sub>2</sub>N<sub>2</sub>O<sub>5</sub> (MW = 530.26); mass spectroscopy


(MH<sup>+</sup>) = 531.3


Step H



- 5 The ester from **Step G** (0.500 g, 0.00094 mol) was dissolved in methanol (8 ml) and treated with a solution of LiOH in water (2 ml). The reaction was stirred overnight. The reaction was cooled and water was added to the solution. The solution was then acidified using 1N HCl to pH = 3 then
- 10 extracted with ethyl acetate. The organic layer was concentrated to afford the desired carboxylic acid (0.112 g, 47 %). Purification of the crude material using reverse phase chromatography (7:3 acetonitrile: water) afforded the desired acid (0.033 mg, 7%) C<sub>27</sub>H<sub>32</sub>F<sub>2</sub>N<sub>2</sub>O<sub>5</sub> (MW = 502.23); mass
- 15 spectroscopy (MH<sup>+</sup>) = 503.3


**Example 133**



Step A


The 4-(4-methoxyphenyl) butyric acid (10.04 g, 0.052 mol) and hydroxylamine (5.04 g, 0.052 mol) were combined in THF and treated with EDAC (11.86 g, 0.062 mol), HOBT (7.0 g, 0.052 mol) and diisopropyl ethyl amine (16.9 ml, 0.124 mol). The reaction was stirred overnight. The reaction was concentrated to one-half its original volume. Ethyl acetate (100 ml) was added and the solution was extracted with 1N HCl. The organic layer was washed with 2N NaOH then concentrated. Purification of the crude material by flash chromatography (2: 1 hexanes: ethyl acetate) gave the Weinreb amide (9.77 g, 78 %).  $C_{13}H_{19}NO_3$  (MW = 237.14); mass spectroscopy (MH+) = 238.0

15


Step B

The Weinreb amide from **Step A** (4.7 g, 0.0198 mol) was dissolved in ether (164 ml) and cooled to 0°C. Methyl magnesium bromide (3.0 M in ether, 33 ml, 99 mmol) was added slowly to the reaction mixture. The reaction was stirred overnight. Hydrochloric acid (1N, 50 ml) was added and the mixture was extracted with ether. The organic layer was stripped to afford the methyl ketone as a clear oil (3.66 g, 96%).  $C_{12}H_{16}O_2$  (MW = 192.12); mass spectroscopy (MH+) = 192.1


25

Step C


The methyl ketone from **Step B** (3.66 g, 0.0190 mol) was combined with potassium cyanide (2.73 g, 0.042 mol) and ammonium carbonate (9.13 g, 0.095 mol) in 1:1 methanol: water (35:35 ml) and stirred overnight at 50°C. The reaction mixture was poured into water and extracted with ethyl acetate. The organic layer was washed with brine then concentrated. Purification of the crude material by flash chromatography (2:1 hexanes: ethyl acetate) yielded the desired hydantoin as a thick, clear oil.  $C_{14}H_{18}N_2O_3$  (MW = 262.13); mass spectroscopy (MH<sup>+</sup>) = 263.1

Step D

The hydantoin from **Step C** (1 g, 0.0038 mol) was dissolved in DMF (20 ml) and treated with  $K_2CO_3$  (2.09 g, 0.0152 mol) and magnesium sulfate (0.689 g, 0.0057 mol) followed by 3, 4 dimethyl-chlorobenzene (0.607 ml, 0.0042 mol). The reaction was stirred overnight. Hydrochloric acid was added and the mixture was extracted with ethyl acetate. The organic layer was concentrated. Purification of the crude material by flash chromatography (2: 1 ethyl acetate: hexanes) gave the alkylated hydantoin (1.30 g, 91%).  $C_{23}H_{28}N_2O_3$  (MW = 380.21); mass spectroscopy (MH<sup>+</sup>) = 381.2

Step E


Sodium hydride (0.150 g, 0.0038 mol) was suspended in DMF (5 ml) and cooled to 0°C. The hydantoin from **Step D** (1.30 g, 0.00340 mol) was added as a solution in DMF (5 ml).  
5 Iodopropane (0.367 ml, 0.0038 mol) was syringed into solution and the reaction was stirred overnight at room temperature. The reaction mixture was poured into 1N HCl and extracted with ethyl acetate. Purification of the crude  
10 material by flash chromatography (2:1 hexanes: ethyl acetate) gave the desired product as a clear oil (1.27 g, 89%).  $C_{26}H_{34}N_2O_3$  (MW = 422.26); mass spectroscopy (MH+) = 423.2

Step F

15 The methoxy ether from **Step E** (1.27 g, 0.0030 mol) was dissolved in methylene chloride (10 ml) and cooled to 0°C. To this solution was added, dropwise, a solution of  $BBr_3$  (0.569 ml, 0.0060 mol) in methylene chloride (5 ml). After  
20 stirring for about twenty minutes, the reaction mixture was

cooled to 0°C and quenched by the dropwise addition of methanol/methylene chloride. The solvent was concentrated and the resulting material was dissolved in methylene chloride. The organic layer was extracted with water followed by brine. Upon evaporation of the solvent, the phenol (1.19 g, 98%) was obtained and was carried forth without further purification.  $C_{25}H_{32}N_2O$  (MW = 408.24); mass spectroscopy (MH<sup>+</sup>) = 409.2


10

Step G

15



The phenol from **Step F** (1.19 g, 0.00290 mol) was dissolved in EtOH and treated with ethyl 2-bromoisobutyrate (1.28 ml, 0.0087 mol), powdered  $K_2CO_3$  (1.610 g, 0.0117 mol), and  $MgSO_4$  (0.348 g, 0.0029 mol). The reaction was stirred overnight at 77.7°C. Upon cooling, the reaction mixture was poured into 5N HCl and combined with EtOAc. The organic layer was extracted with water followed by brine then concentrated to dryness. Purification by flashed chromatography (5:1 hexanes: ethyl acetate) gave the ester (1.26 g, 83 %).  $C_{31}H_{42}N_2O_3$  (MW = 522.31); mass spectroscopy (MH<sup>+</sup>) =

20

Step H

The ester from **Step G** (0.500 g, 0.00095 mol) was dissolved in methanol (8 ml) and treated with a solution of LiOH in water (2 ml). The reaction was stirred overnight. The reaction was cooled and water was added to the solution. The solution was then acidified using 1N HCl to pH = 3 then extracted with ethyl acetate. The organic layer was concentrated to afford the desired carboxylic acid (0.239 g, 51 %). Purification of the crude material using reverse phase chromatography (7:3 acetonitrile: water) afforded the desired acid (0.033 mg, 7%).  $C_{29}H_{38}N_2O_5$  (MW = 494.28); mass spectroscopy ( $MH^+$ ) = 495.3

15

**Example 134**Step A


The diester from **Example 14, Step B** (10 g, 0.032 mol) was dissolved in dioxane. A solution of LiOH in water was

20




added, dropwise, over a period of ten minutes. The reaction was allowed to stir for two hours. The solvent was concentrated and the resulting residue was redissolved in water then washed with ether. The aqueous solution was acidified with 5N HCl and extracted with ethyl acetate (2X). The combined organic layers were washed with brine, dried over Na<sub>2</sub>SO<sub>4</sub>, filtered, and concentrated to afford the desired acid as a yellow oil (9.61 g, 93%). C<sub>18</sub>H<sub>26</sub>O<sub>5</sub> (MW = 322.18); mass spectroscopy (MH<sup>+</sup>) = 321.0

Step B



The acid from **Step A** (9.60 g, 0.0298 mol) was combined with *N*-methyl, *O*-methyl hydroxyl amine, EDAC (6.83 g, 0.0358 mol), HOBT (4.02 g, 0.0298 mol), and DIEA (10 ml, 0.075 mol) in THF (80 ml). The reaction was allowed to stir overnight at room temperature. The solvent was concentrated and the resulting material was redissolved in ethyl acetate and extracted with 1N HCl followed by brine. The organic layer was concentrated. Purification by flash chromatography (4:1 hexane: ethyl acetate) gave the Weinreb amide (7.23 g, 66%). C<sub>26</sub>H<sub>31</sub>NO<sub>5</sub> (MW = 365.22); mass spectroscopy (MH<sup>+</sup>) = 366.22

Step C




A THF solution (50 ml) of the Weinreb amide from **Step B** (3.4 g, 0.0093 mol) was cooled to -78°C and methyl magnesium bromide (3.0 M in ether, 15 ml, 0.046 mol) was added to the

solution. After stirring for two hours, 1N HCl was added to quench the reaction. The reaction mixture was extracted with ether. The ethereal layer was concentrated.


Purification by flash chromatography (9:1 hexanes: ethyl acetate) yielded the methyl ketone as a colorless oil (1.77 g, 60%).  $C_{19}H_{28}O_4$  (MW = 320.20); mass spectroscopy (MH+) = 321.0

#### Step D



The methyl ketone from **Step C** (1.77 g, 0.0055 mol) was combined with potassium cyanide (0.792 g, 0.0122 mol) and ammonium carbonate (2.65 g, 0.0277 mol) in 1:1 methanol: water (20:20 ml) and stirred overnight at 50°C. The reaction mixture was poured into water and extracted with ethyl acetate. The organic layer was washed with brine then concentrated to afford the hydantoin as a white solid (2.11 g, 99%) which was carried forth without further purification.  $C_{21}H_{30}N_2O_5$  (MW = 390.22); mass spectroscopy (MH+) = 391.3

#### Step E




The hydantoin from **Step D** (0.600 g, 0.0015 mol) was dissolved in DMF (10 ml) and treated with  $K_2CO_3$  (0.849 g, 0.0060 mol) and magnesium sulfate (0.271 g, 0.0023 mol)



followed by 3, 4 dimethyl benzyl chloride (0.245 ml, 0.00169 mol). The reaction was stirred overnight at 45°C .

Hydrochloric acid (5N) was added and the mixture was extracted with ethyl acetate (2X). The organic layer was concentrated. Purification of the crude material by flash chromatography (4: 1 ethyl acetate: hexanes; 2:1 ethyl acetate: hexanes) give the alkylated hydantoin (0.560 g, 73%).  $C_{30}H_{40}N_2O_5$  (MW = 508.29); mass spectroscopy (MH<sup>+</sup>) = 509.4

#### Step F



The ester from **Step E** (0.560 g, 0.0011 mol) was dissolved in dichloromethane (10 ml) and treated with trifluoroacetic acid (0.425 ml). The reaction was stirred overnight. The solution was concentrated and the resulting residue was redissolved in water and extracted with methylene chloride. The organic layer was extracted with 2N NaOH. While allowing the layers to separate, a white precipitate crashed out of solution. The solid was filtered; spectral data indicated that the solid was the desired carboxylic acid (0.170 g, 34 %).  $C_{26}H_{32}N_2O_5$  (MW = 452.23); mass spectroscopy (MH<sup>+</sup>) = 453.3


**Example 135**Step A

5

The diester from **Example 14, Step B** (10 g, 0.032 mol) was dissolved in dioxane. A solution of LiOH in water was added, dropwise, over a period of ten minutes. The reaction was allowed to stir for two hours. The solvent was concentrated and the resulting residue was redissolved in water then washed with ether. The aqueous solution was acidified with 5N HCl and extracted with ethyl acetate (2X). The combined organic layers were washed with brine, dried over Na<sub>2</sub>SO<sub>4</sub>, filtered, and concentrated to afford the desired acid as a yellow oil (9.61 g, 93%). C<sub>18</sub>H<sub>26</sub>O<sub>5</sub> (MW = 322.18); mass spectroscopy (MH<sup>-</sup>) = 321.0

10

15

Step B


20

The acid from **Step A** (9.60 g, 0.0298 mol) was combined with *N*-methyl, *O*-methyl hydroxyl amine, EDAC (6.83 g, 0.0358 mol), HOBt (4.02 g, 0.0298 mol), and DIEA (10 ml, 0.075 mol) in THF (80 ml). The reaction was allowed to stir overnight at room temperature. The solvent was concentrated and the

resulting material was redissolved in ethyl acetate and extracted with 1N HCl followed by brine. The organic layer was concentrated. Purification by flash chromatography (4:1 hexane: ethyl acetate) gave the Weinreb amide (7.23 g, 66%).

5  $C_{20}H_{31}NO_5$  (MW = 365.22); mass spectroscopy (MH+) = 366.22


Step C



A THF solution (50 ml) of the Weinreb amide from **Step B** (3.4 g, 0.0093 mol) was cooled to  $-78^{\circ}\text{C}$  and methyl magnesium bromide (3.0 M in ether, 15 ml, 0.046 mol) was added to the solution. After stirring for two hours, 1N HCl was added to quench the reaction. The reaction mixture was extracted with ether. The ethereal layer was concentrated.

15 Purification by flash chromatography (9:1 hexanes: ethyl acetate) yielded the methyl ketone as a colorless oil (1.77 g, 60%).  $C_{19}H_{28}O_4$  (MW = 320.20); mass spectroscopy (MH+) = 321.0


Step D



The methyl ketone from **Step C** (1.77 g, 0.0055 mol) was combined with potassium cyanide (0.792 g, 0.0122 mol) and ammonium carbonate (2.65 g, 0.0277 mol) in 1:1 methanol: water (20:20 ml) and stirred overnight at  $50^{\circ}\text{C}$ . The reaction mixture was poured into water and extracted with ethyl acetate. The organic layer was washed with brine then concentrated to afford the hydantoin as a white solid (2.11

g, 99%) which was carried forth without further purification.  $C_{21}H_{30}N_2O_5$  (MW = 390.22); mass spectroscopy (MH+) = 391.3


5

Step E

The hydantoin from **Step D** (0.600 g, 0.0015 mol) was dissolved in DMF (10 ml) and treated with  $K_2CO_3$  (0.849 g, 0.0060 mol) and magnesium sulfate (0.271 g, 0.0023 mol) followed by 4-ethyl benzyl chloride (0.252 ml, 0.00169 mol). The reaction was stirred overnight at room temperature. The reaction mixture was poured into 5N HCl and extracted with ethyl acetate (2X). The organic layer was concentrated. Purification of the crude material by flash chromatography (2:1 ethyl acetate: hexanes) gave the alkylated hydantoin as a colorless oil (0.752 g, 98%).  $C_{30}H_{40}N_2O_5$  (MW = 508.29); mass spectroscopy (MH+) = 509.3



15

20

Step F

The ester from **Step E** (0.752 g, 0.00148 mol) was dissolved in dichloromethane (20 ml) and treated with trifluoroacetic acid (0.570 ml, 0.0074 mol). The reaction was stirred

overnight. The solution was concentrated. Purification of the crude material by flash chromatography (2:1 hexanes: ethyl acetate) afforded the desired product as a white solid (0.358 g, 54%).  $C_{26}H_{32}N_2O_5$  (MW = 452.23); mass spectroscopy (MH<sup>+</sup>) = 453.3

**Example 136**Step A

10


A THF (15 mL) solution of the hydantoin from **Example 22**, **Step A** (855.1 mg, 1.34 mmol) under an atmosphere of nitrogen was treated with BH<sub>3</sub>-THF (1.0 M in THF, 13.5 mL, 13.5 mmol). The reaction mixture was stirred at room temperature for 3 hours then cooled to 0 °C and carefully treated with saturated aqueous sodium bicarbonate (25 mL). After 2 hr additional bicarbonate (20 mL) was added along with H<sub>2</sub>O (100 mL). The resulting solution was extracted with ethyl acetate (2x100 mL). The combined organic extracts were dried over Na<sub>2</sub>SO<sub>4</sub> and concentrated. Purification by

20

chromatography (2:1 hexanes:ethyl acetate) gave the desired product as a colorless oil (595.4 mg, 71%).

$C_{39}H_{46}N_2O_5$  (MW = 622.81); mass spectroscopy:  $(MH^+) = 623.3$

5

Step B


A TFA solution (10 mL) of the product from **Step A** (595.4 mg, 0.956 mmol) was treated with triethylsilane (618 mL, 3.8 mmol) and stirred at room temperature for 1.5 hour. The reaction mixture was concentrated and the residue diluted with 0.5 N NaOH (50 mL). The aqueous solution was washed with diethyl ether (50 mL), acidified with 5 N HCl, then extracted with ethyl acetate (2x50 mL). The combined organic extracts were dried over  $Na_2SO_4$  and concentrated to afford the desired carboxylic acid as a foam-like solid (417.8 mg, 98%).

10


15

$C_{27}H_{30}N_2O_4$  (MW = 446.55); mass spectroscopy:  $(MH^+) = 447.2$ ,  $(MH^-) = 445.4$


20

**Example 137**



Step A


4-Hydroxyphenethyl alcohol (15.0g, 109 mmol) and ethyl 2-bromoisobutyrate (24ml, 163.5 mmol) were stirred together in  
5 DMF (250ml). Potassium carbonate (44.9g, 325 mmol) and magnesium sulfate (13.1g, 109 mmol) added and the mixture was stirred at 75 °C overnight. The reaction mixture was cooled, filtered over Celite, washing the filter cake with ethyl acetate. The combined filtrates were diluted with  
10 ethyl acetate, washed with water and 2N hydrochloric acid, dried over sodium sulfate, filtered and concentrated. The obtained residue was dissolved in ether and washed with 2N sodium hydroxide and aqueous brine, dried over sodium sulfate. Evaporation of the solvent and subsequent  
15 purification by flash chromatography (hexanes:ethyl acetate) gave the desired product (11.8g).  
 $C_{14}H_{20}O_4$  (MW = 252.3); MS ( $M^+$ , 253.1)

Step B

20 Carbonyldiimidazole (10.0g, 61.7mmol) was stirred in methylene chloride (100ml) and benzylamine (13.5ml, 23.4mmol, in  $CH_2Cl_2$ , 20ml) was added dropwise. An ice bath was placed under the flask during the addition. After  
25 complete addition the mixture was allowed to stir overnight at ambient temperature. The resulting solid was washed with ether and filtered to yield the desired product as a white solid (13.9g).

$C_{15}H_{16}N_2O$  (MW = 240.3); MS ( $M^+$ , 241.0)


Step C



- 5 The product from **Example 137, Step B** (10.0g, 41.7mmol) was stirred in dioxane and oxalyl chloride (3.6ml, 41.7mmol) was added via syringe. The mixture was heated to 75 °C overnight. Evaporation of the solvent and washing the obtained solid with ether gave the desired product as a
- 10 white solid (12.1g).


$C_{17}H_{14}N_2O_3$  (MW = 294.3);  $^1\text{NMR}$

Step D




- 15 The product from **Example 137, Step C** (2.0g, 6.8mmol) was stirred in THF (50ml) and methanol (30ml). Sodium borohydride was added and stirred for 5 minutes. Water (1ml) and concentrated hydrochloric acid (1ml) were added then allowed to stand over sodium sulfate overnight.
- 20 Evaporation of the solvent gave a residue that was dissolved in water and ethyl acetate. Layers were separated and the organic layer was washed with aqueous brine and dried over sodium sulfate. Evaporation of the solvent gave the desired product (2.0g).

25  $C_{17}H_{16}N_2O_3$  (MW = 296.3);  $^1\text{NMR}$

Step E

The products from **Example 137, Step A** (1.7g, 6.8mmol) and **Step D** (1.0g, 3.4mmol) were stirred together in acetonitrile (10ml) with p-toluenesulfonic acid (0.1g, 0.5mmol) and heated to 80 °C for 6 hours. The resulting mixture was concentrated and methylene chloride added. This was washed with water and aqueous brine then dried over sodium sulfate. Evaporation of the solvent and subsequent purification by flash chromatography (hexanes/ethyl acetate) gave the desired product (0.8g).

$C_{31}H_{34}N_2O_6$  (MW = 530.6);  $^1\text{NMR}$

Step F

The product from **Example 137, Step E** (0.25g, 0.47mmol) was dissolved in methanol (2ml), added to lithium hydroxide (0.02g, 1.0mmol, in water, 5ml) and heated to 50 °C for 1 hour, then stirred over two days at ambient temperature. Hydrolysis completed by heating to reflux for 1 hour. The reaction was added to water and extracted with ether. The aqueous layer was acidified with concentrated hydrochloric acid and extracted twice with ethyl acetate. The organic


layers were combined, washed with aqueous brine and dried over sodium sulfate. Evaporation of the solvent gave the desired product (0.19g).

$C_{29}H_{30}N_2O_6$  (MW = 502.6); MS (M-, 501.1)

5

### Example 138

#### 1-Isocyanatomethyl-4-methyl-benzene




A mixture of *p*-methyl benzylamine (25 g, 0.21 mol) in  $CH_2Cl_2$  (200 mL) and aqueous saturated solution of sodium bicarbonate (200 mL) was cooled to 0 °C and then treated dropwise with a 20% solution of phosgene in toluene (100 mL). The reaction was stirred at room temperature for 20 minutes and then the organic layer removed and the aqueous layer back-extracted with  $CH_2Cl_2$ . The organic layer was dried ( $Na_2SO_4$ ) and the solvent removed *in vacuo* to give 20.6 g 1-isocyanatomethyl-4-methyl-benzene that was utilized directly in the next reaction without purification.

20

#### Step B

#### 1-(4-Methyl-benzyl)-3-(2-oxo-tetrahydro-furan-3-yl)-urea



A 0 °C slurry of  $\alpha$ -amino- $\gamma$ -butyrolactone hydrobromide (25 g, 0.137 mol) in THF (200 mL) was combined with 1-isocyanatomethyl-4-methyl-benzene (20.6 g, 0.139 mol) and then treated dropwise with *N,N*-diisopropylethylamine (19.5 g, 0.151 mol). The reaction mixture was stirred for at room temperature for 2 h. The solvent was removed *in vacuo* to


25

give 48.5 g crude 1-(4-methyl-benzyl)-3-(2-oxo-tetrahydro-furan-3-yl)-urea that was utilized directly in the next reaction without purification. MS (ES<sup>+</sup>) Calc'd for C<sub>13</sub>H<sub>17</sub>N<sub>2</sub>O<sub>3</sub> (M +1) 249. Found m/z 249 (100%). <sup>1</sup>H NMR.

5

### Step C


5-(2-Hydroxy-ethyl)-3-(4-methyl-benzyl)-imidazolidine-2,4-dione



- 10 A solution of crude 1-(4-methyl-benzyl)-3-(2-oxo-tetrahydro-furan-3-yl)-urea (48.5 g, assume 0.137 mol) in MeOH (1.5 L) was treated with sodium methoxide (21.1 g, 0.391 mol) and heated to reflux under N<sub>2</sub> for 2 h until done by thin layer chromatography (9:1 CH<sub>2</sub>Cl<sub>2</sub>:MeOH). The reaction was cooled
- 15 and the solvent removed in vacuo to give a residue that was combined with aqueous 1 N HCl (640 mL) and extracted with EtOAc. The organic layer was dried (MgSO<sub>4</sub>) and the solvent removed in vacuo to afford 24.72 g (73%) of crude 5-(2-hydroxy-ethyl)-3-(4-methyl-benzyl)-imidazolidine-2,4-dione
- 20 as a yellow solid which was carried on in the next step without purification. MS (ES<sup>+</sup>) Calc'd for C<sub>13</sub>H<sub>17</sub>N<sub>2</sub>O<sub>3</sub> (M +1) 249. Found m/z 249 (100%). <sup>1</sup>H NMR.

### Step D

- 25 5-[2-(*tert*-Butyl-diphenyl-silanyloxy)-ethyl]-3-(4-methyl-benzyl)-imidazolidine-2,4-dione




A solution of crude 5-(2-hydroxy-ethyl)-3-(4-methyl-benzyl)-imidazolidine-2,4-dione (24.68 g, 99.4 mmol) and imidazole (10.15 g, 0.149 mol) in DMF (250 mL) was treated dropwise with *tert*-butyldiphenylsilyl chloride (27.37 g, 99.6 mmol) and then stirred at room temperature under N<sub>2</sub> for 7 h. The reaction was worked up extractively with aqueous 1 N HCl (300 mL), diethyl ether and brine and the organic layer was dried (MgSO<sub>4</sub>) and the solvent removed *in vacuo* to give crude product which was purified by flash chromatography using a gradient of 5:1 to 2:1 hexanes:EtOAc to afford 23.82 g (49%) of 5-[2-(*tert*-butyl-diphenyl-silanyloxy)-ethyl]-3-(4-methyl-benzyl)-imidazolidine-2,4-dione. MS (ES<sup>+</sup>) Calc'd for C<sub>29</sub>H<sub>35</sub>N<sub>2</sub>O<sub>3</sub>Si (M + 1) 487. Found *m/z* 487 (100%). <sup>1</sup>H NMR.

15

Step E

5-[2-(*tert*-Butyl-diphenyl-silanyloxy)-ethyl]-3-(4-methyl-benzyl)-1-propyl-imidazolidine-2,4-dione




A 0 °C solution of 5-[2-(*tert*-butyl-diphenyl-silanyloxy)-ethyl]-3-(4-methyl-benzyl)-imidazolidine-2,4-dione (14.26 g, 29.3 mmol) in DMF (250 mL) was treated with sodium hydride (60% dispersion, 1.29 g, 32.2 mmol) and warmed to room temperature and stirred under N<sub>2</sub> for 30 minutes. The resultant mixture was cooled to 0 °C and then treated with 1-propyl iodide (5.47 g, 32.2 mmol) and then warmed to room temperature and stirred for 3 h. The reaction was quenched with aqueous 1 N HCl (60 mL) and then worked up extractively with diethyl ether and water. The organic layer was dried (MgSO<sub>4</sub>) and the solvent removed *in vacuo* to give crude product that was purified by flash chromatography using 6:1

hexanes:acetone to afford 14.8 g (96%) 5-[2-(*tert*-butyl-diphenyl-silanyloxy)-ethyl]-3-(4-methyl-benzyl)-1-propyl-imidazolidine-2,4-dione. MS (ES<sup>+</sup>) Calc'd for C<sub>32</sub>H<sub>40</sub>N<sub>2</sub>O<sub>3</sub>Si (M<sup>+</sup>) 529. Found *m/z* 529 (100%). <sup>1</sup>H NMR.

5

#### Step F

4-(2-Hydroxy-ethyl)-1-(4-methyl-benzyl)-3-propyl-1,3-dihydro-imidazol-2-one



10

A solution of 5-[2-(*tert*-butyl-diphenyl-silanyloxy)-ethyl]-3-(4-methyl-benzyl)-1-propyl-imidazolidine-2,4-dione (14.71 g, 27.8 mmol) in EtOH (500 mL) was treated with sodium borohydride (15.78 g, 0.417 mol) in three portions over 6 h and stirred at room temperature under N<sub>2</sub> for an additional 16 h. The thick reaction mixture was cooled in an ice bath and slowly quenched with aqueous 1 N HCl (~1L) over a 1 h period. The reaction mixture was then worked up extractively with EtOAc, water and brine and the organic layer dried (MgSO<sub>4</sub>) and the solvent removed *in vacuo* to give 16.79 g crude 4-[2-(*tert*-butyl-diphenyl-silanyloxy)-ethyl]-1-(4-methyl-benzyl)-3-propyl-1,3-dihydro-imidazol-2-one that was de-silylated immediately.

15

20

25


30

The 4-[2-(*tert*-butyl-diphenyl-silanyloxy)-ethyl]-1-(4-methyl-benzyl)-3-propyl-1,3-dihydro-imidazol-2-one (16.79 g, assumed 27.8 mmol) was dissolved in THF (160 mL) and then treated with a 1 M solution of tetrabutylammonium fluoride in THF (55.6 mL, 55.5 mmol) and the reaction stirred at room temperature under N<sub>2</sub> for 20 minutes. The reaction was worked up extractively with EtOAc and water and the organic

layer dried ( $\text{MgSO}_4$ ) and the solvent removed in vacuo to give crude product that was purified by flash chromatography using 3% of MeOH in  $\text{CH}_2\text{Cl}_2$  to afford 4.65 g (61%) 4-(2-hydroxy-ethyl)-1-(4-methyl-benzyl)-3-propyl-1,3-dihydro-imidazol-2-one. MS ( $\text{ES}^+$ ) Calc'd for  $\text{C}_{16}\text{H}_{23}\text{N}_2\text{O}_2$  ( $M + 1$ ) 275. Found  $m/z$  275 (100%).  $^1\text{H}$  NMR.

#### Step G


10 **Toluene-4-sulfonic acid 2-[1-(4-methyl-benzyl)-2-oxo-3-propyl-2,3-dihydro-1H-imidazol-4-yl]-ethyl ester**



A solution of 4-(2-hydroxy-ethyl)-1-(4-methyl-benzyl)-3-propyl-1,3-dihydro-imidazol-2-one (0.126 g, 0.459 mmol), pyridine (0.146 g, 1.85 mmol) and 4-dimethyl amino pyridine (0.017 g, 0.139 mmol) in  $\text{CH}_2\text{Cl}_2$  (10 mL) was treated with *p*-toluenesulfonic anhydride (0.240 g, 0.735 mmol) and the reaction stirred at room temperature for under  $\text{N}_2$  for 1.5 h. The reaction mixture was washed with aqueous 1 N HCl (4 mL), the organic layer was dried ( $\text{MgSO}_4$ ), and the solvent removed in vacuo to afford crude product that was purified by flash chromatography using 100%  $\text{CH}_2\text{Cl}_2$  then 2.5% MeOH in  $\text{CH}_2\text{Cl}_2$  to afford 0.194 g (98%) toluene-4-sulfonic acid 2-[1-(4-methyl-benzyl)-2-oxo-3-propyl-2,3-dihydro-1H-imidazol-4-yl]-ethyl ester. MS ( $\text{ES}^+$ ) Calc'd for  $\text{C}_{23}\text{H}_{29}\text{N}_2\text{O}_4\text{S}$  ( $M + 1$ ) 429. Found  $m/z$  429 (100%).  $^1\text{H}$  NMR.

#### Step H

**2-Methyl-2-(4-{2-[1-(4-methyl-benzyl)-2-oxo-3-propyl-2,3-dihydro-1H-imidazol-4-yl]-ethoxy}-phenoxy)-propionic acid**






A mixture of toluene-4-sulfonic acid 2-[1-(4-methyl-benzyl)-2-oxo-3-propyl-2,3-dihydro-1H-imidazol-4-yl]-ethyl ester (0.194 g, 1.51 mmol), 2-(4-hydroxy-phenoxy)-2-methyl-propionic acid ethyl ester (0.088 g, 0.392mmol) and Cs<sub>2</sub>CO<sub>3</sub> (0.156 g, 0.478 mmol) in DMF (15 mL) was heated at 55 °C for 5 under N<sub>2</sub> for 16 h. The reaction mixture was cooled to room temperature, quenched with aqueous 1 N HCl (2 mL), and worked up extractively with EtOAc and water. The organic layer was dried (MgSO<sub>4</sub>) and the solvent removed *in vacuo* to 10 afford crude product that was purified by flash chromatography using a gradient of 4:1 to 1:1 hexanes:acetone to afford 0.036 g (19%) 2-methyl-2-(4-{2-[1-(4-methyl-benzyl)-2-oxo-3-propyl-2,3-dihydro-1H-imidazol-4-yl]-ethoxy}-phenoxy)-propionic acid ethyl ester.

15 The 2-methyl-2-(4-{2-[1-(4-methyl-benzyl)-2-oxo-3-propyl-2,3-dihydro-1H-imidazol-4-yl]-ethoxy}-phenoxy)-propionic acid ethyl ester (0.036g, 0.075 mmol) was combined with aqueous 5 N NaOH (0.15 mL) in EtOH (5 mL) and heated to reflux under N<sub>2</sub> for 1 h. The reaction was cooled and the 20 solvent removed *in vacuo* to give a residue that was acidified with aqueous 1 N HCl (1 mL). The mixture was extracted with Et<sub>2</sub>O and water, the organic layer dried (MgSO<sub>4</sub>), and the solvent removed *in vacuo* to afford 0.024 g (71%) 2-methyl-2-(4-{2-[1-(4-methyl-benzyl)-2-oxo-3-propyl-2,3-dihydro-1H-imidazol-4-yl]-ethoxy}-phenoxy)-propionic 25 acid. HRMS (ES<sup>+</sup>) *m/z* exact mass calcd for C<sub>26</sub>H<sub>33</sub>N<sub>2</sub>O<sub>5</sub> [M + 1] 453.2389, found 453.2387. <sup>1</sup>H NMR.

#### Example 139

30 **2-Methoxy-2-methyl-3-(4-{2-[1-(4-methyl-benzyl)-2-oxo-3-propyl-2,3-dihydro-1H-imidazol-4-yl]-ethoxy}-phenyl)-propionic acid**




A mixture of toluene-4-sulfonic acid 2-[1-(4-methyl-benzyl)-2-oxo-3-propyl-2,3-dihydro-1H-imidazol-4-yl]-ethyl ester (0.079 g, 0.184 mmol), 3-(4-hydroxy-phenyl)-2-methoxy-2-methyl-propionic acid ethyl ester (0.038 g, 0.169mmol) and  $K_2CO_3$  (0.047 g, 0.340 mmol) in EtOH (8 mL) was heated at 75 °C under  $N_2$  for 16 h. The reaction mixture was cooled to room temperature and quenched with aqueous 1 N HCl (1 mL). The solvent was removed *in vacuo* to give a residue that was diluted with water and extracted with EtOAc. The organic layer was dried ( $MgSO_4$ ) and the solvent removed *in vacuo* to afford crude product that was purified by flash chromatography using a 2:1 hexanes:acetone to afford 0.036 g (44%) 2-methoxy-2-methyl-3-(4-{2-[1-(4-methyl-benzyl)-2-oxo-3-propyl-2,3-dihydro-1H-imidazol-4-yl]-ethoxy}-phenyl)-propionic acid ethyl ester.

The 2-methoxy-2-methyl-3-(4-{2-[1-(4-methyl-benzyl)-2-oxo-3-propyl-2,3-dihydro-1H-imidazol-4-yl]-ethoxy}-phenyl)-propionic acid ethyl ester (0.015g, 0.031 mmol) was combined with aqueous 5 N NaOH (0.1 mL) in EtOH (3 mL) and heated to reflux under  $N_2$  for 1 h. The reaction was cooled and the solvent removed *in vacuo* to give a residue that was acidified with aqueous 1 N HCl (3 mL). The mixture was extracted with  $Et_2O$  and water, the organic layer dried ( $MgSO_4$ ), and the solvent removed *in vacuo* to afford 0.015 g (100%) 2-methoxy-2-methyl-3-(4-{2-[1-(4-methyl-benzyl)-2-oxo-3-propyl-2,3-dihydro-1H-imidazol-4-yl]-ethoxy}-phenyl)-propionic acid. MS ( $ES^+$ )  $m/z$  calcd for  $C_{26}H_{32}N_2O_5$  [ $M + 1$ ] 453, found 453.  $^1H$  NMR.

## Example 140

2-Methyl-3-(4-{2-[1-(4-methyl-benzyl)-2-oxo-3-propyl-2,3-dihydro-1H-imidazol-4-yl]-ethoxy}-phenyl)-2-phenoxy-propionic acid




5 A mixture of toluene-4-sulfonic acid 2-[1-(4-methyl-benzyl)-2-oxo-3-propyl-2,3-dihydro-1H-imidazol-4-yl]-ethyl ester (0.029 g, 0.0677 mmol), 3-(4-Hydroxy-phenyl)-2-methyl-2-phenoxy-propionic acid ethyl ester (0.018 g, 0.0599mmol) and  
10  $\text{Cs}_2\text{CO}_3$  (0.024 g, 0.0736 mmol) in DMF (3 mL) was heated at 55 °C under  $\text{N}_2$  for 16 h. The reaction mixture was cooled to room temperature and quenched with aqueous 1 N HCl (0.3 mL) and then was diluted with water and extracted with  $\text{Et}_2\text{O}$ . The organic layer was dried ( $\text{MgSO}_4$ ) and the solvent removed  
15 *in vacuo* to afford crude product that was purified by flash chromatography using a gradient of 4:1 to hexanes:EtOAc to 2:1 EtOAc:hexanes to afford 0.011 g (32%) 2-methyl-3-(4-{2-[1-(4-methyl-benzyl)-2-oxo-3-propyl-2,3-dihydro-1H-imidazol-4-yl]-ethoxy}-phenyl)-2-phenoxy-propionic acid ethyl ester.  
20 The 2-methyl-3-(4-{2-[1-(4-methyl-benzyl)-2-oxo-3-propyl-2,3-dihydro-1H-imidazol-4-yl]-ethoxy}-phenyl)-2-phenoxy-propionic acid ethyl ester (0.011g, 0.020 mmol) was combined with aqueous 5 N NaOH (0.2 mL) in EtOH (3 mL) and heated to reflux under  $\text{N}_2$  for 1 h. The reaction was cooled  
25 and the solvent removed *in vacuo* to give a residue that was acidified with aqueous 1 N HCl (2 mL). The mixture was extracted with  $\text{Et}_2\text{O}$  and water, the organic layer dried ( $\text{MgSO}_4$ ), and the solvent removed *in vacuo* to afford 0.007 g (67%) 2-methyl-3-(4-{2-[1-(4-methyl-benzyl)-2-oxo-3-propyl-

2,3-dihydro-1H-imidazol-4-yl]-ethoxy}-phenyl)-2-phenoxy-propionic acid. HRMS (ES<sup>+</sup>) *m/z* exact mass calcd for C<sub>32</sub>H<sub>37</sub>N<sub>2</sub>O<sub>5</sub> [M + 1] 529.2702, found 529.2715. <sup>1</sup>H NMR.

5

**Example 141**Step A

**1-(3,4-Dimethyl-benzyl)-4-[3-(4-methoxy-phenyl)-propyl]-5-methyl-3-propyl-1,3-dihydro-imidazol-2-one**




10

A 0 °C solution of 3-(3,4-dimethyl-benzyl)-5-[3-(4-methoxy-phenyl)-propyl]-1-propyl-imidazolidine-2,4-dione (0.155 g, 0.379 mmol) in THF (6 mL) was treated dropwise with a 3 molar solution of methyl magnesium bromide in Et<sub>2</sub>O (0.76 mL, 2.28 mmol) and then stirred at 0 °C under N<sub>2</sub> for 1 h. The reaction was quenched with 1 N HCl (5 mL) and stirred at room temperature for 15 minutes to effect elimination and then was worked up extractively with EtOAc and water. The organic layer dried (MgSO<sub>4</sub>), and the solvent removed in vacuo to afford 0.155 g (100%) 1-(3,4-dimethyl-benzyl)-4-[3-(4-methoxy-phenyl)-propyl]-5-methyl-3-propyl-1,3-dihydro-imidazol-2-one that was utilized without purification. MS (ES<sup>+</sup>) Calc'd for C<sub>26</sub>H<sub>35</sub>N<sub>2</sub>O<sub>2</sub> (M + 1) 407. Found *m/z* 407 (100%). <sup>1</sup>H NMR.

25

Step B

**2-(4-{3-[1-(3,4-Dimethyl-benzyl)-5-methyl-2-oxo-3-propyl-2,3-dihydro-1H-imidazol-4-yl]-propyl}-phenoxy)-2-methyl-propionic acid ethyl ester**




A 0 °C solution of 1-(3,4-dimethyl-benzyl)-4-[3-(4-methoxy-phenyl)-propyl]-5-methyl-3-propyl-1,3-dihydro-imidazol-2-one (0.145 g, 0.357 mmol) in CH<sub>2</sub>Cl<sub>2</sub> (6 mL) was treated dropwise with BBr<sub>3</sub> (0.10 mL, 1.06 mmol) and then stirred at 0 °C  
5 under N<sub>2</sub> for 30 minutes. The reaction was diluted with Et<sub>2</sub>O and then quenched with ice and then water. The organic layer was dried (MgSO<sub>4</sub>), and the solvent removed *in vacuo* to afford 0.169 g (100%) 1-(3,4-dimethyl-benzyl)-4-[3-(4-hydroxy-phenyl)-propyl]-5-methyl-3-propyl-1,3-dihydro-  
10 imidazol-2-one that was utilized without purification.

The crude 1-(3,4-dimethyl-benzyl)-4-[3-(4-hydroxy-phenyl)-propyl]-5-methyl-3-propyl-1,3-dihydro-imidazol-2-one (0.169 g) was combined with ethyl 2-bromoisobutyrate (0.366 g, 1.87 mmol), MgSO<sub>4</sub> (0.043 g, 0.357 mmol) and 325 mesh K<sub>2</sub>CO<sub>3</sub>  
15 (0.197 g, 1.43 mmol) in ethanol (8 mL) and heated at 75 °C for 16 h under N<sub>2</sub>. The reaction was cooled and the solvent removed *in vacuo*. The residue was acidified with 1 N HCl (10 mL) and extracted with EtOAc and water. The organic layer was dried (MgSO<sub>4</sub>), and the solvent removed *in vacuo* to  
20 afford crude product that was purified by flash chromatography using 2:1 to hexanes:acetone to afford 0.124 g (69%) 2-(4-{3-[1-(3,4-dimethyl-benzyl)-5-methyl-2-oxo-3-propyl-2,3-dihydro-1H-imidazol-4-yl]-propyl}-phenoxy)-2-methyl-propionic acid ethyl ester. MS (ES<sup>+</sup>) Calc'd for  
25 C<sub>31</sub>H<sub>43</sub>N<sub>2</sub>O<sub>4</sub> (M +1) 507. Found *m/z* 507 (100%). <sup>1</sup>H NMR.

#### Step C

2-(4-{3-[1-(3,4-Dimethyl-benzyl)-5-methyl-2-oxo-3-propyl-2,3-dihydro-1H-imidazol-4-yl]-propyl}-phenoxy)-2-methyl-  
30 propionic acid




A solution of 2-(4-{3-[1-(3,4-dimethyl-benzyl)-5-methyl-2-oxo-3-propyl-2,3-dihydro-1H-imidazol-4-yl]-propyl}-phenoxy)-2-methyl-propionic acid ethyl ester (0.124 g, 0.244 mmol) in  
 5 EtOH (10 mL) was treated with aqueous 5 N NaOH (0.5 mL) and heated to reflux under N<sub>2</sub> for 1 h. The reaction was cooled and the solvent removed *in vacuo* to give a residue that was acidified with aqueous 1 N HCl. The mixture was extracted with Et<sub>2</sub>O and water, the organic layer dried (MgSO<sub>4</sub>), and  
 10 the solvent removed *in vacuo* to afford 0.104 g (89%) 2-(4-{3-[1-(3,4-dimethyl-benzyl)-5-methyl-2-oxo-3-propyl-2,3-dihydro-1H-imidazol-4-yl]-propyl}-phenoxy)-2-methyl-propionic acid. HRMS (ES<sup>+</sup>) *m/z* exact mass calcd for C<sub>29</sub>H<sub>39</sub>N<sub>2</sub>O<sub>4</sub> [M + 1] 479.2910, found 479.2917. <sup>1</sup>H NMR.

15

**Example 142**

**2-(4-{3-[1-(3,4-Dimethyl-benzyl)-5-ethyl-2-oxo-3-propyl-2,3-dihydro-1H-imidazol-4-yl]-propyl}-phenoxy)-2-methyl-propionic acid**

20




The procedures herein were utilized with ethyl magnesium bromide to afford 2-(4-{3-[1-(3,4-dimethyl-benzyl)-5-ethyl-2-oxo-3-propyl-2,3-dihydro-1H-imidazol-4-yl]-propyl}-phenoxy)-2-methyl-propionic acid. HRMS (ES<sup>+</sup>) *m/z* exact mass  
 25 calcd for C<sub>30</sub>H<sub>41</sub>N<sub>2</sub>O<sub>4</sub> [M + 1] 493.3066, found 493.3079. <sup>1</sup>H NMR.

## Cyclic Urea Experimentals

## General Procedure

## Step A


- 5    2-(4-{2-[1-(4-Methoxy-benzyl)-3-methyl-2-oxo-imidazolidin-4-yl]-ethoxy}-phenoxy)-2-methyl-propionic acid ethyl ester



- A mixture of toluene-4-sulfonic acid 2-[1-(4-methoxy-benzyl)-3-methyl-2-oxo-imidazolidin-4-yl]-ethyl ester (3.10 g, 7.41 mmol), 2-(4-hydroxy-phenoxy)-2-methyl-propionic acid ethyl ester (1.50 g, 6.69 mmol) and  $\text{Cs}_2\text{CO}_3$  (2.62 g, 8.04 mmol) in DMF (70 mL) was heated at 65 °C for under  $\text{N}_2$  for 16 h. The reaction mixture was cooled to room temperature, quenched with aqueous 1 N HCl (25 mL), and worked up  
15 extractively with EtOAc and water. The organic layer was dried ( $\text{MgSO}_4$ ) and the solvent removed *in vacuo* to afford crude product that was purified by flash chromatography using 5:1 hexanes:acetone to afford 2.92 g (93%) 2-(4-{2-[1-(4-methoxy-benzyl)-3-methyl-2-oxo-imidazolidin-4-yl]-ethoxy}-phenoxy)-2-methyl-propionic acid ethyl ester. MS (ES<sup>+</sup>) Calc'd for  $\text{C}_{36}\text{H}_{47}\text{N}_2\text{O}_6$  (M +1) 603. Found  $m/z$  603 (100%).  
20 <sup>1</sup>H NMR.

## Step B

- 25    2-Methyl-2-{4-[2-(3-methyl-2-oxo-imidazolidin-4-yl)-ethoxy]-phenoxy}-propionic acid ethyl ester




- A mixture of 2-(4-{2-[1-(4-methoxy-benzyl)-3-methyl-2-oxo-imidazolidin-4-yl]-ethoxy}-phenoxy)-2-methyl-propionic acid ethyl ester (2.72 g, 5.78 mmol) and triethylsilane (1.34 g, 11.5 mmol) was treated with trifluoroacetic acid (70 mL) and  
30

stirred at room temperature under N<sub>2</sub> for 5 h. The reaction mixture was diluted with water and then extracted with ethyl acetate. The organic layer was dried (MgSO<sub>4</sub>) and the solvent removed *in vacuo* to afford crude product that was purified by flash chromatography using 98:2 CH<sub>2</sub>Cl<sub>2</sub>:MeOH to afford 2.40 g (100%) 2-methyl-2-{4-[2-(3-methyl-2-oxo-imidazolidin-4-yl)-ethoxy]-phenoxy}-propionic acid ethyl ester. R<sub>f</sub> = 0.09 (1:1 hexanes:acetone); MS (ES<sup>+</sup>) Calc'd for C<sub>18</sub>H<sub>27</sub>N<sub>2</sub>O<sub>5</sub> (M +1) 351. Found *m/z* 351 (100%). <sup>1</sup>H NMR.

#### Step C

**2-(4-{2-[1-(4-Methanesulfonyl-benzyl)-3-methyl-2-oxo-imidazolidin-4-yl]-ethoxy}-phenoxy)-2-methyl-propionic acid ethyl ester**



A solution of 2-methyl-2-{4-[2-(3-methyl-2-oxo-imidazolidin-4-yl)-ethoxy]-phenoxy}-propionic acid ethyl ester (0.143 g, 0.408 mmol) in dry DMF (4 mL) was treated a 60% suspension of NaH (0.033 g, 0.825 mmol) and the resultant mixture was stirred at room temperature for 20 minutes under N<sub>2</sub>. The reaction mixture was cooled to 0 °C and then treated with *p*-methanesulfonylbenzyl chloride and then warmed to room temperature and stirred for 16 h.

The reaction was acidified with aqueous 1 N HCl (4 mL), diluted with water and extracted with Et<sub>2</sub>O. The organic layer was dried (MgSO<sub>4</sub>) and the solvent removed *in vacuo* to afford crude product that was purified by column chromatography using a gradient of 4:1 to 1:1 hexanes:acetone to afford 0.090 g (42%) 2-(4-{2-[1-(4-methanesulfonyl-benzyl)-3-methyl-2-oxo-imidazolidin-4-yl]-




ethoxy}-phenoxy)-2-methyl-propionic acid ethyl ester.  $R_f$  = 0.40 (1:1 hexanes:acetone)MS (ES<sup>+</sup>) Calc'd for C<sub>26</sub>H<sub>35</sub>N<sub>2</sub>O<sub>7</sub>S (M+1) 518. Found  $m/z$  518 (100%). <sup>1</sup>H NMR.

5

Step D

2-(4-{2-[1-(4-Methanesulfonyl-benzyl)-3-methyl-2-oxo-imidazolidin-4-yl]-ethoxy}-phenoxy)-2-methyl-propionic acid




A solution of 2-(4-{2-[1-(4-methanesulfonyl-benzyl)-3-methyl-2-oxo-imidazolidin-4-yl]-ethoxy}-phenoxy)-2-methyl-propionic acid ethyl ester (0.087 g, 0.168 mmol) in ethanol (8 mL) was treated with aqueous 5 N NaOH (1 mL) and heated to reflux 1 h. The reaction mixture was cooled, the solvent removed *in vacuo*. The resultant residue was acidified with aqueous 1 N HCl (5 mL) and extracted with CH<sub>2</sub>Cl<sub>2</sub>. The organic layer was dried (MgSO<sub>4</sub>) and the solvent removed *in vacuo* to afford 0.043 g (52%) 2-(4-{2-[1-(4-methanesulfonyl-benzyl)-3-methyl-2-oxo-imidazolidin-4-yl]-ethoxy}-phenoxy)-2-methyl-propionic acid. HRMS (ES<sup>+</sup>)  $m/z$  exact mass calcd for C<sub>24</sub>H<sub>31</sub>N<sub>2</sub>O<sub>7</sub>S 491.1852, found 491.1879. <sup>1</sup>H NMR.

20

**Example 143**

2-(4-{2-[1-(4-Methoxy-benzyl)-3-methyl-2-oxo-imidazolidin-4-yl]-ethoxy}-phenoxy)-2-methyl-propionic acid



25


A solution of 2-(4-{2-[1-(4-methoxy-benzyl)-3-methyl-2-oxo-imidazolidin-4-yl]-ethoxy}-phenoxy)-2-methyl-propionic acid ethyl ester (0.198 g, 0.421 mmol) in ethanol (10 mL) was

treated with aqueous 5 N NaOH (1 mL) and heated to reflux 1.5 h. The reaction mixture was cooled, the solvent removed *in vacuo*. The resultant residue was acidified with aqueous 1 N HCl (10 mL) and extracted with Et<sub>2</sub>O. The organic layer  
5 was dried (MgSO<sub>4</sub>) and the solvent removed *in vacuo* to afford 0.147 g (79%) 2-(4-{2-[1-(4-methoxy-benzyl)-3-methyl-2-oxo-imidazolidin-4-yl]-ethoxy}-phenoxy)-2-methyl-propionic acid. MS (ES<sup>+</sup>) Calc'd for C<sub>24</sub>H<sub>31</sub>N<sub>2</sub>O<sub>6</sub> (M+1) 443. Found *m/z* 443 (100%). <sup>1</sup>H NMR.

10

**Example 144**


**(R)-2-Methyl-2-(4-{2-[3-methyl-2-oxo-1-(4-trifluoromethyl-benzyl)-imidazolidin-4-yl]-ethoxy}-phenoxy)-propionic acid**



15 The procedures herein were utilized with 4-trifluoromethylbenzyl bromide to prepare (R)-2-methyl-2-(4-{2-[3-methyl-2-oxo-1-(4-trifluoromethyl-benzyl)-imidazolidin-4-yl]-ethoxy}-phenoxy)-propionic acid. HRMS (ES<sup>+</sup>) *m/z* exact mass calcd for C<sub>24</sub>H<sub>28</sub>N<sub>2</sub>O<sub>5</sub>F<sub>3</sub> 481.1950, found  
20 481.1960. <sup>1</sup>H NMR.

**Example 145**

**(R)-2-(4-{2-[1-(3,4-Difluoro-benzyl)-3-methyl-2-oxo-imidazolidin-4-yl]-ethoxy}-phenoxy)-2-methyl-propionic acid**




25

The procedures herein were utilized with 3,4-difluorobenzyl bromide to prepare (R)- 2-(4-{2-[1-(3,4-difluoro-benzyl)-3-methyl-2-oxo-imidazolidin-4-yl]-ethoxy}-phenoxy)-2-methyl-

propionic acid. HRMS (ES<sup>+</sup>) *m/z* exact mass calcd for C<sub>23</sub>H<sub>27</sub>N<sub>2</sub>O<sub>5</sub>F<sub>2</sub> 449.1888, found 449.1888. <sup>1</sup>H NMR.

#### Example 146


5     **(R)-2-Methyl-2-{4-[2-(3-methyl-2-oxo-1-quinolin-2-ylmethyl-imidazolidin-4-yl)-ethoxy]-phenoxy}-propionic acid**



The procedures herein were utilized with 2-(chloromethyl)quinoline to prepare (R)-2-methyl-2-{4-[2-(3-methyl-2-oxo-1-quinolin-2-ylmethyl-imidazolidin-4-yl)-ethoxy]-phenoxy}-propionic acid. HRMS (ES<sup>+</sup>) *m/z* exact mass calcd for C<sub>26</sub>H<sub>30</sub>N<sub>3</sub>O<sub>5</sub> 464.2185, found 464.2193. <sup>1</sup>H NMR.

#### Example 147


15     **(R)-2-(4-{2-[1-(3,5-Difluoro-benzyl)-3-methyl-2-oxo-imidazolidin-4-yl]-ethoxy}-phenoxy)-2-methyl-propionic acid**



The procedures herein were utilized with 3,5-difluorobenzyl bromide to prepare (R)-2-(4-{2-[1-(3,5-difluoro-benzyl)-3-methyl-2-oxo-imidazolidin-4-yl]-ethoxy}-phenoxy)-2-methyl-propionic acid. HRMS (ES<sup>+</sup>) *m/z* exact mass calcd for C<sub>23</sub>H<sub>27</sub>N<sub>2</sub>O<sub>5</sub>F<sub>2</sub> 449.1888, found 449.1913. <sup>1</sup>H NMR.

#### Example 148


25     **(R)-2-(4-{2-[1-(3-Methoxy-benzyl)-3-methyl-2-oxo-imidazolidin-4-yl]-ethoxy}-phenoxy)-2-methyl-propionic acid**



The procedures herein were utilized with 3-methoxybenzyl  
bromide to prepare (R)-2-(4-{2-[1-(3-methoxy-benzyl)-3-  
methyl-2-oxo-imidazolidin-4-yl]-ethoxy}-phenoxy)-2-methyl-  
5 propionic acid. HRMS (ES<sup>+</sup>) *m/z* exact mass calcd for  
C<sub>24</sub>H<sub>31</sub>N<sub>2</sub>O<sub>6</sub> 443.2182, found 443.2182. <sup>1</sup>H NMR.

#### Example 149


(R)-2-(4-{2-[1-(4-Chloro-benzyl)-3-methyl-2-oxo-  
10 imidazolidin-4-yl]-ethoxy}-phenoxy)-2-methyl-propionic acid




The procedures herein were utilized with 4-chlorobenzyl  
bromide to prepare (R)-2-(4-{2-[1-(4-chloro-benzyl)-3-  
methyl-2-oxo-imidazolidin-4-yl]-ethoxy}-phenoxy)-2-methyl-  
15 propionic acid. HRMS (ES<sup>+</sup>) *m/z* exact mass calcd for  
C<sub>23</sub>H<sub>28</sub>N<sub>2</sub>O<sub>5</sub>Cl 447.1687, found 447.1667. <sup>1</sup>H NMR.

#### Example 150


(R)-2-Methyl-2-{4-[2-(3-methyl-2-oxo-1-pyridin-2-ylmethyl-  
20 imidazolidin-4-yl)-ethoxy]-phenoxy}-propionic acid



The procedures herein were utilized with 2-picolyl chloride  
hydrochloride to prepare (R)-2-methyl-2-{4-[2-(3-methyl-2-  
oxo-1-pyridin-2-ylmethyl-imidazolidin-4-yl)-ethoxy]-  
25 phenoxy}-propionic acid. HRMS (ES<sup>+</sup>) *m/z* exact mass calcd  
for C<sub>22</sub>H<sub>28</sub>N<sub>3</sub>O<sub>5</sub> 414.2029, found 414.2028. <sup>1</sup>H NMR.

**Example 151**Step A**[1-(3,4-Dimethyl-benzyl)-2,5-dioxo-imidazolidin-4-yl]-acetic acid methyl ester**


5 A solution of (2,5-dioxo-imidazolidin-4-yl)-acetic acid methyl ester (6.0 g, 34.8 mmol) in DMF (240 mL) was treated with 3,4-dimethyl benzyl bromide (5.93 g, 38.3 mmol), MgSO<sub>4</sub> (8.40 g, 60.8 mmol) and then 325 mesh K<sub>2</sub>CO<sub>3</sub> (9.15 g, 66.2  
10 mmol) at 0 °C. The resultant mixture was warmed to room temperature under N<sub>2</sub> and then heated at 50 °C for 5 h. The reaction mixture was cooled to room temperature then filtered, and then aqueous 1N HCl (140 mL) was added to the filtrate. The filtrate was extracted with EtOAc and the  
15 organic layer dried (MgSO<sub>4</sub>). The solvent was removed in vacuo to give 11.43 g crude product which was purified by flash chromatography using 3:1 hexanes:acetone to give 6.35 g of product that was triturated in Et<sub>2</sub>O/hexanes and filtered to give 2.24 g (22%) [1-(3,4-dimethyl-benzyl)-2,5-  
20 dioxo-imidazolidin-4-yl]-acetic acid methyl ester. MS (ES<sup>+</sup>) Calc'd for C<sub>15</sub>H<sub>19</sub>N<sub>2</sub>O<sub>4</sub> (M +1) 291. Found m/z 291 (100%). <sup>1</sup>H NMR.

Step B**[1-(3,4-Dimethyl-benzyl)-3-(4-methoxy-benzyl)-2,5-dioxo-imidazolidin-4-yl]-acetic acid methyl ester**

A 0 °C solution of compound [1-(3,4-dimethyl-benzyl)-2,5-dioxo-imidazolidin-4-yl]-acetic acid methyl ester (2.24 g, 7.71 mmol) in DMF (30 mL) was treated with sodium hydride (60% dispersion, 0.37 g, 9.25 mmol) and warmed to room temperature and stirred under N<sub>2</sub> for 20 minutes. The resultant mixture was cooled to 0 °C and then treated with 4-methoxybenzyl chloride (2.41 g, 15.4 mmol) and then warmed to room temperature and stirred for 16 h. The reaction was quenched with 25 mL of aqueous 1 N HCl and then worked up extractively with Et<sub>2</sub>O and water. The organic layer was dried (MgSO<sub>4</sub>) and the solvent removed *in vacuo* to give crude product that was purified by flash chromatography using a gradient of 5:1 hexanes:acetone to afford 3.23 g (100%) [1-(3,4-dimethyl-benzyl)-3-(4-methoxy-benzyl)-2,5-dioxo-imidazolidin-4-yl]-acetic acid methyl ester. MS (ES<sup>+</sup>) Calc'd for C<sub>23</sub>H<sub>26</sub>N<sub>2</sub>O<sub>5</sub> (M +1) 411. Found *m/z* 411 (100%). <sup>1</sup>H NMR.

#### Step C

**1-(3,4-Dimethyl-benzyl)-4-(2-hydroxy-ethyl)-3-(4-methoxy-benzyl)-imidazolidin-2-one**




A solution of [1-(3,4-dimethyl-benzyl)-3-(4-methoxy-benzyl)-2,5-dioxo-imidazolidin-4-yl]-acetic acid methyl ester (3.22 g, 7.84 mmol) in methanol (50 mL) was treated with aqueous 5 N NaOH (25.7 mL) and heated to reflux 1.5 h. The reaction mixture was cooled, the solvent was removed *in vacuo*. The resultant residue was acidified with aqueous 1 N HCl (200 mL) and extracted with Et<sub>2</sub>O and water. The organic layer was dried (MgSO<sub>4</sub>) and the solvent was removed *in vacuo* to afford 3.18 g (100%) of acid that was utilized without

purification. A solution of crude acid (3.18 g, assume 7.84 mmol) in THF (50 mL) was treated dropwise with 1 M solution of borane-THF complex in THF (47 mL, 47 mmol) and then stirred at room temperature under N<sub>2</sub> for 16 h. The reaction was quenched with methanol (30 mL) and stirred at room temperature for 30 minutes. The solvent was removed in vacuo to give crude product that was purified by flash chromatography using 3:1 hexanes:acetone to afford 1.89 g (65%) 1-(3,4-dimethyl-benzyl)-4-(2-hydroxy-ethyl)-3-(4-methoxy-benzyl)-imidazolidin-2-one. MS (ES<sup>+</sup>) Calc'd for C<sub>22</sub>H<sub>28</sub>N<sub>2</sub>O<sub>3</sub> (M +1) 369. Found m/z 369 (100%). <sup>1</sup>H NMR.

#### Step D


**Toluene-4-sulfonic acid 2-[1-(3,4-dimethyl-benzyl)-3-(4-methoxy-benzyl)-2-oxo-imidazolidin-4-yl]-ethyl ester**



A solution of 1-(3,4-dimethyl-benzyl)-4-(2-hydroxy-ethyl)-3-(4-methoxy-benzyl)-imidazolidin-2-one (1.89 g, 5.13 mmol), pyridine (1.42 g, 17.9 mmol) and 4-dimethyl amino pyridine (0.188 g, 1.54 mmol) in CH<sub>2</sub>Cl<sub>2</sub> was treated with p-toluenesulfonic anhydride (2.68 g, 8.21 mmol) and the reaction stirred at room temperature for under N<sub>2</sub> for 1.5 h. The reaction mixture was washed with aqueous 0.5 N HCl (100 mL), the organic layer was dried (MgSO<sub>4</sub>), and the solvent removed in vacuo to afford crude product that was purified by flash chromatography using 3:1 hexanes:actetone to afford 2.57 g (96%) toluene-4-sulfonic acid 2-[1-(3,4-dimethyl-benzyl)-3-(4-methoxy-benzyl)-2-oxo-imidazolidin-4-yl]-ethyl ester. MS (ES<sup>+</sup>) Calc'd for C<sub>29</sub>H<sub>34</sub>N<sub>2</sub>O<sub>5</sub>S (M +1) 523. Found m/z 523 (100%). <sup>1</sup>H NMR.

Step E


**2-(4-{2-[1-(3,4-Dimethyl-benzyl)-3-(4-methoxy-benzyl)-2-oxo-imidazolidin-4-yl]-ethoxy}-phenoxy)-2-methyl-propionic acid ethyl ester**



5 A mixture of toluene-4-sulfonic acid 2-[1-(3,4-dimethyl-benzyl)-3-(4-methoxy-benzyl)-2-oxo-imidazolidin-4-yl]-ethyl ester (2.54 g, 4.86 mmol), 2-(4-hydroxy-phenoxy)-2-methyl-propionic acid ethyl ester (0.99 g, 4.01 mmol) and Cs<sub>2</sub>CO<sub>3</sub>  
 10 (1.73 g, 5.31 mmol) in DMF (60 mL) was heated at 55 °C for under N<sub>2</sub> for 16 h. The reaction mixture was cooled to room temperature, quenched with aqueous 1 N HCl (25 mL), and worked up extractively with EtOAc and water. The organic layer was dried (MgSO<sub>4</sub>) and the solvent removed *in vacuo* to  
 15 afford crude product that was purified by flash chromatography using 4:1 hexanes:acetone to afford 2.51 g (99%) 2-(4-{2-[1-(3,4-dimethyl-benzyl)-3-(4-methoxy-benzyl)-2-oxo-imidazolidin-4-yl]-ethoxy}-phenoxy)-2-methyl-propionic acid ethyl ester. MS (ES<sup>+</sup>) Calc'd for C<sub>34</sub>H<sub>43</sub>N<sub>2</sub>O<sub>6</sub> (M +1) 575.  
 20 Found *m/z* 575 (100%). <sup>1</sup>H NMR.

Step F

**2-(4-{2-[1-(3,4-Dimethyl-benzyl)-2-oxo-imidazolidin-4-yl]-ethoxy}-phenoxy)-2-methyl-propionic acid ethyl ester**



25 A solution of 2-(4-{2-[1-(3,4-dimethyl-benzyl)-2-oxo-imidazolidin-4-yl]-ethoxy}-phenoxy)-2-methyl-propionic acid ethyl ester (2.51 g, 4.36 mmol) in




trifluoroacetic acid (70 mL) was stirred at room temperature under N<sub>2</sub> for 1 h. The reaction was diluted with water, and extracted with Et<sub>2</sub>O. The organic layer was dried (MgSO<sub>4</sub>) and the solvent removed *in vacuo* to afford crude product that was purified by flash chromatography using 98:2 CH<sub>2</sub>Cl<sub>2</sub>:MeOH to afford 1.20 g (60%) 2-(4-{2-[1-(3,4-dimethyl-benzyl)-2-oxo-imidazolidin-4-yl]-ethoxy}-phenoxy)-2-methyl-propionic acid ethyl ester. MS (ES<sup>+</sup>) Calc'd for C<sub>26</sub>H<sub>35</sub>N<sub>2</sub>O<sub>5</sub> (M + 1) 455. Found *m/z* 455 (100%). <sup>1</sup>H NMR.

10

Step G

**2-(4-{2-[1-(3,4-Dimethyl-benzyl)-2-oxo-imidazolidin-4-yl]-ethoxy}-phenoxy)-2-methyl-propionic acid**




15 A solution of 2-(4-{2-[1-(3,4-dimethyl-benzyl)-2-oxo-imidazolidin-4-yl]-ethoxy}-phenoxy)-2-methyl-propionic acid ethyl ester (0.087 g, 0.191 mmol) in ethanol (8 mL) was treated with aqueous 5 N NaOH (0.4 mL) and heated to reflux 1.5 h. The reaction mixture was cooled, the solvent removed *in vacuo*. The resultant residue was acidified with aqueous 1 N HCl (5 mL) and extracted with Et<sub>2</sub>O. The organic layer was dried (MgSO<sub>4</sub>) and the solvent removed *in vacuo* to afford 0.059 g (72%) 2-(4-{2-[1-(3,4-dimethyl-benzyl)-2-oxo-imidazolidin-4-yl]-ethoxy}-phenoxy)-2-methyl-propionic acid.

25 HRMS (ES<sup>+</sup>) *m/z* exact mass calcd for C<sub>24</sub>H<sub>31</sub>N<sub>2</sub>O<sub>5</sub> [M + 1] 427.2233, found 427.2232. <sup>1</sup>H NMR.

30

**Example 152****Step A**


**2-(4-{2-[1-(3,4-Dimethyl-benzyl)-3-methyl-2-oxo-  
imidazolidin-4-yl]-ethoxy}-phenoxy)-2-methyl-propionic acid  
ethyl ester**



A solution of 2-(4-{2-[1-(3,4-dimethyl-benzyl)-2-oxo-  
imidazolidin-4-yl]-ethoxy}-phenoxy)-2-methyl-propionic acid  
ethyl ester (0.128 g, 0.280 mmol) in DMF (4mL) was treated  
10 with 60% oil suspension of NaH (0.028 g, 0.70 mmol) and  
stirred at room temperature under N<sub>2</sub> for 15 minutes. The  
reaction was cooled to 0 °C and treated with iodomethane  
(0.16 g, 1.13 mmol) and then warmed to room temperature and  
stirred for 1 h. The reaction was quenched with 1 N HCl (3  
15 mL) and worked up extractively with Et<sub>2</sub>O and water. The  
organic layer was dried (MgSO<sub>4</sub>) and the solvent removed *in*  
*vacuo* to afford crude product that was purified by flash  
chromatography using 5:1 hexanes:acetone to afford 0.098 g  
(74%) 2-(4-{2-[1-(3,4-dimethyl-benzyl)-3-methyl-2-oxo-  
20 imidazolidin-4-yl]-ethoxy}-phenoxy)-2-methyl-propionic acid  
ethyl ester. MS (ES<sup>+</sup>) Calc'd for C<sub>27</sub>H<sub>37</sub>N<sub>2</sub>O<sub>5</sub> (M +1) 469.  
Found *m/z* 469 (100%). <sup>1</sup>H NMR.

**Step B**


**2-(4-{2-[1-(3,4-Dimethyl-benzyl)-3-methyl-2-oxo-  
imidazolidin-4-yl]-ethoxy}-phenoxy)-2-methyl-propionic acid**



A solution of 2-(4-{2-[1-(3,4-dimethyl-benzyl)-3-methyl-2-oxo-imidazolidin-4-yl]-ethoxy}-phenoxy)-2-methyl-propionic acid ethyl ester (0.098 g, 0.209 mmol) in ethanol (8 mL) was treated with aqueous 5 N NaOH (0.5 mL) and heated to reflux 1 h. The reaction mixture was cooled, the solvent removed *in vacuo*. The resultant residue was acidified with aqueous 1 N HCl (5 mL) and extracted with Et<sub>2</sub>O. The organic layer was dried (MgSO<sub>4</sub>) and the solvent removed *in vacuo* to afford 0.050 g (54%) 2-(4-{2-[1-(3,4-dimethyl-benzyl)-3-methyl-2-oxo-imidazolidin-4-yl]-ethoxy}-phenoxy)-2-methyl-propionic acid. HRMS (ES<sup>+</sup>) *m/z* exact mass calcd for C<sub>25</sub>H<sub>33</sub>N<sub>2</sub>O<sub>5</sub> [M + 1] 441.2389, found 441.2390. <sup>1</sup>H NMR.

#### Example 153


2-(4-{2-[1-(3,4-Dimethyl-benzyl)-3-ethyl-2-oxo-imidazolidin-4-yl]-ethoxy}-phenoxy)-2-methyl-propionic acid



The procedures herein utilized with ethyl iodide to afford 2-(4-{2-[1-(3,4-dimethyl-benzyl)-3-ethyl-2-oxo-imidazolidin-4-yl]-ethoxy}-phenoxy)-2-methyl-propionic acid. MS (ES<sup>+</sup>) Calc'd for C<sub>26</sub>H<sub>35</sub>N<sub>2</sub>O<sub>5</sub> (M + 1) 455. Found *m/z* 455 (100%). <sup>1</sup>H NMR.

#### Example 154

2-(4-{2-[1-(3,4-Dimethyl-benzyl)-2-oxo-3-propyl-imidazolidin-4-yl]-ethoxy}-phenoxy)-2-methyl-propionic acid




The procedures herein utilized with propyl iodide to afford 2-(4-{2-[1-(3,4-dimethyl-benzyl)-2-oxo-3-propyl-

imidazolidin-4-yl]-ethoxy}-phenoxy)-2-methyl-propionic acid.  
MS (ES<sup>+</sup>) Calc'd for C<sub>27</sub>H<sub>37</sub>N<sub>2</sub>O<sub>5</sub> (M + 1) 469. Found m/z 469  
(100%). <sup>1</sup>H NMR.

**Example 155**


5      **(R)-2-Methyl-2-(4-{2-[1-(4-methyl-benzyl)-2-oxo-3-propyl-imidazolidin-4-yl]-ethoxy}-phenoxy)-propionic acid**



The procedures herein were utilized to prepare 2-methyl-2-(4-{2-[1-(4-methyl-benzyl)-2-oxo-3-propyl-imidazolidin-4-yl]-ethoxy}-phenoxy)-propionic acid. HRMS (ES<sup>+</sup>) m/z exact  
10      mass calcd for C<sub>26</sub>H<sub>35</sub>N<sub>2</sub>O<sub>5</sub> [M + 1] 455.2546, found 455.2565.  
<sup>1</sup>H NMR.

**Example 156**


15      **(R)-2-Methyl-2-(4-{2-[3-methyl-1-(4-methyl-benzyl)-2-oxo-imidazolidin-4-yl]-ethoxy}-phenoxy)-propionic acid**



The procedures herein were utilized to prepare 2-methyl-2-(4-{2-[3-methyl-1-(4-methyl-benzyl)-2-oxo-imidazolidin-4-yl]-ethoxy}-phenoxy)-propionic acid. HRMS  
20      (ES<sup>+</sup>) m/z exact mass calcd for C<sub>24</sub>H<sub>31</sub>N<sub>2</sub>O<sub>5</sub> [M + 1] 427.2233,  
found 427.2233. <sup>1</sup>H NMR.

**Example 157**


25      **2-(4-{2-[1-(4-tert-Butyl-benzyl)-2-oxo-imidazolidin-4-yl]-ethoxy}-phenoxy)-2-methyl-propionic acid**



The procedures herein were utilized to prepare 2-(4-{2-[1-(4-*tert*-butyl-benzyl)-2-oxo-imidazolidin-4-yl]-ethoxy}-phenoxy)-2-methyl-propionic acid. HRMS (ES<sup>+</sup>) *m/z* exact mass  
5 calcd for C<sub>26</sub>H<sub>35</sub>N<sub>2</sub>O<sub>5</sub> [M + 1] 455.2546, found 455.2538. <sup>1</sup>H NMR.

#### Example 158


2-(4-{2-[1-(4-*tert*-Butyl-benzyl)-3-methyl-2-oxo-  
10 imidazolidin-4-yl]-ethoxy}-phenoxy)-2-methyl-propionic acid



The procedures herein were utilized to prepare 2-(4-{2-[1-(4-*tert*-Butyl-benzyl)-3-methyl-2-oxo-imidazolidin-4-yl]-ethoxy}-phenoxy)-2-methyl-propionic acid. HRMS (ES<sup>+</sup>) *m/z*  
15 exact mass calcd for C<sub>27</sub>H<sub>37</sub>N<sub>2</sub>O<sub>5</sub> [M + 1] 469.2702, found 469.2690. <sup>1</sup>H NMR.

#### Example 159


2-(4-{2-[1-(4-*tert*-Butyl-benzyl)-3-ethyl-2-oxo-imidazolidin-  
20 4-yl]-ethoxy}-phenoxy)-2-methyl-propionic acid



The procedures herein were utilized to prepare 2-(4-{2-[1-(4-*tert*-butyl-benzyl)-3-ethyl-2-oxo-imidazolidin-4-yl]-ethoxy}-phenoxy)-2-methyl-propionic acid. MS (ES<sup>+</sup>) Calc'd  
25 for C<sub>28</sub>H<sub>39</sub>N<sub>2</sub>O<sub>5</sub> (M + 1) 483. Found *m/z* 483 (100%). <sup>1</sup>H NMR.

**Example 160**


**2-(4-{2-[1-(4-*tert*-Butyl-benzyl)-2-oxo-3-propyl-imidazolidin-4-yl]-ethoxy}-phenoxy)-2-methyl-propionic acid**



The procedures herein were utilized to prepare 2-(4-{2-[1-(4-*tert*-butyl-benzyl)-2-oxo-3-propyl-imidazolidin-4-yl]-ethoxy}-phenoxy)-2-methyl-propionic acid. MS (ES<sup>+</sup>) Calc'd for C<sub>29</sub>H<sub>41</sub>N<sub>2</sub>O<sub>5</sub> (M +1) 497. Found *m/z* 497 (100%). <sup>1</sup>H NMR.

**Example 161**


**2-Methyl-2-{4-[2-(1-naphthalen-2-ylmethyl-2-oxo-imidazolidin-4-yl)-ethoxy]-phenoxy}-propionic acid**



The procedures herein were utilized to prepare 2-methyl-2-{4-[2-(1-naphthalen-2-ylmethyl-2-oxo-imidazolidin-4-yl)-ethoxy]-phenoxy}-propionic acid. MS (ES<sup>+</sup>) Calc'd for C<sub>26</sub>H<sub>29</sub>N<sub>2</sub>O<sub>5</sub> (M +1) 449. Found *m/z* 449 (100%). <sup>1</sup>H NMR.

**Example 162**

**2-Methyl-2-{4-[2-(3-methyl-1-naphthalen-2-ylmethyl-2-oxo-imidazolidin-4-yl)-ethoxy]-phenoxy}-propionic acid**




The procedures herein were utilized to prepare 2-methyl-2-{4-[2-(3-methyl-1-naphthalen-2-ylmethyl-2-oxo-imidazolidin-

4-yl)-ethoxy]-phenoxy}-propionic acid. MS (ES<sup>+</sup>) Calc'd for C<sub>27</sub>H<sub>31</sub>N<sub>2</sub>O<sub>5</sub> (M + 1) 463. Found *m/z* 463 (100%). <sup>1</sup>H NMR.

#### Example 163


5     **2-Methyl-2-{4-[2-(1-naphthalen-2-ylmethyl-2-oxo-3-propyl-imidazolidin-4-yl)-ethoxy]-phenoxy}-propionic acid**



The procedures herein were utilized to prepare 2-methyl-2-{4-[2-(1-naphthalen-2-ylmethyl-2-oxo-3-propyl-imidazolidin-4-yl)-ethoxy]-phenoxy}-propionic acid. MS (ES<sup>+</sup>) Calc'd for C<sub>29</sub>H<sub>35</sub>N<sub>2</sub>O<sub>5</sub> (M + 1) 491. Found *m/z* 491 (100%). <sup>1</sup>H NMR.

#### Example 164


15     **2,2-Dimethyl-3-(4-{2-[1-(4-methyl-benzyl)-2-oxo-3-propyl-imidazolidin-4-yl]-ethoxy}-phenyl)-propionic acid**



The procedures herein were utilized to prepare 2,2-dimethyl-3-(4-{2-[1-(4-methyl-benzyl)-2-oxo-3-propyl-imidazolidin-4-yl]-ethoxy}-phenyl)-propionic acid. HRMS (ES<sup>+</sup>) *m/z* exact mass calcd for C<sub>27</sub>H<sub>37</sub>N<sub>2</sub>O<sub>4</sub> [M + 1] 453.2753, found 453.2754. <sup>1</sup>H NMR.

#### Example 165


25     **2-Methyl-2-(3-{2-[1-(4-methyl-benzyl)-2-oxo-3-propyl-imidazolidin-4-yl]-ethoxy}-phenoxy)-propionic acid**



The procedures herein were utilized to prepare 2-methyl-2-(3-{2-[1-(4-methyl-benzyl)-2-oxo-3-propyl-imidazolidin-4-yl]-ethoxy}-phenoxy)-propionic acid. HRMS (ES<sup>+</sup>) *m/z* exact mass calcd for C<sub>26</sub>H<sub>34</sub>N<sub>2</sub>O<sub>5</sub> [M + 1] 455.2546, found 455.2549. <sup>1</sup>H NMR.

#### Example 166


2-Methoxy-2-methyl-3-(4-{2-[1-(4-methyl-benzyl)-2-oxo-3-propyl-imidazolidin-4-yl]-ethoxy}-phenyl)-propionic acid



The procedures herein were utilized to prepare 2-methoxy-2-methyl-3-(4-{2-[1-(4-methyl-benzyl)-2-oxo-3-propyl-imidazolidin-4-yl]-ethoxy}-phenyl)-propionic acid. HRMS (ES<sup>+</sup>) *m/z* exact mass calcd for C<sub>27</sub>H<sub>36</sub>N<sub>2</sub>O<sub>5</sub> [M + 1] 469.2702, found 469.2709. <sup>1</sup>H NMR.

#### Example 167

2-(4-{2-[1-(4-Methyl-benzyl)-2-oxo-3-propyl-imidazolidin-4-yl]-ethoxy}-benzyl)-butyric acid




The procedures herein were utilized to prepare 2-(4-{2-[1-(4-methyl-benzyl)-2-oxo-3-propyl-imidazolidin-4-yl]-ethoxy}-benzyl)-butyric acid. HRMS (ES<sup>+</sup>) *m/z* exact mass calcd for C<sub>27</sub>H<sub>36</sub>N<sub>2</sub>O<sub>4</sub> [M + 1] 453.2753, found 453.2749. <sup>1</sup>H NMR.



**Example 168**

**4-{2-[1-(4-Methyl-benzyl)-2-oxo-3-propyl-imidazolidin-4-yl]-ethoxy}-benzoic acid**

5




The procedures herein were utilized to prepare 4-{2-[1-(4-methyl-benzyl)-2-oxo-3-propyl-imidazolidin-4-yl]-ethoxy}-benzoic acid. HRMS (ES<sup>+</sup>) *m/z* exact mass calcd for C<sub>23</sub>H<sub>29</sub>N<sub>2</sub>O<sub>4</sub> [M + 1] 397.2127, found 397.2112. <sup>1</sup>H NMR.

10

**Example 169**

**(4-{2-[1-(4-Methyl-benzyl)-2-oxo-3-propyl-imidazolidin-4-yl]-ethoxy}-phenyl)-acetic acid**

15




The procedures herein were utilized to prepare (4-{2-[1-(4-methyl-benzyl)-2-oxo-3-propyl-imidazolidin-4-yl]-ethoxy}-phenyl)-acetic acid. HRMS (ES<sup>+</sup>) *m/z* exact mass calcd for C<sub>24</sub>H<sub>31</sub>N<sub>2</sub>O<sub>4</sub> [M + 1] 411.2284, found 411.2269. <sup>1</sup>H NMR.

20

**Example 170**

**3-(4-{2-[1-(4-Methyl-benzyl)-2-oxo-3-propyl-imidazolidin-4-yl]-ethoxy}-phenyl)-propionic acid**




The procedures herein were utilized to prepare 3-(4-{2-[1-(4-methyl-benzyl)-2-oxo-3-propyl-imidazolidin-4-yl]-ethoxy}-

25

phenyl)-propionic acid. HRMS (ES<sup>+</sup>) *m/z* exact mass calcd for C<sub>25</sub>H<sub>33</sub>N<sub>2</sub>O<sub>4</sub> [M + 1] 425.2440, found 425.2427. <sup>1</sup>H NMR.

#### Example 171

- 5    **3-{2-[1-(4-Methyl-benzyl)-2-oxo-3-propyl-imidazolidin-4-yl]-ethoxy}-benzoic acid**




The procedures herein were utilized to prepare 3-{2-[1-(4-methyl-benzyl)-2-oxo-3-propyl-imidazolidin-4-yl]-ethoxy}-  
10 benzoic acid. HRMS (ES<sup>+</sup>) *m/z* exact mass calcd for C<sub>23</sub>H<sub>29</sub>N<sub>2</sub>O<sub>4</sub> [M + 1] 397.2127, found 397.2118. <sup>1</sup>H NMR.

#### Example 172

##### Step A

- 15    **1-(4-Hydroxy-phenyl)-cyclopentanecarboxylic acid ethyl ester**




A solution of 1-(*p*-methoxyphenyl)-cyclopentanecarboxylic acid 093909 (5.0 g, 22.7 mmol) in ethanol (50 mL) was treated with conc. H<sub>2</sub>SO<sub>4</sub> (4.40 g, 44.9 mmol) and heated to  
20 reflux under N<sub>2</sub> for 12 h. The reaction was cooled and the solvent removed *in vacuo* to give a residue that was extracted with EtOAc and water. The organic layer was dried (MgSO<sub>4</sub>) and the solvent removed *in vacuo* to 5.93 g crude ester (100%) that was utilized without purification. The  
25 ester (5.93 g, assume 22.7 mmol) was dissolved in dry CH<sub>2</sub>Cl<sub>2</sub> (75 mL), cooled to - 78 °C and then treated dropwise with BBr<sub>3</sub> (11.4 g, 45.6 mmol). The reaction was warmed to 0 °C for 15 minutes and then quenched with methanol then water. The reaction mixture was extracted with CH<sub>2</sub>Cl<sub>2</sub> and water,

the organic layer was dried ( $\text{MgSO}_4$ ), and the solvent removed in *vacuo* to give crude product that was purified by flash chromatography using 6:1 hexanes:EtOAc to afford 4.34 g (79%) 1-(4-hydroxy-phenyl)-cyclopentanecarboxylic acid ethyl ester. MS ( $\text{ES}^+$ ) Calc'd for  $\text{C}_{14}\text{H}_{19}\text{O}_3$  ( $M + 1$ ) 235. Found  $m/z$  235 (100%).  $^1\text{H}$  NMR.

#### Step B

1-(4-{2-[1-(4-Methyl-benzyl)-2-oxo-3-propyl-imidazolidin-4-yl]-ethoxy}-phenyl)-cyclopentanecarboxylic acid



10

A mixture of 1-(4-hydroxy-phenyl)-cyclopentanecarboxylic acid ethyl ester (0.042 g, 0.179 mmol), toluene-4-sulfonic acid 2-[1-(4-methyl-benzyl)-2-oxo-3-propyl-imidazolidin-4-yl]-ethyl ester (0.085 g, 0.197 mmol) and 325 mesh  $\text{K}_2\text{CO}_3$  (0.050 g, 0.362 mmol) in ethanol (5 mL) was heated at reflux for under  $\text{N}_2$  for 16 h. The reaction was then treated with aqueous 5 N NaOH (0.5 mL) and heated to reflux 1.5 h. The reaction mixture was cooled, the solvent removed in *vacuo*. The resultant residue was acidified with aqueous 1 N HCl (10 mL) and extracted with  $\text{Et}_2\text{O}$ . The organic layer was dried ( $\text{MgSO}_4$ ) and the solvent removed in *vacuo* to afford 0.058 g crude acid that was purified by preparative HPLC to afford 0.033 g (40%) 1-(4-{2-[1-(4-methyl-benzyl)-2-oxo-3-propyl-imidazolidin-4-yl]-ethoxy}-phenyl)-cyclopentanecarboxylic acid. MS ( $\text{ES}^+$ ) Calc'd for  $\text{C}_{28}\text{H}_{37}\text{N}_2\text{O}_4$  ( $M + 1$ ) 465. Found  $m/z$  465 (100%).  $^1\text{H}$  NMR.

15


20

25

30

#### Example 173


2-Methyl-2-(4-{2-[1-(4-methyl-benzyl)-2-oxo-3-propyl-imidazolidin-4-yl]-ethoxy}-phenoxy)-3-phenyl-propionic acid



The procedures herein were utilized to prepare 2-methyl-2-(4-{2-[1-(4-methyl-benzyl)-2-oxo-3-propyl-imidazolidin-4-yl]-ethoxy}-phenoxy)-3-phenyl-propionic acid. HRMS (ES<sup>+</sup>)  $m/z$  exact mass calcd for C<sub>32</sub>H<sub>38</sub>N<sub>2</sub>O<sub>5</sub> [M + 1] 531.2859, found 531.2855. <sup>1</sup>H NMR.

#### Example 174

2-Methoxy-3-(4-{2-[1-(4-methyl-benzyl)-2-oxo-3-propyl-imidazolidin-4-yl]-ethoxy}-phenyl)-propionic acid




The procedures herein were utilized to prepare 2-Methoxy-3-(4-{2-[1-(4-methyl-benzyl)-2-oxo-3-propyl-imidazolidin-4-yl]-ethoxy}-phenyl)-propionic acid. MS (ES<sup>+</sup>) Calc'd for C<sub>26</sub>H<sub>35</sub>N<sub>2</sub>O<sub>5</sub> (M + 1) 455. Found  $m/z$  455 (100%). <sup>1</sup>H NMR.

#### Example 175

##### Step A

5-[3-(4-Methoxy-phenyl)-propyl]-3-(4-methyl-benzyl)-imidazolidine-2,4-dione




A solution of 5-[3-(4-methoxy-phenyl)-propyl]-imidazolidine-2,4-dione (2.00 g, 8.05 mmol) in DMF (80 mL) was treated with *p*-methyl benzyl bromide (1.64 g, 8.86 mmol), MgSO<sub>4</sub> (1.94 g, 16.1 mmol) and then 325 mesh K<sub>2</sub>CO<sub>3</sub> (2.12 g, 15.3

mmol). The resultant mixture was stirred under N<sub>2</sub> in an ice bath for 30 minutes and then warmed to room temperature for 16 h. The reaction mixture was then filtered, and then aqueous 1N HCl (40 mL) was added to the filtrate. The filtrate was extracted with EtOAc and the organic layer dried (MgSO<sub>4</sub>). The solvent was removed *in vacuo* to give a crude product which was purified by flash chromatography using 3:1 hexanes:EtOAc to afford 2.41 g (85%) 5-[3-(4-methoxy-phenyl)-propyl]-imidazolidine-2,4-dione. MS (ES<sup>+</sup>) Calc'd for C<sub>21</sub>H<sub>25</sub>N<sub>2</sub>O<sub>3</sub> (M +1) 353. Found *m/z* 353 (100%). <sup>1</sup>H NMR.

#### Step B

#### 5-[3-(4-Methoxy-phenyl)-propyl]-3-(4-methyl-benzyl)-1-propyl-imidazolidine-2,4-dione



15


A 0 °C solution of compound 5-[3-(4-methoxy-phenyl)-propyl]-imidazolidine-2,4-dione (2.41 g, 6.84 mmol) in DMF (35 mL) was treated with sodium hydride (60% dispersion, 0.23 g, 5.75 mmol) and warmed to room temperature and stirred under N<sub>2</sub> for 15 minutes. The resultant mixture was cooled to 0 °C and then treated with 1-propyl iodide (0.96 g, 5.64 mmol) and then warmed to room temperature and stirred for 1 h. The reaction was quenched with 1 N HCl (20 mL) and then worked up extractively with Et<sub>2</sub>O and water. The organic layer was dried (MgSO<sub>4</sub>) and the solvent removed *in vacuo* to give crude product that was purified by flash chromatography using a 6:1 hexanes:EtOAc to afford 1.58 g (71%) 5-[3-(4-methoxy-phenyl)-propyl]-3-(4-methyl-benzyl)-1-propyl-imidazolidine-2,4-dione. MS (ES<sup>+</sup>) Calc'd for C<sub>24</sub>H<sub>31</sub>N<sub>2</sub>O<sub>3</sub> (M

25

+1) 395. Found  $m/z$  395 (100%).  $^1\text{H}$  NMR.

Step C


5           **4-[3-(4-Methoxy-phenyl)-propyl]-1-(4-methyl-benzyl)-3-propyl-imidazolidin-2-one**



A solution of 5-[3-(4-methoxy-phenyl)-propyl]-3-(4-methyl-benzyl)-1-propyl-imidazolidine-2,4-dione (1.26 g, 3.19 mmol) in THF (20 mL) was treated dropwise with 1 M solution of borane-THF complex in THF (16 mL, 16 mmol) and then stirred at room temperature under  $\text{N}_2$  for 22 h. The reaction was quenched with methanol (20 mL) and stirred at room temperature for 2 hr. The solvent was removed *in vacuo* to give 1.25 g (100%) crude 4-[3-(4-methoxy-phenyl)-propyl]-1-(4-methyl-benzyl)-3-propyl-imidazolidin-2-one that was carried on without purification. MS ( $\text{ES}^+$ ) Calc'd for  $\text{C}_{24}\text{H}_{33}\text{N}_2\text{O}_2$  ( $\text{M} + 1$ ) 381. Found  $m/z$  381 (100%).  $^1\text{H}$  NMR.

Step D

20           **4-[3-(4-Hydroxy-phenyl)-propyl]-1-(4-methyl-benzyl)-3-propyl-imidazolidin-2-one**




A solution of 4-[3-(4-methoxy-phenyl)-propyl]-1-(4-methyl-benzyl)-3-propyl-imidazolidin-2-one (1.22 g, assume 3.21 mmol) was dissolved in dry  $\text{CH}_2\text{Cl}_2$  (25 mL), cooled to 0 °C and then treated dropwise with  $\text{BBr}_3$  (2.41 g, 9.63 mmol). The reaction was stirred at 0 °C for 1 h. The reaction mixture

was diluted with Et<sub>2</sub>O and water, the organic layer was dried (MgSO<sub>4</sub>), and the solvent removed *in vacuo* to give crude product that was purified by flash chromatography using 1:1 hexanes:EtOAc to afford 0.687 g (68%) 4-[3-(4-hydroxy-phenyl)-propyl]-1-(4-methyl-benzyl)-3-propyl-imidazolidin-2-one. MS (ES<sup>+</sup>) Calc'd for C<sub>23</sub>H<sub>31</sub>N<sub>2</sub>O<sub>2</sub> (M +1) 367. Found *m/z* 367 (100%). <sup>1</sup>H NMR.

#### Step E


2-Methyl-2-(4-{3-[1-(4-methyl-benzyl)-2-oxo-3-propyl-imidazolidin-4-yl]-propyl}-phenoxy)-propionic acid ethyl ester



A mixture of 4-[3-(4-hydroxy-phenyl)-propyl]-1-(4-methyl-benzyl)-3-propyl-imidazolidin-2-one (0.687 g, 1.87 mmol), ethyl 2-bromoisobutyrate (2.55 g, 13.1 mmol), MgSO<sub>4</sub> (0.22 g, 1.83 mmol) and 325 mesh K<sub>2</sub>CO<sub>3</sub> (0.77 g, 5.57 mmol) in ethanol (60 mL) was heated at 70 °C for under N<sub>2</sub> for 16 h. The reaction mixture was cooled and the solvent removed *in vacuo* to give a residue that was acidified with 1 N HCl (20 mL). The reaction was diluted with EtOAc and washed with water. The organic layer was dried (MgSO<sub>4</sub>) and the solvent removed *in vacuo* to give crude product that was purified by flash chromatography using 5:1 hexanes:acetone to afford 0.648 g (72%) 2-methyl-2-(4-{3-[1-(4-methyl-benzyl)-2-oxo-3-propyl-imidazolidin-4-yl]-propyl}-phenoxy)-propionic acid ethyl ester. MS (ES<sup>+</sup>) Calc'd for C<sub>29</sub>H<sub>41</sub>N<sub>2</sub>O<sub>4</sub> (M +1) 481. Found *m/z* 481 (100%). <sup>1</sup>H NMR.

#### Step F


2-Methyl-2-(4-{3-[1-(4-methyl-benzyl)-2-oxo-3-propyl-imidazolidin-4-yl]-propyl}-phenoxy)-propionic acid



A solution of 2-methyl-2-(4-{3-[1-(4-methyl-benzyl)-2-oxo-3-propyl-imidazolidin-4-yl]-propyl}-phenoxy)-propionic acid ethyl ester (0.191 g, 0.397 mmol) in ethanol (10 mL) was  
 5 treated with aqueous 5 N NaOH (0.8 mL) and heated to reflux 1.5 h. The reaction mixture was cooled, the solvent removed *in vacuo*. The resultant residue was acidified with aqueous 1 N HCl (10 mL) and extracted with Et<sub>2</sub>O. The organic layer was dried (MgSO<sub>4</sub>) and the solvent removed *in vacuo* to afford  
 10 0.178g (99%) 2-methyl-2-(4-{3-[1-(4-methyl-benzyl)-2-oxo-3-propyl-imidazolidin-4-yl]-propyl}-phenoxy)-propionic acid. HRMS (ES<sup>+</sup>) *m/z* exact mass calcd for C<sub>27</sub>H<sub>37</sub>N<sub>2</sub>O<sub>4</sub> [M + 1] 453.2753, found 453.2751. <sup>1</sup>H NMR.

#### Example 176

15 **(R)-2-Methyl-2-(4-{3-[1-(4-methyl-benzyl)-2-oxo-3-propyl-imidazolidin-4-yl]-propyl}-phenoxy)-propionic acid**




The procedures herein were utilized to prepare (R)-2-methyl-2-(4-{3-[1-(4-methyl-benzyl)-2-oxo-3-propyl-imidazolidin-4-yl]-propyl}-phenoxy)-propionic acid. HRMS (ES<sup>+</sup>) *m/z* exact  
 20 mass calcd for C<sub>27</sub>H<sub>37</sub>N<sub>2</sub>O<sub>4</sub> [M + 1] 453.2753, found 453.2757. <sup>1</sup>H NMR.

#### Example 177

##### Step A

25 **2-(4-{3-[1-(3,4-Dimethyl-benzyl)-2,5-dioxo-imidazolidin-4-yl]-propyl}-phenoxy)-2-methyl-propionic acid tert-butyl ester**






Compound 2-(4-[3-(2,5-dioxo-imidazolidin-4-yl)-propyl]-phenoxy)-2-methyl-propionic acid tert-butyl ester (0.29 g, 0.770 mmol) in DMF (10 mL) was treated with 3,4-dimethyl benzyl chloride (0.131 g, 0.847 mmol), MgSO<sub>4</sub> (0.185 g, 1.54 mmol) and then 325 mesh K<sub>2</sub>CO<sub>3</sub> (0.213 g, 1.54 mmol). The resultant mixture was heated to 50 °C under N<sub>2</sub> for 4 h. The reaction mixture was cooled and quenched with 1N HCl (8 mL) and then extracted with EtOAc and water. The organic layer was dried (MgSO<sub>4</sub>) and the solvent was removed in vacuo to give a crude product which was purified by flash chromatography using a gradient of 4:1 then 1:1 hexanes:EtOAc to afford 0.202 g (53%) 2-(4-{3-[1-(3,4-dimethyl-benzyl)-2,5-dioxo-imidazolidin-4-yl]-propyl}-phenoxy)-2-methyl-propionic acid tert-butyl ester. MS (ES<sup>+</sup>) Calc'd for C<sub>29</sub>H<sub>39</sub>N<sub>2</sub>O<sub>5</sub> (M +1) 495. Found m/z 495 (100%). <sup>1</sup>H NMR.

#### Step B

2-(4-{3-[1-(3,4-Dimethyl-benzyl)-2-oxo-imidazolidin-4-yl]-propyl}-phenoxy)-2-methyl-propionic acid tert-butyl ester




A solution of 2-(4-{3-[1-(3,4-dimethyl-benzyl)-2,5-dioxo-imidazolidin-4-yl]-propyl}-phenoxy)-2-methyl-propionic acid tert-butyl ester (0.113 g, 0.228 mmol) in THF (6 mL) was treated dropwise with 1 M solution of borane-THF complex in THF (4.4 mL, 4.4 mmol) and then stirred at room temperature under N<sub>2</sub> for 32 h. The reaction was quenched with methanol (6 mL) and stirred at room temperature for 2

hr. The solvent was removed *in vacuo* to give crude product that was purified by flash chromatography using 97:3 CH<sub>2</sub>Cl<sub>2</sub>:MeOH to afford 0.051 g (47%) 2-(4-{3-[1-(3,4-dimethyl-benzyl)-2-oxo-imidazolidin-4-yl]-propyl}-phenoxy)-2-methyl-propionic acid *tert*-butyl ester. MS (ES<sup>+</sup>) Calc'd for C<sub>29</sub>H<sub>41</sub>N<sub>2</sub>O<sub>4</sub> (M +1) 481. Found *m/z* 481 (100%). <sup>1</sup>H NMR.

### Step C


2-(4-{3-[1-(3,4-Dimethyl-benzyl)-2-oxo-imidazolidin-4-yl]-propyl}-phenoxy)-2-methyl-propionic acid



A solution of 2-(4-{3-[1-(3,4-dimethyl-benzyl)-2-oxo-imidazolidin-4-yl]-propyl}-phenoxy)-2-methyl-propionic acid *tert*-butyl ester (0.051 g, 0.106 mmol) in CH<sub>2</sub>Cl<sub>2</sub> (3 mL) was treated with trifluoroacetic acid (0.5 mL) and stirred at room temperature under N<sub>2</sub> for 5 h. The solvent removed *in vacuo* to give crude acid that was purified by flash chromatography using 94:6 CH<sub>2</sub>Cl<sub>2</sub>:MeOH to afford 0.011 g (24%) 2-(4-{3-[1-(3,4-dimethyl-benzyl)-2-oxo-imidazolidin-4-yl]-propyl}-phenoxy)-2-methyl-propionic acid. MS (ES<sup>+</sup>) Calc'd for C<sub>25</sub>H<sub>33</sub>N<sub>2</sub>O<sub>4</sub> (M +1) 425. Found *m/z* 425 (100%). <sup>1</sup>H NMR.

### Example 178

2-(4-{3-[1-(4-*tert*-Butyl-benzyl)-2-oxo-imidazolidin-4-yl]-propyl}-phenoxy)-2-methyl-propionic acid




The procedures herein were utilized to prepare 2-(4-{3-[1-

(4-*tert*-butyl-benzyl)-2-oxo-imidazolidin-4-yl]-propyl}-phenoxy)-2-methyl-propionic acid. MS (ES<sup>+</sup>) Calc'd for C<sub>27</sub>H<sub>37</sub>N<sub>2</sub>O<sub>4</sub> (M +1) 453. Found *m/z* 453 (100%). <sup>1</sup>H NMR.

5

**Example 179**

**2-(4-{3-[1-(4-Chloro-benzyl)-2-oxo-imidazolidin-4-yl]-propyl}-phenoxy)-2-methyl-propionic acid**




10 The procedures herein were utilized to prepare 2-(4-{3-[1-(4-chloro-benzyl)-2-oxo-imidazolidin-4-yl]-propyl}-phenoxy)-2-methyl-propionic acid. MS (ES<sup>+</sup>) Calc'd for C<sub>23</sub>H<sub>28</sub>ClN<sub>2</sub>O<sub>4</sub> (M+1) 431. Found *m/z* 431 (100%).

15

**Example 180**


**2-(4-{3-[1-(4-Chloro-benzyl)-3-methyl-2-oxo-imidazolidin-4-yl]-propyl}-phenoxy)-2-methyl-propionic acid**



20 The procedures herein were utilized to prepare 2-(4-{3-[1-(4-chloro-benzyl)-3-methyl-2-oxo-imidazolidin-4-yl]-propyl}-phenoxy)-2-methyl-propionic acid. MS (ES<sup>+</sup>) Calc'd for C<sub>24</sub>H<sub>30</sub>ClN<sub>2</sub>O<sub>4</sub> (M+1) 445. Found *m/z* 445 (100%).

**Example 181**

25 **2-Methyl-2-{4-[3-(3-methyl-1-naphthalen-2-ylmethyl-2-oxo-imidazolidin-4-yl)-propyl]-phenoxy}-propionic acid**




The procedures herein were utilized to prepare 2-methyl-2-{4-[3-(3-methyl-1-naphthalen-2-ylmethyl-2-oxo-imidazolidin-4-yl)-propyl]-phenoxy}-propionic acid. MS (ES<sup>+</sup>) Calc'd for C<sub>28</sub>H<sub>33</sub>N<sub>2</sub>O<sub>4</sub> (M+1) 461. Found m/z 461 (100%).

5

**Example 182**

2-{4-[3-(3-Ethyl-1-naphthalen-2-ylmethyl-2-oxo-imidazolidin-4-yl)-propyl]-phenoxy}-2-methyl-propionic acid




10 The procedures herein were utilized to prepare 2-{4-[3-(3-ethyl-1-naphthalen-2-ylmethyl-2-oxo-imidazolidin-4-yl)-propyl]-phenoxy}-2-methyl-propionic acid. MS (ES<sup>+</sup>) Calc'd for C<sub>29</sub>H<sub>35</sub>N<sub>2</sub>O<sub>4</sub> (M+1) 475. Found m/z 475 (100%).

15

**Example 183**

2-Methyl-2-{4-[3-(1-naphthalen-2-ylmethyl-2-oxo-3-propyl-imidazolidin-4-yl)-propyl]-phenoxy}-propionic acid




The procedures herein were utilized to prepare 2-methyl-2-{4-[3-(1-naphthalen-2-ylmethyl-2-oxo-3-propyl-imidazolidin-4-yl)-propyl]-phenoxy}-propionic acid. MS (ES<sup>+</sup>) Calc'd for C<sub>30</sub>H<sub>37</sub>N<sub>2</sub>O<sub>4</sub> (M+1) 489. Found m/z 489 (100%).

20

**Example 184**


25 2-(4-{3-[1-(4-tert-butyl-benzyl)-3-methyl-2-oxo-imidazolidin-4-yl]-propyl}-phenoxy)-2-ethyl-butyric acid



A mixture of 1-(4-*tert*-butyl-benzyl)-4-[3-(4-hydroxy-phenyl)-propyl]-3-methyl-imidazolidin-2-one (0.077 g, 0.202 mmol), potassium *tert*-butoxide (0.091 g, 0.745 mmol) and  
 5 ethyl  $\alpha$ -bromodiethylacetate (0.361 g, 1.61 mmol) in 2-methyl-2-propanol (8 mL) was heated to 100 °C under N<sub>2</sub> for 16 h. Aqueous 5 N NaOH (1.5 mL) was added to the reaction mixture and it was heated at reflux an additional 1 h. The reaction mixture was cooled and quenched with 1 N HCl. The  
 10 mixture was then diluted with water and extracted with CH<sub>2</sub>Cl<sub>2</sub>. The organic layer was dried (Na<sub>2</sub>SO<sub>4</sub>) and the solvent removed *in vacuo* to afford 0.288 g of crude product that was purified by preparative HPLC to give 0.014 g (14%) 2-(4-{3-[1-(4-*tert*-butyl-benzyl)-3-methyl-2-oxo-imidazolidin-4-yl]-propyl}-phenoxy)-2-ethyl-butyric acid. <sup>1</sup>H NMR. HRMS (ES<sup>+</sup>) *m/z* exact mass calcd for C<sub>30</sub>H<sub>43</sub>N<sub>2</sub>O<sub>4</sub> [M + 1] 495.3223, found 495.3252.

#### Example 185

20 1-(4-{3-[1-(4-*tert*-butyl-benzyl)-3-methyl-2-oxo-imidazolidin-4-yl]-propyl}-phenoxy)-cyclohexanecarboxylic acid




A mixture of 1-(4-*tert*-butyl-benzyl)-4-[3-(4-hydroxy-phenyl)-propyl]-3-methyl-imidazolidin-2-one (0.074 g, 0.194 mmol) and potassium *tert*-butoxide (0.044 g, 0.392 mmol) in

2-methyl-2-propanol (8 mL) was heated to 60 °C under N<sub>2</sub> and then and methyl 1-bromocyclohexanecarboxylate (0.274 g, 1.16 mmol) was added dropwise to the to 60 °C solution. Additional potassium *tert*-butoxide (0.044 g, 0.392 mmol) and  
5 methyl 1-bromocyclohexanecarboxylate (0.274 g, 1.16 mmol) was added to the reaction and it was stirred for 2 h. The reaction mixture was cooled and quenched with 1 N HCl. The mixture was then diluted with water and extracted with CH<sub>2</sub>Cl<sub>2</sub>. The organic layer was dried (Na<sub>2</sub>SO<sub>4</sub>) and the solvent  
10 removed *in vacuo* to afford 0.400 g of crude product that was purified by preparative HPLC to give 0.059 g (60%) 1-(4-{3-[1-(4-*tert*-butyl-benzyl)-3-methyl-2-oxo-imidazolidin-4-yl]-propyl}-phenoxy)-cyclohexanecarboxylic acid. <sup>1</sup>H NMR. HRMS (ES<sup>+</sup>) *m/z* exact mass calcd for C<sub>31</sub>H<sub>43</sub>N<sub>2</sub>O<sub>4</sub> [M + 1] 507.3223,  
15 found 507.3237.

#### Example 186

2-(4-{3-[1-(4-*tert*-butyl-benzyl)-3-methyl-2-oxo-imidazolidin-4-yl]-propyl}-phenoxy)-2-ethyl-hexanoic acid




20 The procedures herein were utilized to prepare 2-(4-{3-[1-(4-*tert*-butyl-benzyl)-3-methyl-2-oxo-imidazolidin-4-yl]-propyl}-phenoxy)-2-ethyl-hexanoic acid. <sup>1</sup>H NMR. HRMS (ES<sup>+</sup>) *m/z* exact mass calcd for C<sub>32</sub>H<sub>47</sub>N<sub>2</sub>O<sub>4</sub> [M + 1] 523.3536, found 523.3555.

25

#### Example 187


2-(4-{3-[1-(4-*tert*-butyl-benzyl)-3-methyl-2-oxo-imidazolidin-4-yl]-propyl}-phenoxy)-pentanoic acid



The procedures herein were utilized to prepare 2-(4-{3-[1-(4-*tert*-butyl-benzyl)-3-methyl-2-oxo-imidazolidin-4-yl]-propyl}-phenoxy)-pentanoic acid.  $^1\text{H}$  NMR. HRMS ( $\text{ES}^+$ )  $m/z$  exact mass calcd for  $\text{C}_{29}\text{H}_{42}\text{N}_2\text{O}_4$  [ $\text{M} + 1$ ] 481.3066, found 481.3062.

#### Example 188

2-(4-{3-[1-(4-*tert*-butyl-benzyl)-3-methyl-2-oxo-imidazolidin-4-yl]-propyl}-phenoxy)-4-methyl-pentanoic acid




The procedures herein were utilized to prepare 2-(4-{3-[1-(4-*tert*-butyl-benzyl)-3-methyl-2-oxo-imidazolidin-4-yl]-propyl}-phenoxy)-4-methyl-pentanoic acid.  $^1\text{H}$  NMR. HRMS ( $\text{ES}^+$ )  $m/z$  exact mass calcd for  $\text{C}_{30}\text{H}_{43}\text{N}_2\text{O}_4$  [ $\text{M} + 1$ ] 495.3223, found 495.3244.

#### Example 189

##### Step A

2-bromo-2-methyl-pentanoic acid




A mixture of 2-methylpentanoic acid (10.0 g, 86.1 mmol), *N*-bromosuccinimide (22.98 g, 129 mmol) and conc.  $\text{H}_2\text{SO}_4$  (2.5 mL) in trifluoroacetic acid (50 mL) was heated to reflux under  $\text{N}_2$  for 16 h. The reaction was cooled and the solvent removed *in vacuo* and the crude oil was purified *via vacuum*

distillation (1 mm, 135-140 °C) to give a slowly crystallizing oil that was diluted with CH<sub>2</sub>Cl<sub>2</sub> and then filtered to remove solids. The solvent was removed *in vacuo* to afford 9.08 g (54%) of 2-bromo-2-methyl-pentanoic acid.

5 <sup>1</sup>H NMR.

Step B


**4-[3-(4-hydroxy-phenyl)-propyl]-3-methyl-1-(4-trifluoromethyl-benzyl)-imidazolidin-2-one**



10 A solution of 4-[3-(4-methoxy-phenyl)-propyl]-3-methyl-1-(4-trifluoromethyl-benzyl)-imidazolidin-2-one (3.51 g, 8.63 mmol) in dry CH<sub>2</sub>Cl<sub>2</sub> (60 mL), cooled to -78 °C and then treated dropwise with BBr<sub>3</sub> (6.49 g, 25.9 mmol). The reaction was warmed 0 °C and stirred for 2 h under N<sub>2</sub>. The  
15 reaction mixture was diluted with water and extracted with CH<sub>2</sub>Cl<sub>2</sub>. The organic layer was dried (MgSO<sub>4</sub>), and the solvent removed *in vacuo* to give crude product that was purified by flash chromatography using 2:1 hexanes:acetone to afford  
20 3.01 g (89%) 4-[3-(4-hydroxy-phenyl)-propyl]-3-methyl-1-(4-trifluoromethyl-benzyl)-imidazolidin-2-one. <sup>1</sup>H NMR. MS (ES<sup>+</sup>) Calc'd for C<sub>21</sub>H<sub>24</sub>N<sub>2</sub>O<sub>2</sub>F<sub>3</sub> (M +1) 393. Found *m/z* 393 (100%).

Step C

**2-methyl-2-(4-{3-[3-methyl-2-oxo-1-(4-trifluoromethyl-benzyl)-imidazolidin-4-yl]-propyl}-phenoxy)-pentanoic acid ethyl ester**






A mixture of 4-[3-(4-hydroxy-phenyl)-propyl]-3-methyl-1-(4-trifluoromethyl-benzyl)-imidazolidin-2-one (0.51 g, 1.30 mmol) and 2-bromo-2-methyl-pentanoic acid (2.54 g, 13.0 mmol) in 2-methyl-2-propanol (10 mL) was heated to 45 °C under N<sub>2</sub> and then a solution of 1 M potassium *tert*-butoxide in 2-methyl-2-propanol (27.3 mL, 27.3 mmol) was added dropwise and the reaction was stirred at 45 °C for 40 minutes. The reaction mixture was cooled and quenched with 1 N HCl (50 mL). The mixture was then diluted with water and extracted with Et<sub>2</sub>O. The organic layer was dried (Na<sub>2</sub>SO<sub>4</sub>) and the solvent removed *in vacuo* to afford crude product that was combined with EtOH (50 mL) and conc. H<sub>2</sub>SO<sub>4</sub> (6 mL). The mixture was heated to reflux for 3 h. The reaction mixture was cooled and the solvent removed *in vacuo*. The residue was diluted with water and extracted with EtOAc and the organic layer was dried (Na<sub>2</sub>SO<sub>4</sub>). The solvent was removed *in vacuo* to give crude product that was purified by flash chromatography using 5:1 hexanes:acetone to afford 0.491 g (71%) 2-methyl-2-(4-{3-[3-methyl-2-oxo-1-(4-trifluoromethyl-benzyl)-imidazolidin-4-yl]-propyl}-phenoxy)-pentanoic acid ethyl ester. This material could be purified by chiral preparative HPLC (conditions, ) <sup>1</sup>H NMR. MS (ES<sup>+</sup>) *m/z* calcd for C<sub>29</sub>H<sub>38</sub>N<sub>2</sub>O<sub>4</sub>F<sub>3</sub> [M + 1] 535. Found 535 (100%).

25

Step D

2-methyl-2-(4-{3-[3-methyl-2-oxo-1-(4-trifluoromethyl-benzyl)-imidazolidin-4-yl]-propyl}-phenoxy)-pentanoic acid




A mixture of 2-methyl-2-(4-{3-[3-methyl-2-oxo-1-(4-trifluoromethyl-benzyl)-imidazolidin-4-yl]-propyl}-phenoxy)-pentanoic acid ethyl ester (0.020 g, 0.040 mmol) and 5 N NaOH (0.5 mL) in EtOH (6 mL) was heated at reflux for 1 h.

5 The reaction mixture was cooled and quenched with 1 N HCl. The mixture was then diluted with water and extracted with CH<sub>2</sub>Cl<sub>2</sub>. The organic layer was dried (Na<sub>2</sub>SO<sub>4</sub>) and the solvent removed *in vacuo* to afford 0.010 g (53%) 2-methyl-2-(4-{3-[3-methyl-2-oxo-1-(4-trifluoromethyl-benzyl)-imidazolidin-4-yl]-propyl}-phenoxy)-pentanoic acid. <sup>1</sup>H NMR. HRMS (ES<sup>+</sup>) *m/z* exact mass calcd for C<sub>27</sub>H<sub>34</sub>N<sub>2</sub>O<sub>4</sub>F<sub>3</sub> [M + 1] 507.2471, found 507.2459.

### Example 190

#### Step A

15 **2-methoxy-3-(4-{2-[3-methyl-2-oxo-1-(4-trifluoromethyl-benzyl)-imidazolidin-4-yl]-ethoxy}-phenyl)-propionic acid ethyl ester**



A mixture of 3-(4-hydroxy-phenyl)-2-methoxy-propionic acid ethyl ester 0.027 g, 0.120 mmol), toluene-4-sulfonic acid 2-[3-methyl-2-oxo-1-(4-trifluoromethyl-benzyl)-imidazolidin-4-yl]-ethyl ester (0.061 g, 0.133 mmol) and Cs<sub>2</sub>CO<sub>3</sub> (0.059 g, 0.181 mmol) in DMF (4 mL) was heated to 55 °C under N<sub>2</sub> for 16 h. The reaction was cooled and quenched with 1 N HCl (10 mL) and worked up extractively with Et<sub>2</sub>O and water. The organic layer was dried (MgSO<sub>4</sub>) and the solvent removed *in vacuo* to afford crude product that was purified by flash chromatography using 4:1 then 3:1 hexanes:acetone to afford 0.040 g (64%) 2-methoxy-3-(4-{2-[3-methyl-2-oxo-1-(4-trifluoromethyl-benzyl)-imidazolidin-4-yl]-ethoxy}-phenyl)-

20


25

30

propionic acid ethyl ester.  $^1\text{H}$  NMR. MS ( $\text{ES}^+$ ) Calc'd for  $\text{C}_{26}\text{H}_{32}\text{N}_2\text{O}_5\text{F}_3$  ( $M + 1$ ) 509. Found  $m/z$  509 (100%).

#### Step B

2-methoxy-3-(4-{2-[3-methyl-2-oxo-1-(4-trifluoromethyl-  
5 benzyl)-imidazolidin-4-yl]-ethoxy}-phenyl)-propionic acid




A solution of 2-methoxy-3-(4-{2-[3-methyl-2-oxo-1-(4-trifluoromethyl-benzyl)-imidazolidin-4-yl]-ethoxy}-phenyl)-propionic acid ethyl ester (0.040 g, 0.079 mmol) in ethanol  
10 (5 mL) was treated with aqueous 5 N NaOH (0.5 mL) and stirred at room temperature for 2 h. The solvent removed *in vacuo*. The resultant residue was acidified with aqueous 1 N HCl (10 mL) and extracted with  $\text{CH}_2\text{Cl}_2$ . The organic layer was dried ( $\text{Na}_2\text{SO}_4$ ) and the solvent removed *in vacuo* to  
15 afford 0.026 g (68%) 2-methoxy-3-(4-{2-[3-methyl-2-oxo-1-(4-trifluoromethyl-benzyl)-imidazolidin-4-yl]-ethoxy}-phenyl)-propionic acid.  $^1\text{H}$  NMR. MS ( $\text{ES}^+$ )  $m/z$  calc'd for  $\text{C}_{24}\text{H}_{28}\text{N}_2\text{O}_5\text{F}_3$  ( $M + 1$ ) 481. Found  $m/z$  481.

20

#### Example 191

2-ethoxy-3-(4-{2-[3-methyl-2-oxo-1-(4-trifluoromethyl-benzyl)-imidazolidin-4-yl]-ethoxy}-phenyl)-propionic acid




The procedures herein were utilized to prepare 2-ethoxy-3-(4-{2-[3-methyl-2-oxo-1-(4-trifluoromethyl-benzyl)-imidazolidin-4-yl]-ethoxy}-phenyl)-propionic acid.  $^1\text{H}$  NMR.  
25

MS (ES<sup>+</sup>) *m/z* calc'd for C<sub>25</sub>H<sub>30</sub>N<sub>2</sub>O<sub>5</sub>F<sub>3</sub> (M +1) 495. Found *m/z* 495.

### Example 192

5        3-(4-{2-[3-methyl-2-oxo-1-(4-trifluoromethyl-benzyl)-  
imidazolidin-4-yl]-ethoxy}-phenyl)-2-propoxy-propionic acid




The procedures herein were utilized to prepare 3-(4-{2-[3-methyl-2-oxo-1-(4-trifluoromethyl-benzyl)-imidazolidin-4-yl]-ethoxy}-phenyl)-2-propoxy-propionic acid. <sup>1</sup>H NMR. HRMS  
10        (ES<sup>+</sup>) *m/z* exact mass calc'd for C<sub>26</sub>H<sub>32</sub>N<sub>2</sub>O<sub>5</sub>F<sub>3</sub> (M +1) 509.2263. Found *m/z* 509.2268.

### Example 193

15

#### Step A

3-(4-benzyloxy-phenyl)-2-cyclopentyloxy-propionic acid  
cyclopentyl ester



A mixture of 3-(4-benzyloxy-phenyl)-2-hydroxy-propionic acid  
20        (1.0 g, 3.67 mmol), cyclopentyl iodide (7.2 g, 36.7 mmol)  
and silver (I) oxide (4.25 g, 18.3 mmol) in DMF (15 mL) was  
heated at 55 °C under N<sub>2</sub> for 72 h. The reaction mixture was  
cooled, diluted with Et<sub>2</sub>O and filtered through hyflo. The  
filtrate was extracted with water and the organic layer was  
25        dried (MgSO<sub>4</sub>) and the solvent removed *in vacuo* to afford  
crude product that was purified by flash chromatography  
using 10:1 hexanes:acetone to afford 0.255 g (17%) 3-(4-  
benzyloxy-phenyl)-2-cyclopentyloxy-propionic acid


cyclopentyl ester.  $R_f = 0.67$  (1:1 hexanes:acetone).  $^1\text{H}$  NMR.

### Step B

#### 2-cyclopentyloxy-3-(4-hydroxy-phenyl)-propionic acid

5

#### cyclopentyl ester




A mixture of 3-(4-benzyloxy-phenyl)-2-cyclopentyloxy-propionic acid cyclopentyl ester (0.137 g, 0.335 mmol) and 10% Pd/C (0.14 g) in EtOAc (70 mL) was purged with  $\text{N}_2$  then  $\text{H}_2$  and then stirred under a  $\text{H}_2$  balloon atmosphere for 3 h at room temperature. Upon reaction completion,  $\text{MgSO}_4$  was added and the mixture filtered through hyflo. The solvent was removed *in vacuo* to afford 0.090 g (84%) 2-cyclopentyloxy-3-(4-hydroxy-phenyl)-propionic acid cyclopentyl ester.  $^1\text{H}$  NMR. MS ( $\text{ES}^-$ ) Calc'd for  $\text{C}_{19}\text{H}_{25}\text{O}_4$  (M - 1) 317. Found  $m/z$  317 (100%).

### Step C

#### 2-cyclopentyloxy-3-(4-{2-[3-methyl-2-oxo-1-(4-trifluoromethyl-benzyl)-imidazolidin-4-yl]-ethoxy}-phenyl)-propionic acid cyclopentyl ester

20




A mixture of 2-cyclopentyloxy-3-(4-hydroxy-phenyl)-propionic acid cyclopentyl ester (0.090 g, 0.283 mmol), toluene-4-sulfonic acid 2-[3-methyl-2-oxo-1-(4-trifluoromethyl-benzyl)-imidazolidin-4-yl]-ethyl ester (0.140 g, 0.307 mmol) and  $\text{Cs}_2\text{CO}_3$  (0.138 g, 0.423 mmol) in DMF (10 mL) was heated to  $65^\circ\text{C}$  under  $\text{N}_2$  for 16 h. The reaction was cooled and quenched with 1 N HCl (10 mL) and worked up extractively

with Et<sub>2</sub>O and water. The organic layer was dried (MgSO<sub>4</sub>) and the solvent removed *in vacuo* to afford crude product that was purified by flash chromatography using 3:1 hexanes:acetone to afford 0.108 g (64%) 2-cyclopentyloxy-3-(4-{2-[3-methyl-2-oxo-1-(4-trifluoromethyl-benzyl)-imidazolidin-4-yl]-ethoxy}-phenyl)-propionic acid cyclopentyl ester. <sup>1</sup>H NMR. MS (ES<sup>+</sup>) Calc'd for C<sub>33</sub>H<sub>42</sub>N<sub>2</sub>O<sub>5</sub>F<sub>3</sub> (M +1) 603. Found m/z 603 (100%).

#### Step D

2-cyclopentyloxy-3-(4-{2-[3-methyl-2-oxo-1-(4-trifluoromethyl-benzyl)-imidazolidin-4-yl]-ethoxy}-phenyl)-propionic acid




A solution of 2-cyclopentyloxy-3-(4-{2-[3-methyl-2-oxo-1-(4-trifluoromethyl-benzyl)-imidazolidin-4-yl]-ethoxy}-phenyl)-propionic acid cyclopentyl ester (0.108 g, 0.179 mmol) in ethanol (10 mL) was treated with aqueous 5 N NaOH (1 mL) and stirred at room temperature for 3 h. The solvent removed *in vacuo*. The resultant residue was acidified with aqueous 1 N HCl (10 mL) and extracted with CH<sub>2</sub>Cl<sub>2</sub>. The organic layer was dried (Na<sub>2</sub>SO<sub>4</sub>) and the solvent removed *in vacuo* to afford 0.106 g of crude acid that was purified by chiral HPLC (20 x 250 mm Chiralpak AD, 3:2 heptane:IPA with 0.1% trifluoroacetic acid mobile phase, 14 mL/min, 225 nm) to give 0.080 g (83%) of 100% de 2-cyclopentyloxy-3-(4-{2-[3-methyl-2-oxo-1-(4-trifluoromethyl-benzyl)-imidazolidin-4-yl]-ethoxy}-phenyl)-propionic acid. <sup>1</sup>H NMR. HRMS (ES<sup>+</sup>) m/z exact mass calc'd for C<sub>28</sub>H<sub>34</sub>N<sub>2</sub>O<sub>5</sub>F<sub>3</sub> (M +1) 535.2420. Found m/z 535.2408.

## Example 194

## Step A


## 5-(4-benzyloxy-benzyl)-2,2-dimethyl-[1,3]dioxolan-4-one



- 5 A mixture of 3-(4-benzyloxy-phenyl)-2-hydroxy-propionic acid (2.0 g, 7.34 mmol), 2,2-dimethoxypropane (18.63 g, 179 mmol) and pyridinium *p*-toluene sulfonate (0.92 g, 3.66 mmol) in  $\text{CHCl}_3$  (80 mL) was heated to reflux for 40 minutes under  $\text{N}_2$ . The reaction was cooled and worked up extractively with
- 10  $\text{CH}_2\text{Cl}_2$  and water. The organic layer was dried ( $\text{Na}_2\text{SO}_4$ ) and the solvent removed *in vacuo* to afford crude product that was purified by flash chromatography using 10:1 hexanes:acetone to afford 2.01 g (88%) 5-(4-benzyloxy-benzyl)-2,2-dimethyl-[1,3]dioxolan-4-one.  $R_f = 0.53$  (1:1
- 15 hexanes:acetone).  $^1\text{H}$  NMR (500 MHz,  $\text{CDCl}_3$ )  $\delta$  7.43-7.35 (m, 4H), 7.34-7.29 (m, 1H), 7.16 (d, 2H,  $J = 8.80$  Hz), 6.91 (d, 2H,  $J = 8.80$  Hz), 5.04 (s, 2H), 4.61 (dd, 1H,  $J = 6.36$  Hz,  $J = 4.40$  Hz), 3.13 (dd, 1H,  $J = 14.67$  Hz,  $J = 4.40$  Hz), 2.99 (dd, 1H,  $J = 14.67$  Hz,  $J = 4.40$  Hz), 1.50 (s, 3H), 1.36 (s,
- 20 3H); MS ( $\text{ES}^+$ ) Calc'd for  $\text{C}_{19}\text{H}_{20}\text{O}_4$  ( $\text{M} + \text{NH}_4$ ) 330. Found  $m/z$  330 (100%).

## Step B

## 5-(4-hydroxy-benzyl)-2,2-dimethyl-[1,3]dioxolan-4-one




- 25 A mixture of 5-(4-benzyloxy-benzyl)-2,2-dimethyl-[1,3]dioxolan-4-one (1.0 g, 3.20 mmol) and 10% Pd/C (0.75 g) in EtOAc (40 mL) was purged with  $\text{N}_2$  then  $\text{H}_2$  and then stirred under a  $\text{H}_2$  balloon atmosphere for 3 h at room temperature. Upon reaction completion,  $\text{Na}_2\text{SO}_4$  was added and
- 30 the mixture filtered through hyflo. The solvent was removed

in vacuo to afford 0.747 g (100%) 5-(4-hydroxy-benzyl)-2,2-dimethyl-[1,3]dioxolan-4-one .  $^1\text{H}$  NMR (500 MHz,  $\text{CDCl}_3$ )  $\delta$  7.11 (d, 2H,  $J = 8.31$  Hz), 6.76 (d, 2H,  $J = 8.31$  Hz), 4.92 (bs, 1H), 4.61 (dd, 1H,  $J = 6.36$  Hz,  $J = 4.40$  Hz), 3.11 (dd, 1H,  $J = 14.67$  Hz,  $J = 4.40$  Hz), 2.98 (dd, 1H,  $J = 14.67$  Hz,  $J = 4.40$  Hz), 1.50 (s, 3H), 1.36 (s, 3H); MS ( $\text{ES}^-$ ) Calc'd for  $\text{C}_{12}\text{H}_{13}\text{O}_4$  (M - 1) 221. Found  $m/z$  221 (100%).

### Step C

#### 10 3-(4-hydroxy-phenyl)-2-isopropoxy-propionic acid ethyl ester




A 0  $^{\circ}\text{C}$  solution of 5-(4-hydroxy-benzyl)-2,2-dimethyl-[1,3]dioxolan-4-one (0.20 g, 0.900 mmol) and triethylsilane (1.05 g, 9.0 mmol) in dry  $\text{CH}_2\text{Cl}_2$  (10 mL) was treated dropwise with a 1 molar solution of  $\text{TiCl}_4$  in  $\text{CH}_2\text{Cl}_2$  (0.90 mL, 0.900 mmol) under  $\text{N}_2$ . The resultant red slurry was stirred at 0  $^{\circ}\text{C}$  for 15 minutes and then warmed to room temperature for 45 minutes. The reaction mixture was quenched with water and extracted with EtOAc. The organic layer was dried ( $\text{Na}_2\text{SO}_4$ ) and the solvent removed in vacuo to afford 0.320 g of crude acid that was carried on as is. The oil was dissolved in EtOH (25 mL) and treated with conc.  $\text{H}_2\text{SO}_4$  (1 mL) and then stirred at room temperature for 17 h under  $\text{N}_2$ . The solvent was removed in vacuo and the oil extracted with EtOAc and water. The organic layer was dried ( $\text{Na}_2\text{SO}_4$ ) and the solvent removed in vacuo to afford crude product that was purified by flash chromatography using 5:1 hexanes:acetone to give 0.158 g (70%) 3-(4-hydroxy-phenyl)-2-isopropoxy-propionic acid ethyl ester.  $R_f = 0.48$  (1:1 hexanes:acetone).  $^1\text{H}$  NMR (500 MHz,  $\text{CDCl}_3$ )  $\delta$  7.10 (d, 2H,  $J = 8.80$  Hz), 6.73 (d, 2H,  $J = 8.31$  Hz), 4.79 (bs, 1H) 4.20-



4.12 (m, 2H), 3.99 (dd, 1H,  $J = 8.31$  Hz,  $J = 4.89$  Hz), 3.49 (hp, 1H,  $J = 5.87$  Hz), 2.95-2.83 (m, 2H), 1.23 (t, 3H,  $J = 6.85$  Hz), 1.14 (d, 3H,  $J = 6.36$  Hz), 0.97 (d, 3H,  $J = 6.36$  Hz); MS (ES<sup>-</sup>) Calc'd for C<sub>14</sub>H<sub>19</sub>O<sub>4</sub> (M - 1) 251. Found  $m/z$  251 (100%). 97.6% ee by chiral HPLC assay (Chiralcel OJ, 4.6 x 250 mm, 90/10 heptane/IPA eluent, 1 mL/min, 266 nm).

#### Step D

3-(4-{2-[1-(4-hydroxy-benzyl)-3-methyl-2-oxo-imidazolidin-4-yl]-ethoxy}-phenyl)-2-isopropoxy-propionic acid ethyl ester




A mixture of 3-(4-hydroxy-phenyl)-2-isopropoxy-propionic acid ethyl ester (0.034 g, 0.134 mmol), toluene-4-sulfonic acid 2-[3-methyl-2-oxo-1-(4-trifluoromethyl-benzyl)-imidazolidin-4-yl]-ethyl ester (0.068 g, 0.149 mmol) and Cs<sub>2</sub>CO<sub>3</sub> (0.066 g, 0.202 mmol) in DMF (6 mL) was heated to 65 °C under N<sub>2</sub> for 16 h. The reaction was cooled and quenched with 1 N HCl (10 mL) and worked up extractively with Et<sub>2</sub>O and water. The organic layer was dried (MgSO<sub>4</sub>) and the solvent removed in vacuo to afford crude product that was purified by flash chromatography using 4:1 hexanes:acetone to afford 0.048 g (67%) 3-(4-{2-[1-(4-hydroxy-benzyl)-3-methyl-2-oxo-imidazolidin-4-yl]-ethoxy}-phenyl)-2-isopropoxy-propionic acid ethyl ester.  $R_f = 0.36$  (1:1 hexanes:acetone). <sup>1</sup>H NMR (500 MHz, CDCl<sub>3</sub>) δ <sup>1</sup>H NMR (500 MHz, CDCl<sub>3</sub>) □ 7.56 (d, 2H,  $J = 8.07$  Hz), 7.36 (d, 2H,  $J = 8.07$  Hz), 7.13 (d, 2H,  $J = 8.80$  Hz), 6.72 (d, 2H,  $J = 8.80$  Hz), 4.47, 4.36 (AB<sub>q</sub>, 2H,  $J = 15.16$  Hz), 4.20-4.12 (m, 2H), 4.00-3.94 (m, 3H), 3.66-3.59 (m, 1H), 3.48 (hp, 1H,  $J = 5.87$  Hz), 3.37 (t, 1H,  $J = 8.80$  Hz), 2.98 (t, 1H,  $J = 8.80$  Hz), 2.90-

2.85 (m, 2H), 2.84 (s, 3H), 2.27-2.23 (m, 1H), 1.93-1.84 (m, 1H), 1.23 (t, 3H,  $J = 6.85$  Hz), 1.14 (d, 3H,  $J = 5.87$  Hz), 0.97 (d, 3H,  $J = 5.87$  Hz). MS (ES<sup>+</sup>) Calc'd for C<sub>28</sub>H<sub>36</sub>N<sub>2</sub>O<sub>5</sub>F<sub>3</sub> (M +1) 537. Found  $m/z$  537 (100%).

5

### Step E

#### 3-(4-{2-[1-(4-hydroxy-benzyl)-3-methyl-2-oxo-imidazolidin-4-yl]-ethoxy}-phenyl)-2-isopropoxy-propionic acid




10 A solution of 3-(4-{2-[1-(4-hydroxy-benzyl)-3-methyl-2-oxo-imidazolidin-4-yl]-ethoxy}-phenyl)-2-isopropoxy-propionic acid ethyl ester (0.048 g, 0.089 mmol) in ethanol (8 mL) was treated with aqueous 5 N NaOH (1.5 mL) and stirred at room temperature for 2 h. The solvent removed in  
15 *vacuo*. The resultant residue was acidified with aqueous 1 N HCl (15 mL) and extracted with CH<sub>2</sub>Cl<sub>2</sub>. The organic layer was dried (Na<sub>2</sub>SO<sub>4</sub>) and the solvent removed *in vacuo* to afford 0.038 g (84%) 3-(4-{2-[1-(4-hydroxy-benzyl)-3-methyl-2-oxo-imidazolidin-4-yl]-ethoxy}-phenyl)-2-isopropoxy-  
20 propionic acid. <sup>1</sup>H NMR (500 MHz, CDCl<sub>3</sub>)  $\delta$  7.49 (d, 2H,  $J = 8.07$  Hz), 7.30 (d, 2H,  $J = 8.07$  Hz), 7.06 (d, 2H,  $J = 8.31$  Hz), 6.66 (d, 2H,  $J = 8.31$  Hz), 4.40, 4.31 (AB<sub>q</sub>, 2H,  $J = 15.65$  Hz), 4.03-4.01 (m, 1H), 3.90 (t, 2H,  $J = 5.87$  Hz), 3.59-3.53 (m, 1H), 3.48 (hp, 1H,  $J = 5.87$  Hz), 3.31 (t, 1H,  $J = 8.80$  Hz), 3.12-2.96 (m, 1H), 2.95 (t, 1H,  $J = 8.80$  Hz),  
25 2.90-2.79 (m, 1H), 2.78 (s, 3H), 2.27-2.23 (m, 1H), 1.93-1.84 (m, 1H), 1.14 (d, 3H,  $J = 5.87$  Hz), 0.97 (d, 3H,  $J = 5.87$  Hz). HRMS (ES<sup>+</sup>)  $m/z$  exact mass calc'd for C<sub>26</sub>H<sub>32</sub>N<sub>2</sub>O<sub>5</sub>F<sub>3</sub> (M +1) 509.2274. Found  $m/z$  509.2263.

30

**Example 195**


**2-(1-ethyl-propoxy)-3-(4-{2-[1-(4-hydroxy-benzyl)-3-methyl-2-oxo-imidazolidin-4-yl]-ethoxy}-phenyl)-propionic acid**



The procedures herein were utilized to prepare 2-(1-ethyl-propoxy)-3-(4-{2-[1-(4-hydroxy-benzyl)-3-methyl-2-oxo-imidazolidin-4-yl]-ethoxy}-phenyl)-propionic acid.  $^1\text{H}$  NMR. MS ( $\text{ES}^+$ )  $m/z$  calc'd for  $\text{C}_{28}\text{H}_{35}\text{N}_2\text{O}_5\text{F}_3$  ( $M + 1$ ) 537. Found  $m/z$  537.

**Example 196**


**2-methoxy-3-(4-{3-[3-methyl-2-oxo-1-(4-trifluoromethyl-benzyl)-imidazolidin-4-yl]-propyl}-phenyl)-propionic acid**



The procedures herein were utilized to prepare 2-methoxy-3-(4-{3-[3-methyl-2-oxo-1-(4-trifluoromethyl-benzyl)-imidazolidin-4-yl]-propyl}-phenyl)-propionic acid.  $^1\text{H}$  NMR. HRMS ( $\text{ES}^+$ )  $m/z$  exact mass calc'd for  $\text{C}_{25}\text{H}_{30}\text{N}_2\text{O}_4\text{F}_3$  ( $M + 1$ ) 479.2158. Found  $m/z$  479.2171.

**Example 197**

**2-methoxy-2-methyl-3-(4-{3-[3-methyl-2-oxo-1-(4-trifluoromethyl-benzyl)-imidazolidin-4-yl]-propyl}-phenyl)-propionic acid**




The procedures herein were utilized to prepare 2-methoxy-2-methyl-3-(4-{3-[3-methyl-2-oxo-1-(4-trifluoromethyl-benzyl)-imidazolidin-4-yl]-propyl}-phenyl)-propionic acid. <sup>1</sup>H NMR. HRMS (ES<sup>+</sup>) *m/z* exact mass calc'd for C<sub>26</sub>H<sub>32</sub>N<sub>2</sub>O<sub>4</sub>F<sub>3</sub> (M +1) 493.2314. Found *m/z* 493.2321.

### Example 198

#### Step A

#### 2-(4-trifluoromethyl-phenoxy)-propionic acid ethyl ester




10

A mixture of trifluoromethyl-*p*-cresol (9.74 g, 60.1 mmol), ethyl 2-bromopropionate (10.9 g, 60.1 mmol) and Cs<sub>2</sub>CO<sub>3</sub> (39.15 g, 120 mmol) in DMF (450 mL) was heated at 90 °C under N<sub>2</sub> for 16 h. The reaction was cooled and filtered using Et<sub>2</sub>O to rinse the solids. The filtrate was washed with 1 N HCl (200 mL) and water. The organic layer was dried (Na<sub>2</sub>SO<sub>4</sub>) and the solvent removed *in vacuo* to afford 19.03 g of crude product that was purified by flash chromatography using 6:1 hexanes:acetone to afford 13.96 g (89%) 2-(4-trifluoromethyl-phenoxy)-propionic acid ethyl ester. *R*<sub>f</sub> = 0.63 (1:1 hexanes:acetone). <sup>1</sup> NMR.

20

#### Step B

#### 3-(3-benzyloxy-phenyl)-3-hydroxy-2-methyl-2-(4-trifluoromethyl-phenoxy)-propionic acid ethyl ester




25

A -78 °C solution of 2-(4-trifluoromethyl-phenoxy)-propionic acid ethyl ester (10.0 g, 38.1 mmol) in THF (80 mL) was

transferred via cannula to a  $-78^{\circ}\text{C}$  solution of 1.5 M LDA-THF complex (45.8 mL, 68.7 mmol) in THF (80 mL). The resultant mixture was stirred for 5 minutes at  $-78^{\circ}\text{C}$  and then 3-benzyloxybenzaldehyde (7.3 g, 34.4 mmol) was added in one portion and the reaction mixture stirred at  $-78^{\circ}\text{C}$  for 5 minutes and was then quenched with a  $-78^{\circ}\text{C}$  THF solution of acetic acid (6.82 g, 113.5 mmol). The reaction mixture was diluted with aqueous saturated  $\text{NH}_4\text{Cl}$  and extracted with  $\text{Et}_2\text{O}$ . The organic layer was dried ( $\text{Na}_2\text{SO}_4$ ) and the solvent removed *in vacuo* to afford 19.84 g of crude product that was purified by flash chromatography using 8:1 then 4:1 hexanes:acetone to afford 7.88 g (44%) 3-(3-benzyloxy-phenyl)-3-hydroxy-2-methyl-2-(4-trifluoromethyl-phenoxy)-propionic acid ethyl ester as a mixture of diastereomers.  $R_f = 0.55$  (1:1 hexanes:acetone).  $^1\text{NMR}$ .

#### Step C

#### 3-(3-hydroxy-phenyl)-2-methyl-2-(4-trifluoromethyl-phenoxy)-propionic acid ethyl ester



20

Trifluoroacetic anhydride (6.99 g, 33.3 mmol) was added dropwise to a  $0^{\circ}\text{C}$  solution of 3-(3-benzyloxy-phenyl)-3-hydroxy-2-methyl-2-(4-trifluoromethyl-phenoxy)-propionic acid ethyl ester (7.88 g, 16.6 mmol) and pyridine (13.10 g, 0.166 mol) in  $\text{CH}_2\text{Cl}_2$  (150 mL). The reaction solution was warmed to room temperature and stirred under  $\text{N}_2$  for 3 h. The reaction solution was washed twice with aqueous 1 N  $\text{HCl}$  (175 mL each) and the organic layer dried ( $\text{MgSO}_4$ ). The solvent was removed *in vacuo* to give 9.03 g (95%) of crude

25


3-(3-benzyloxy-phenyl)-2-methyl-3-(2,2,2-trifluoro-acetoxy)-  
2-(4-trifluoromethyl-phenoxy)-propionic acid ethyl ester  
which was carried on immediately.

The 3-(3-benzyloxy-phenyl)-2-methyl-3-(2,2,2-trifluoro-  
5 acetoxymethyl)-2-(4-trifluoromethyl-phenoxy)-propionic acid ethyl  
ester (9.03 g, 15.8 mmol) was combined with 10% Pd/C (9.0 g)  
in ethyl acetate (200 mL). The mixture was purged with N<sub>2</sub>  
then H<sub>2</sub> and then stirred at room temperature under a H<sub>2</sub>  
balloon for approximately 48 h. The reaction mixture was  
10 filtered through hyflo to remove the catalyst and the  
resultant filtrate dried (MgSO<sub>4</sub>). The solvent was removed  
in vacuo to give crude product that was purified by flash  
chromatography using 3:1 hexanes:acetone to afford 1.96 g  
(32%) 3-(3-hydroxy-phenyl)-2-methyl-2-(4-trifluoromethyl-  
15 phenoxy)-propionic acid ethyl ester. The racemic phenol was  
purified by preparative chiral HPLC (Chiralcel OD, 5 x 37  
cm, 95:5 heptane:IPA mobile phase, 150 ml/min, 285 nm, GK8-  
A015150-031). R<sub>f</sub> = 0.32 (1:1 hexanes:acetone). <sup>1</sup> NMR. MS  
(ES<sup>-</sup>) Calc'd for C<sub>19</sub>H<sub>18</sub>O<sub>4</sub>F<sub>3</sub> (M - 1) 367. Found m/z 367 (100%).

20

#### Step D

**2-methyl-3-(3-{2-[3-methyl-2-oxo-1-(4-trifluoromethyl-  
benzyl)-imidazolidin-4-yl]-ethoxy}-phenyl)-2-(4-  
trifluoromethyl-phenoxy)-propionic acid ethyl ester**




25 A mixture of 3-(3-hydroxy-phenyl)-2-methyl-2-phenoxy-  
propionic acid ethyl ester (0.100 g, 0.271 mmol), toluene-4-  
sulfonic acid 2-[3-methyl-2-oxo-1-(4-trifluoromethyl-  
benzyl)-imidazolidin-4-yl]-ethyl ester (0.136 g, 0.298  
mmol) and Cs<sub>2</sub>CO<sub>3</sub> (0.133 g, 0.408 mmol) in DMF (8 mL) was

heated to 65 °C under N<sub>2</sub> for 17 h. The reaction was cooled and quenched with 1 N HCl (10 mL) and worked up extractively with Et<sub>2</sub>O and water. The organic layer was dried (MgSO<sub>4</sub>) and the solvent removed in vacuo to afford crude product  
 5 that was purified by flash chromatography using 4:1 hexanes:acetone to afford 0.113 g (64%) 2-methyl-3-(3-{2-[3-methyl-2-oxo-1-(4-trifluoromethyl-benzyl)-imidazolidin-4-yl]-ethoxy}-phenyl)-2-(4-trifluoromethyl-phenoxy)-propionic acid ethyl ester. <sup>1</sup>H NMR. MS (ES<sup>+</sup>) Calc'd for C<sub>33</sub>H<sub>35</sub>N<sub>2</sub>O<sub>5</sub>F<sub>6</sub> (M  
 10 +1) 653. Found m/z 653 (100%).

#### Step E

2-methyl-3-(3-{2-[3-methyl-2-oxo-1-(4-trifluoromethyl-benzyl)-imidazolidin-4-yl]-ethoxy}-phenyl)-2-(4-trifluoromethyl-phenoxy)-propionic acid  
 15




A solution of 2-methyl-3-(3-{2-[3-methyl-2-oxo-1-(4-trifluoromethyl-benzyl)-imidazolidin-4-yl]-ethoxy}-phenyl)-2-(4-trifluoromethyl-phenoxy)-propionic acid ethyl ester  
 20 (0.113 g, 0.173 mmol) in ethanol (12 mL) was treated with aqueous 5 N NaOH (2 mL) and heated to reflux 1 h. The reaction mixture was cooled, the solvent removed in vacuo. The resultant residue was acidified with aqueous 1 N HCl (15 mL) and extracted with CH<sub>2</sub>Cl<sub>2</sub>. The organic layer was dried  
 25 (Na<sub>2</sub>SO<sub>4</sub>) and the solvent removed in vacuo to afford 0.108 g (100%) 2-methyl-3-(3-{2-[3-methyl-2-oxo-1-(4-trifluoromethyl-benzyl)-imidazolidin-4-yl]-ethoxy}-phenyl)-2-(4-trifluoromethyl-phenoxy)-propionic acid. <sup>1</sup>H NMR. HRMS

(ES<sup>+</sup>) *m/z* exact mass calc'd for C<sub>31</sub>H<sub>31</sub>N<sub>2</sub>O<sub>5</sub>F<sub>6</sub> (M +1) 625.2137.  
Found *m/z* 625.2134.

#### Example 199


5        **2-methyl-3-(3-{2-[3-methyl-2-oxo-1-(4-trifluoromethyl-benzyl)-imidazolidin-4-yl]-ethoxy}-phenyl)-2-(3-trifluoromethyl-phenoxy)-propionic acid**



The procedures herein were utilized to prepare 2-methyl-3-  
10 (3-{2-[3-methyl-2-oxo-1-(4-trifluoromethyl-benzyl)-imidazolidin-4-yl]-ethoxy}-phenyl)-2-(3-trifluoromethyl-phenoxy)-propionic acid. <sup>1</sup>H NMR. HRMS (ES<sup>+</sup>) *m/z* exact mass calc'd for C<sub>31</sub>H<sub>31</sub>N<sub>2</sub>O<sub>5</sub>F<sub>6</sub> (M +1) 625.2137. Found *m/z* 625.2128.

#### 15        Example 200

**3-{4-[2-(1-hexyl-3-methyl-2-oxo-imidazolidin-4-yl)-ethoxy]-phenyl}-2-methyl-2-phenoxy-propionic acid**




The procedures herein were utilized to prepare 3-{4-[2-(1-  
20 hexyl-3-methyl-2-oxo-imidazolidin-4-yl)-ethoxy]-phenyl}-2-methyl-2-phenoxy-propionic acid. <sup>1</sup>H NMR. HRMS (ES<sup>+</sup>) *m/z* exact mass calc'd for C<sub>28</sub>H<sub>39</sub>N<sub>2</sub>O<sub>5</sub> (M +1) 483.2859. Found *m/z* 483.2882.



**Example 201**

**3-{4-[2-(1-benzo[1,3]dioxol-5-ylmethyl-3-methyl-2-oxo-imidazolidin-4-yl)-ethoxy]-phenyl}-2-methyl-2-phenoxy-propionic acid**




5

The procedures herein were utilized to prepare 3-{4-[2-(1-benzo[1,3]dioxol-5-ylmethyl-3-methyl-2-oxo-imidazolidin-4-yl)-ethoxy]-phenyl}-2-methyl-2-phenoxy-propionic acid. <sup>1</sup>H NMR. HRMS (ES<sup>+</sup>) m/z exact mass calc'd for C<sub>30</sub>H<sub>33</sub>N<sub>2</sub>O<sub>7</sub> (M +1)

10 533.2288. Found m/z 533.2305.

**Example 202**

**2-methyl-3-{4-[2-(3-methyl-2-oxo-1-quinolin-2-ylmethyl-imidazolidin-4-yl)-ethoxy]-phenyl}-2-phenoxy-propionic acid**




15

The procedures herein were utilized to prepare 2-methyl-3-{4-[2-(3-methyl-2-oxo-1-quinolin-2-ylmethyl-imidazolidin-4-yl)-ethoxy]-phenyl}-2-phenoxy-propionic acid. <sup>1</sup>H NMR. HRMS (ES<sup>+</sup>) m/z exact mass calc'd for C<sub>32</sub>H<sub>34</sub>N<sub>3</sub>O<sub>5</sub> (M +1) 540.2498.

20 Found m/z 540.2523.

**Example 203**


**2-methyl-3-(4-{2-[3-methyl-2-oxo-1-(4-trifluoromethoxy-benzyl)-imidazolidin-4-yl]-ethoxy}-phenyl)-2-phenoxy-propionic acid**



The procedures herein were utilized to prepare 2-methyl-3-(4-{2-[3-methyl-2-oxo-1-(4-trifluoromethoxy-benzyl)-imidazolidin-4-yl]-ethoxy}-phenyl)-2-phenoxy-propionic acid ethyl ester.  $^1\text{H}$  NMR. MS ( $\text{ES}^+$ )  $m/z$  calc'd for  $\text{C}_{30}\text{H}_{32}\text{N}_2\text{O}_6\text{F}_3$  ( $\text{M} + 1$ ) 573. Found  $m/z$  573.

#### Example 204


2-methyl-3-(4-{2-[3-methyl-2-oxo-1-(4-trifluoromethyl-benzyl)-imidazolidin-4-yl]-ethoxy}-phenyl)-2-p-tolyloxy-propionic acid



The procedures herein were utilized to prepare 2-methyl-3-(4-{2-[3-methyl-2-oxo-1-(4-trifluoromethyl-benzyl)-imidazolidin-4-yl]-ethoxy}-phenyl)-2-p-tolyloxy-propionic acid.  $^1\text{H}$  NMR. HRMS ( $\text{ES}^+$ )  $m/z$  exact mass calc'd for  $\text{C}_{31}\text{H}_{34}\text{N}_2\text{O}_5\text{F}_3$  ( $\text{M} + 1$ ) 571.2420. Found  $m/z$  571.2423.

#### Example 205


2-(benzo[1,3]dioxol-5-yloxy)-2-methyl-3-(4-{2-[3-methyl-2-oxo-1-(4-trifluoromethyl-benzyl)-imidazolidin-4-yl]-ethoxy}-phenyl)-propionic acid



The procedures herein were utilized to prepare 2-(benzo[1,3]dioxol-5-yloxy)-2-methyl-3-(4-{2-[3-methyl-2-oxo-1-(4-trifluoromethyl-benzyl)-imidazolidin-4-yl]-ethoxy}-phenyl)-propionic acid.  $^1\text{H}$  NMR. HRMS ( $\text{ES}^+$ )  $m/z$  exact mass  
 5 calc'd for  $\text{C}_{31}\text{H}_{32}\text{N}_2\text{O}_7\text{F}_3$  ( $M + 1$ ) 601.2162. Found  $m/z$  601.2180.

#### Example 206


3-(4-{2-[1-(4-hydroxy-benzyl)-3-methyl-2-oxo-imidazolidin-4-yl]-ethoxy}-phenyl)-2-methyl-2-(4-trifluoromethyl-phenoxy)-  
 10 propionic acid



The procedures herein were utilized to prepare 3-(4-{2-[1-(4-hydroxy-benzyl)-3-methyl-2-oxo-imidazolidin-4-yl]-ethoxy}-phenyl)-2-methyl-2-(4-trifluoromethyl-phenoxy)-  
 15 propionic acid.  $^1\text{H}$  NMR. HRMS ( $\text{ES}^+$ )  $m/z$  exact mass calc'd for  $\text{C}_{31}\text{H}_{31}\text{N}_2\text{O}_5\text{F}_6$  ( $M + 1$ ) 625.2137. Found  $m/z$  625.2144.

#### Example 207

3-(4-{2-[1-(3,4-dimethyl-benzyl)-3-methyl-2-oxo-imidazolidin-4-yl]-ethoxy}-phenyl)-2-methyl-2-phenoxy-  
 20 propionic acid




The procedures herein were utilized to prepare 3-(4-{2-[1-(3,4-dimethyl-benzyl)-3-methyl-2-oxo-imidazolidin-4-yl]-ethoxy}-phenyl)-2-methyl-2-phenoxy-propionic acid.  $^1\text{H}$  NMR.  
 25

HRMS (ES<sup>+</sup>) *m/z* exact mass calc'd for C<sub>31</sub>H<sub>37</sub>N<sub>2</sub>O<sub>5</sub> (M +1)  
517.2702. Found *m/z* 517.2704.

**Example 208**


- 5    2-(2-Butyl-4-{3-[1-(4-tert-butyl-benzyl)-3-methyl-2-oxo-  
imidazolidin-4-yl]-propyl}-phenoxy)-2-methyl-propionic acid



10

**Step A**

2-(2-Butyl-4-{3-[1-(4-tert-butyl-benzyl)-3-methyl-2-oxo-  
imidazolidin-4-yl]-propyl}-phenoxy)-2-methyl-propionic acid  
methyl ester



15

The titled compound was prepared using 2-(4-{3-[1-(4-tert-  
Butyl-benzyl)-3-methyl-2-oxo-imidazolidin-4-yl]-propyl}-2-  
iodo-phenoxy)-2-methyl-propionic acid methyl ester (0.121g,  
0.200 mmole) and n-butyl bornic acid (0.061, 0.600 mmole)

to produce a colorless oil (0.046g, 45%). Mass [EI+] 537 (M+H)<sup>+</sup>.


Step B

2-(2-Butyl-4-{3-[1-(4-tert-butyl-benzyl)-3-methyl-2-oxo-  
5 imidazolidin-4-yl]-propyl}-phenoxy)-2-methyl-propionic acid

The titled compound was prepared using 2-(2-Butyl-4-{3-[1-(4-tert-butyl-benzyl)-3-methyl-2-oxo-imidazolidin-4-yl]-  
propyl}-phenoxy)-2-methyl-propionic acid methyl ester  
10 (0.043g, 0.080 mmole) to produce a colorless oil (0.033g, 81%). Mass [EI+] 523 (M+H)<sup>+</sup>, [EI-] 521 (M-H)<sup>-</sup>.


**Example 209**

2-(4-{3-[1-(4-tert-Butyl-benzyl)-3-methyl-2-oxo-  
15 imidazolidin-4-yl]-propyl}-2-isobutyl-phenoxy)-2-methyl-  
propionic acid



Step A

20 2-(4-{3-[1-(4-tert-Butyl-benzyl)-3-methyl-2-oxo-  
imidazolidin-4-yl]-propyl}-2-isobutyl-phenoxy)-2-methyl-  
propionic acid methyl ester



The titled compound was prepared using 2-(4-{3-[1-(4-tert-Butyl-benzyl)-3-methyl-2-oxo-imidazolidin-4-yl]-propyl}-2-iodo-phenoxy)-2-methyl-propionic acid methyl ester (0.121g, 0.200 mmole) and isobutyl bornic acid (0.061, 0.600 mmole) to produce a colorless oil (0.089g, 83%). Mass [EI+] 537 (M+H)<sup>+</sup>.


#### Step B

2-(4-{3-[1-(4-tert-Butyl-benzyl)-3-methyl-2-oxo-imidazolidin-4-yl]-propyl}-2-isobutyl-phenoxy)-2-methyl-propionic acid

The titled compound was prepared using 2-(4-{3-[1-(4-tert-Butyl-benzyl)-3-methyl-2-oxo-imidazolidin-4-yl]-propyl}-2-isobutyl-phenoxy)-2-methyl-propionic acid methyl ester (0.088g, 0.164 mmole) to produce colorless oil (0.0747g, 87%). Mass [EI+] 523 (M+H)<sup>+</sup>, [EI-] 521 (M-H)<sup>-</sup>.


#### Example 210

2-(5-{3-[1-(4-tert-Butyl-benzyl)-3-methyl-2-oxo-imidazolidin-4-yl]-propyl}-biphenyl-2-yloxy)-2-methyl-propionic acid



### Step A

2-(5-{3-[1-(4-tert-Butyl-benzyl)-3-methyl-2-oxo-  
 5 imidazolidin-4-yl]-propyl}-biphenyl-2-yloxy)-2-methyl-  
 propionic acid methyl ester



The titled compound was prepared using 2-(4-{3-[1-(4-tert-  
 10 Butyl-benzyl)-3-methyl-2-oxo-imidazolidin-4-yl]-propyl}-2-  
 iodo-phenoxy)-2-methyl-propionic acid methyl ester (0.121g,  
 0.200 mmole) and phenyl bornic acid (0.073, 0.600 mmole) to  
 produce a colorless oil (0.103g, 93%). Mass [EI+] 557  
 (M+H)<sup>+</sup>.

15

### Step B

2-(5-{3-[1-(4-tert-Butyl-benzyl)-3-methyl-2-oxo-  
 imidazolidin-4-yl]-propyl}-biphenyl-2-yloxy)-2-methyl-  
 propionic acid


20 The titled compound was prepared using 2-(5-{3-[1-(4-tert-  
 Butyl-benzyl)-3-methyl-2-oxo-imidazolidin-4-yl]-propyl}-

biphenyl-2-yloxy)-2-methyl-propionic acid methyl ester (0.103g, 0.185 mmole) to produce colorless oil (0.0967g, 97%). Mass [EI+] 543 (M+H)<sup>+</sup>, [EI-] 541 (M-H)<sup>-</sup>.

5

**Example 211**


**2-(4-{3-[1-(4-tert-Butyl-benzyl)-3-methyl-2-oxo-imidazolidin-4-yl]-propyl}-2-vinyl-phenoxy)-2-methyl-propionic acid**



10

Step A

**2-(4-{3-[1-(4-tert-Butyl-benzyl)-3-methyl-2-oxo-imidazolidin-4-yl]-propyl}-2-vinyl-phenoxy)-2-methyl-propionic acid methyl ester**



15

2-(4-{3-[1-(4-tert-Butyl-benzyl)-3-methyl-2-oxo-imidazolidin-4-yl]-propyl}-2-iodo-phenoxy)-2-methyl-propionic acid methyl ester (0.264g, 0.435 mmole) and tributyl(vinyl)tin (0.207g, 0.652 mmole) were mixed in toluene (4.0 mL). After bubbled with nitrogen for 15 minutes, tetrakis(triphenylphosphine) palladium(0) (0.050g,

20



0.043 mmole) was added. The reaction was heated at 80°C overnight. The solvent was removed on rota-vapor, and the crude product was purified by column chromatography (silica gel, gradient elution 0-30% acetone in hexane) to provide a  
5 colorless oil (0.158g, 71%). Mass [EI+] 507 (M+H)<sup>+</sup>.


#### Step B

2-(4-{3-[1-(4-tert-Butyl-benzyl)-3-methyl-2-oxo-  
imidazolidin-4-yl]-propyl}-2-vinyl-phenoxy)-2-methyl-  
10 propionic acid

The titled compound was prepared using 2-(4-{3-[1-(4-tert-Butyl-benzyl)-3-methyl-2-oxo-imidazolidin-4-yl]-propyl}-2-vinyl-phenoxy)-2-methyl-propionic acid methyl ester (0.048g,  
15 0.094 mmole) to produce an oil (0.0445g, 95%). Mass [EI+] 493 (M+H)<sup>+</sup>, [EI-] 491 (M-H)<sup>-</sup>.

#### Example 212


2-(4-{3-[1-(4-tert-Butyl-benzyl)-3-methyl-2-oxo-  
20 imidazolidin-4-yl]-propyl}-2-ethyl-phenoxy)-2-methyl-  
propionic acid



Step A

2-(4-{3-[1-(4-tert-Butyl-benzyl)-3-methyl-2-oxo-imidazolidin-4-yl]-propyl}-2-ethyl-phenoxy)-2-methyl-propionic acid methyl ester

5



2-(4-{3-[1-(4-tert-Butyl-benzyl)-3-methyl-2-oxo-imidazolidin-4-yl]-propyl}-2-vinyl-phenoxy)-2-methyl-propionic acid methyl ester (0.110g, 0.217 mmole) was dissolved in absolute ethanol (4 mL). After purged the solution with N<sub>2</sub> for 15 min, 10% Pd/C (0.040g). The reaction was stirred under a hydrogen balloon at room temperature for 2 hours. The catalyst was removed through filtration, and solvent was removed on rota-vapor to provide a colorless oil (0.110g, 100%). Mass [EI+] 509 (M+H)<sup>+</sup>.

10

15

Step B

2-(4-{3-[1-(4-tert-Butyl-benzyl)-3-methyl-2-oxo-imidazolidin-4-yl]-propyl}-2-ethyl-phenoxy)-2-methyl-propionic acid

20


25

The titled compound was prepared using 2-(4-{3-[1-(4-tert-Butyl-benzyl)-3-methyl-2-oxo-imidazolidin-4-yl]-propyl}-2-ethyl-phenoxy)-2-methyl-propionic acid methyl ester (0.110g, 0.216 mmole) to produce an oil (0.0905g, 85%). Mass [EI+] 495 (M+H)<sup>+</sup>, [EI-] 493 (M-H)<sup>-</sup>.


## Example 213

2-(2-Allyl-4-{3-[1-(4-tert-butyl-benzyl)-3-methyl-2-oxo-imidazolidin-4-yl]-propyl}-phenoxy)-2-methyl-propionic acid

5

Step A

10 2-(2-Allyl-4-{3-[1-(4-tert-butyl-benzyl)-3-methyl-2-oxo-imidazolidin-4-yl]-propyl}-phenoxy)-2-methyl-propionic acid methyl ester



15

The titled compound was prepared using 2-(4-{3-[1-(4-tert-Butyl-benzyl)-3-methyl-2-oxo-imidazolidin-4-yl]-propyl}-2-iodo-phenoxy)-2-methyl-propionic acid methyl ester (0.200g, 0.330 mmole) and allyl tributyltin (0.218, 0.659 mmole) to produce a colorless oil (0.102g, 59%). Mass [EI+] 521 (M+H)<sup>+</sup>.

20

Step B

2-(2-Allyl-4-{3-[1-(4-tert-butyl-benzyl)-3-methyl-2-oxo-imidazolidin-4-yl]-propyl}-phenoxy)-2-methyl-propionic acid


5 The titled compound was prepared using 2-(2-Allyl-4-{3-[1-(4-tert-butyl-benzyl)-3-methyl-2-oxo-imidazolidin-4-yl]-propyl}-phenoxy)-2-methyl-propionic acid methyl ester (0.040g, 0.076 mmole) to produce a colorless oil (0.027g, 69%). Mass [EI+] 507 (M+H)<sup>+</sup>, [EI-] 508 (M-H)<sup>-</sup>.

10


**Example 214**

2-(4-{3-[1-(4-tert-Butyl-benzyl)-3-methyl-2-oxo-imidazolidin-4-yl]-propyl}-2-propyl-phenoxy)-2-methyl-propionic acid

15

Step A

20 2-(4-{3-[1-(4-tert-Butyl-benzyl)-3-methyl-2-oxo-imidazolidin-4-yl]-propyl}-2-propyl-phenoxy)-2-methyl-propionic acid methyl ester



2-(2-Allyl-4-{3-[1-(4-tert-butyl-benzyl)-3-methyl-2-oxo-  
imidazolidin-4-yl]-propyl}-phenoxy)-2-methyl-propionic acid  
5 methyl ester (0.060g, 0.115 mmole) was dissolved in absolute  
ethanol (10 mL). After purged the solution with N<sub>2</sub> for 15  
min, 10% Pd/C (0.030g). The reaction was stirred under a  
hydrogen balloon at room temperature for 2 hours. The  
catalyst was removed through filtration, and solvent was  
10 removed on rota-vapor to provide a colorless oil (0.053g,  
88%). Mass [EI+] 523 (M+H)<sup>+</sup>.


#### Step B

2-(4-{3-[1-(4-tert-Butyl-benzyl)-3-methyl-2-oxo-  
15 imidazolidin-4-yl]-propyl}-2-propyl-phenoxy)-2-methyl-  
propionic acid

The titled compound was prepared using 2-(4-{3-[1-(4-tert-  
Butyl-benzyl)-3-methyl-2-oxo-imidazolidin-4-yl]-propyl}-2-  
20 propyl-phenoxy)-2-methyl-propionic acid methyl ester  
(0.052g, 0.100 mmole) to produce an oil (0.050g, 100%).  
Mass [EI+] 509 (M+H)<sup>+</sup>, [EI-] 507 (M-H)<sup>-</sup>.


#### Example 215

25 2-Methyl-2-{2-methyl-4-[3-(3-methyl-2-oxo-imidazolidin-4-  
yl)-propyl]-phenoxy}-propionic acid



### Step A


2-(4-{3-[1-(4-tert-Butyl-benzyl)-3-methyl-2-oxo-  
5 imidazolidin-4-yl]-propyl}-2-methyl-phenoxy)-2-methyl-  
propionic acid methyl ester



10 To a solution of 2-(4-{3-[1-(4-tert-Butyl-benzyl)-3-methyl-  
2-oxo-imidazolidin-4-yl]-propyl}-2-methyl-phenoxy)-2-methyl-  
propionic acid (0.610g, 1.27 mmole) in methanol (10 mL) was  
added 15 drops of concentrated sulfuric acid. The mixture  
was stirred at room temperature overnight. After evaporated  
15 the solvent, the residue was partitioned between ethyl  
acetate (50 mL) and water (50 mL). The organic layer was  
washed by saturated sodium bicarbonate (50 mL), then brine  
(3x50 mL), dried over Na<sub>2</sub>SO<sub>4</sub>, filtered and concentrated in  
vacuo to produce an oil (0.624g, 99%). Mass [EI+] 495  
20 (M+H)<sup>+</sup>.

### Step B

2-Methyl-2-{2-methyl-4-[3-(3-methyl-2-oxo-imidazolidin-4-  
yl)-propyl]-phenoxy}-propionic acid methyl ester



2-(4-{3-[1-(4-tert-Butyl-benzyl)-3-methyl-2-oxo-  
imidazolidin-4-yl]-propyl}-2-methyl-phenoxy)-2-methyl-  
5 propionic acid methyl ester (0.620g, 1.25 mmole) was  
dissolved in 20% HOAc in EtOAc (50 mL). After purged the  
solution with N<sub>2</sub> for 15 min, 10% Pd/C (0.600g). The  
reaction was stirred under a hydrogen balloon at room  
temperature for overnight. The catalyst was removed through  
10 filtration, and solvent was removed on rota-vapor to provide  
a colorless oil (0.337g, 77%). Mass [EI+] 349 (M+H)<sup>+</sup>.


#### Step C

2-Methyl-2-{2-methyl-4-[3-(3-methyl-2-oxo-imidazolidin-4-  
15 yl)-propyl]-phenoxy}-propionic acid

The titled compound was prepared using 2-Methyl-2-{2-methyl-  
4-[3-(3-methyl-2-oxo-imidazolidin-4-yl)-propyl]-phenoxy}-  
propionic acid methyl ester (0.112g, 0.322 mmole) to produce  
20 an oil (0.054g, 50%). Mass [EI+] 335 (M+H)<sup>+</sup>, [EI-] 333 (M-  
H)<sup>-</sup>.

#### Example 216


2-(4-[3-(1-Biphenyl-4-ylmethyl-3-methyl-2-oxo-imidazolidin-  
4-yl)-propyl]-2-methyl-phenoxy)-2-methyl-propionic acid  
25



The titled compound was prepared using 2-Methyl-2-{2-methyl-4-[3-(3-methyl-2-oxo-imidazolidin-4-yl)-propyl]-phenoxy}-propionic acid methyl Ester (0.112g, 0.322 mmole) and 4-phenyl benzyl chloride (0.078g, 0.186 mmole) to produce an oil (0.054g, 34%). Mass [EI+] 501 (M+H)<sup>+</sup>, [EI-] 499 (M-H)<sup>-</sup>.

#### Example 217

2-(4-{3-[1-(4-Bromo-benzyl)-3-methyl-2-oxo-imidazolidin-4-yl]-propyl}-2-methyl-phenoxy)-2-methyl-propionic acid




The titled compound was prepared using 2-Methyl-2-{2-methyl-4-[3-(3-methyl-2-oxo-imidazolidin-4-yl)-propyl]-phenoxy}-propionic acid methyl Ester (0.100g, 0.287 mmole) and 4-bromo benzyl bromide (0.108g, 0.431 mmole) to produce an oil (0.058g, 40%). Mass [EI+] 503, 505 (M+H)<sup>+</sup>, [EI-] 501, 503 (M-H)<sup>-</sup>.



**Example 218**

2-(4-{3-[1-(3-Methoxy-benzyl)-3-methyl-2-oxo-imidazolidin-4-yl]-propyl}-2-methyl-phenoxy)-2-methyl-propionic acid




5

The titled compound was prepared using 2-Methyl-2-{2-methyl-4-[3-(3-methyl-2-oxo-imidazolidin-4-yl)-propyl]-phenoxy}-propionic acid methyl Ester (0.100g, 0.287 mmole) and 3-methoxyl benzyl bromide (0.087g, 0.431 mmole) to produce an oil (0.042g, 32%). Mass [EI+] 455 (M+H)<sup>+</sup>, [EI-] 453 (M-H)<sup>-</sup>.

**Example 219**

2-Methyl-2-{2-methyl-4-[3-(3-methyl-2-oxo-1-quinolin-2-ylmethyl-imidazolidin-4-yl)-propyl]-phenoxy}-propionic acid




The titled compound was prepared using 2-Methyl-2-{2-methyl-4-[3-(3-methyl-2-oxo-imidazolidin-4-yl)-propyl]-phenoxy}-propionic acid methyl Ester (0.100g, 0.287 mmole) and 2-chloromethyl-quinoline (0.092g, 0.431 mmole) to produce an oil (0.023g, 17%). Mass [EI+] 476 (M+H)<sup>+</sup>, [EI-] 474 (M-H)<sup>-</sup>.

20

**Example 220**

**2-Methyl-2-(2-methyl-4-[3-(3-methyl-1-naphthalen-2-ylmethyl-  
2-oxo-imidazolidin-4-yl)-propyl]-phenoxy}-propionic acid**


5



The titled compound was prepared using 2-Methyl-2-(2-methyl-4-[3-(3-methyl-2-oxo-imidazolidin-4-yl)-propyl]-phenoxy)-  
10 propionic acid methyl Ester (0.100g, 0.287 mmole) and 2-Bromomethyl-naphthalene (0.095g, 0.431 mmole) to produce an oil (0.080g, 59%). Mass [EI+] 476 (M+H)<sup>+</sup>, [EI-] 474 (M-H)<sup>-</sup>.

**Example 221**

15 **2-{4-[3-(1-Biphenyl-4-ylmethyl-3-methyl-2-oxo-imidazolidin-4-yl)-propyl]-2-butyl-phenoxy}-2-methyl-propionic acid**




20

Step A

2-(2-Butyl-4-{3-[3-methyl-2-oxo-1-(4-trifluoromethyl-benzyl)-imidazolidin-4-yl]-propyl}-phenoxy)-2-methyl-propionic acid ethyl ester

5




The titled compound was prepared using 4-[3-(3-Butyl-4-hydroxy-phenyl)-propyl]-3-methyl-1-(4-trifluoromethyl-benzyl)-imidazolidin-2-one (0.210g, 0.468 mmole) to produce an oil (0.258g, 98%). Mass [EI+] 563 (M+H)<sup>+</sup>.

10

Step B

2-(2-Butyl-4-{3-[3-methyl-2-oxo-1-(4-trifluoromethyl-benzyl)-imidazolidin-4-yl]-propyl}-phenoxy)-2-methyl-propionic acid ethyl ester

15



The titled compound was prepared using 2-(2-Butyl-4-{3-[3-methyl-2-oxo-1-(4-trifluoromethyl-benzyl)-imidazolidin-4-yl]-propyl}-phenoxy)-2-methyl-propionic acid ethyl ester

20


(0.215g, 0.382 mmole) to produce an oil (0.049g, 32%). Mass [EI+] 405 (M+H)<sup>+</sup>.

### Step C


- 5 2-{4-[3-(1-Biphenyl-4-ylmethyl-3-methyl-2-oxo-imidazolidin-4-yl)-propyl]-2-butyl-phenoxy}-2-methyl-propionic acid

The titled compound was prepared using 2-(2-Butyl-4-{3-[3-methyl-2-oxo-1-(4-trifluoromethyl-benzyl)-imidazolidin-4-yl]-propyl}-phenoxy)-2-methyl-propionic acid ethyl ester  
 10 (0.049g, 0.120 mmole) and 4-phenyl benzyl bromide (0.029g, 0.144 mmole) to produce an oil (0.042g, 65%). Mass [EI+] 543 (M+H)<sup>+</sup>, [EI-] 541 (M-H)<sup>-</sup>.


- 15 Certain compounds of the present invention can be prepared using the methods illustrated by the following schemes:




## Cyclic Urea Synthesis - VI




## Cyclic Urea Synthesis - VII





## Hydantoin and Imidazolone Synthesis - IX



## Cyclic Urea Synthesis - X







## Biological Assays

### Binding and Cotransfection Studies

- 5 The in vitro potency of compounds in modulating PPAR $\gamma$  and PPAR $\alpha$  receptors are determined by the procedures detailed below. DNA-dependent binding (ABCD binding) is carried out using SPA technology with PPAR receptors. Tritium-labeled PPAR $\alpha$  and PPAR $\gamma$  agonists are used as

radioligands for generating displacement curves and  $IC_{50}$  values with compounds of the invention. Cotransfection assays are carried out in CV-1 cells. The reporter plasmid contained an acylCoA oxidase (AOX) PPRE and TK promoter upstream of the luciferase reporter cDNA. Appropriate PPARs and RXR $\alpha$  are constitutively expressed using plasmids containing the CMV promoter. For PPAR $\alpha$  and PPAR $\delta$ , interference by endogenous PPAR $\gamma$  in CV-1 cells is an issue. In order to eliminate such interference, a GAL4 chimeric system is used in which the DNA binding domain of the transfected PPAR is replaced by that of GAL4, and the GAL4 response element is utilized in place of the AOX PPRE. Cotransfection efficacy is determined relative to PPAR $\alpha$  agonist and PPAR $\gamma$  agonist reference molecules. Efficacies are determined by computer fit to a concentration-response curve, or in some cases at a single high concentration of agonist (10  $\mu$ M). For binding or cotransfection studies with receptors other than PPARs, similar assays are carried out using appropriate ligands, receptors, reporter constructs, etc., for that particular receptor.

These studies are carried out to evaluate the ability of compounds of the invention to bind to and/or activate various nuclear transcription factors, particularly huPPAR $\alpha$  ("hu" indicates "human") and huPPAR $\gamma$ . These studies provide in vitro data concerning efficacy and selectivity of compounds of the invention. Furthermore, binding and cotransfection data for compounds of the invention are compared with corresponding data for marketed compounds that act on either huPPAR $\alpha$  or huPPAR $\gamma$ .



Binding and cotransfection data for representative compounds of the invention are compared with corresponding data for reference to determine the binding.

The binding and cotransfection efficacy values found, for compounds of the invention and compounds of this invention which are useful for modulating a PPAR alpha receptor, are  $\leq 100$  nM and  $\geq 50\%$ , respectively. When coagonist modulators are desired, the values may be balanced against selectivity for the alpha, gamma, delta, or another desired PPAR receptor subtype.

Evaluation of Triglyceride Reduction and HDL Cholesterol  
Elevation in HuapoAI Transgenic Mice

Studies are performed to evaluate the effect of compounds of the present invention upon HDL and triglyceride levels in human apoAI mice. For each compound tested, seven to eight week old male mice, transgenic for human apoAI (C57BL/6-tgn(apoal)1rub, Jackson Laboratory, Bar Harbor, ME) are acclimated in individual cages for two weeks with standard chow diet (Purina 5001) and water provided *ad libitum*. After the acclimation, mice and chow are weighed and assigned to test groups ( $n = 5$ ) with randomization by body weight. Mice are dosed daily by oral gavage for 8 days using a 29 gauge, 1-1/2 inch curved feeding needle (Popper & Sons). The vehicle for the controls, test compounds and the positive control (fenofibrate 100mg/kg) is 1% carboxymethylcellulose (w/v) with 0.25% tween 80 (w/v). All mice are dosed daily between 6 and 8 a.m. with a dosing volume of 0.2ml. Prior to termination, animals and diets are weighed and body weight change and food consumption are calculated. Three hours after last dose, mice are euthanized with CO<sub>2</sub> and blood is removed (0.5-1.0 ml) by

cardiac puncture. After sacrifice, the liver, heart, and epididymal fat pad are excised and weighed. Blood is permitted to clot and serum is separated from the blood by centrifugation.

5       Cholesterol and triglycerides are measured colorimetrically using commercially prepared reagents (for example, as available from Sigma #339-1000 and Roche #450061 for triglycerides and cholesterol, respectively). The procedures are modified from published work (McGowan M. W.  
10 et al., Clin Chem 29:538-542,1983; Allain C. C. et al., Clin Chem 20:470-475,1974. Commercially available standards for triglycerides and total cholesterol, respectively, commercial quality control plasma, and samples are measured in duplicate using 200  $\mu$ l of reagent. An additional aliquot  
15 of sample, added to a well containing 200  $\mu$ l water, provides a blank for each specimen. Plates are incubated at room temperature on a plate shaker and absorbance is read at 500 nm and 540 nm for total cholesterol and triglycerides, respectively. Values for the positive control are always  
20 within the expected range and the coefficient of variation for samples is below 10%. All samples from an experiment are assayed at the same time to minimize inter-assay variability.

      Serum lipoproteins are separated and cholesterol  
25 quantitated by fast protein liquid chromatography (FPLC) coupled to an in line detection system. Samples are applied to a Superose 6 HR size exclusion column (Amersham Pharmacia Biotech) and eluted with phosphate buffered saline-EDTA at 0.5 ml/min. Cholesterol reagent (Roche Diagnostics Chol/HP  
30 704036) at 0.16ml/min is mixed with the column effluent through a T-connection and the mixture passed through a 15 m x 0.5 mm id knitted tubing reactor immersed in a 37 C water

bath. The colored product produced in the presence of cholesterol is monitored in the flow stream at 505 nm and the analog voltage from the monitor is converted to a digital signal for collection and analysis. The change in voltage corresponding to change in cholesterol concentration is plotted vs. time and the area under the curve corresponding to the elution of very low density lipoprotein (VLDL), low density lipoprotein (LDL) and high density lipoprotein (HDL) is calculated using Perkin Elmer Turbochrome software.

Triglyceride serum levels in mice dosed with a compound of the invention are compared to mice receiving the vehicle to identify compounds which could be particularly useful for lowering triglycerides. Generally, triglyceride decreases of greater than or equal to 30% (thirty percent) compared to control following a 30 mg/kg dose suggests a compound that can be especially useful for lowering triglyceride levels.

The percent increase of HDLc serum levels in mice receiving a compound of the invention is compared to mice receiving vehicle to identify compounds of the invention that could be particularly useful for elevating HDL levels. Generally, an increase of greater than or equal to 25% (twenty five percent) increase in HDLc level following a 30 mg/kg dose suggests a compound that can be especially useful for elevating HDLc levels.

It may be particularly desirable to select compounds of this invention that both lower triglyceride levels and increase HDLc levels. However, compounds that either lower triglyceride levels or increase HDLc levels may be desirable as well.

Evaluation of Glucose Levels in db/db Mice

The effects, upon plasma glucose of administering various dose levels of different compounds of the present invention is studied using the following methods.

5        Five-week-old male diabetic (db/db) mice [for example, C57BlKs/j-m +/+ Lepr(db), Jackson Laboratory, Bar Harbor, ME] or lean littermates are housed 6 per cage with food and water available at all times. After an acclimation period of 2 weeks, animals are individually identified by ear  
10 notches, weighed, and bled via the tail vein for determination of initial glucose levels. Blood is collected (100 µl) from unfasted animals by wrapping each mouse in a towel, cutting the tip of the tail with a scalpel, and milking blood from the tail into a heparinized capillary  
15 tube. Sample is discharged into a heparinized microtainer with gel separator and retained on ice. Plasma is obtained after centrifugation at 4°C and glucose measured immediately. Remaining plasma is frozen until the completion of the experiment, when glucose and triglycerides  
20 are assayed in all samples. Animals are grouped based on initial glucose levels and body weights. Beginning the following morning, mice are dosed daily by oral gavage for 7 days. Treatments are test compounds (30 mg/kg), a positive control agent (30 mg/kg) or vehicle [1%  
25 carboxymethylcellulose (w/v)/ 0.25% Tween80 (w/v); 0.3 ml/mouse]. On day 7, mice are weighed and bled (tail vein) 3 hours after dosing. Twenty-four hours after the 7<sup>th</sup> dose (i.e., day 8), animals are bled again (tail vein). Samples obtained from conscious animals on days 0, 7 and 8 are  
30 assayed for glucose. After the 24-hour bleed, animals are weighed and dosed for the final time. Three hours after dosing on day 8, animals are anesthetized by inhalation of

isoflurane and blood obtained via cardiac puncture (0.5-0.7 ml). Whole blood is transferred to serum separator tubes, chilled on ice and permitted to clot. Serum is obtained after centrifugation at 4°C and frozen until analysis for compound levels. After sacrifice by cervical dislocation, the liver, heart and epididymal fat pads are excised and weighed.

Glucose is measured colorimetrically using commercially purchased reagents. According to the manufacturers, the procedures are modified from published work (McGowan, M. W., Artiss, J. D., Strandbergh, D. R. & Zak, B. Clin Chem, 20:470-5 (1974) and Keston, A. Specific colorimetric enzymatic analytical reagents for glucose. Abstract of papers 129th Meeting ACS, 31C (1956).); and depend on the release of a mole of hydrogen peroxide for each mole of analyte, coupled with a color reaction first described by Trinder (Trinder, P. Determination of glucose in blood using glucose oxidase with an alternative oxygen acceptor. Ann Clin Biochem, 6:24 (1969)). The absorbance of the dye produced is linearly related to the analyte in the sample. The assays were further modified in our laboratory for use in a 96 well format. The commercially available standard for glucose, commercially available quality control plasma, and samples (2 or 5 µl/well) are measured in duplicate using 200 µl of reagent. An additional aliquot of sample, pipetted to a third well and diluted in 200 µl water, provides a blank for each specimen. Plates are incubated at room temperature for 18 minutes for glucose on a plate shaker (DPC Micormix 5) and absorbance is read at 500 nm on a plate reader. Sample absorbances are compared to a standard curve (100-800 for glucose). Values for the quality control sample are always within the expected range

and the coefficient of variation for samples is below 10%. All samples from an experiment are assayed at the same time to minimize inter-assay variability.

5        Evaluation of the Effects of Compounds of the Present  
         Invention upon A<sup>y</sup> Mice Body Weight, Fat Mass, Glucose and  
                         Insulin Levels

Female A<sup>y</sup> Mice

Female A<sup>y</sup> mice are singly housed, maintained under  
10    standardized conditions (22°C, 12 h light:dark cycle), and  
provided free access to food and water throughout the  
duration of the study. At twenty weeks of age the mice are  
randomly assigned to vehicle control and treated groups  
based on body weight and body fat content as assessed by  
15    DEXA scanning (N=6). Mice are then dosed via oral gavage  
with either vehicle or a Compound of this invention (50  
mg/kg) one hour after the initiation of the light cycle (for  
example, about 7 A.M.) for 18 days. Body weights are  
measured daily throughout the study. On day 14 mice are  
20    maintained in individual metabolic chambers for indirect  
calorimetry assessment of energy expenditure and fuel  
utilization. On day 18 mice are again subjected to DEXA  
scanning for post treatment measurement of body composition.

The results of p.o. dosing of compound for 18 days on  
25    body weight, fat mass, and lean mass are evaluated and  
suggest which compounds of this invention can be especially  
useful for maintaining desirable weight and/or promoting  
desired lean to fat mass.

Indirect calorimetry measurements reveal a significant  
30    reduction in respiratory quotient (RQ) in treated animals  
during the dark cycle [ $0.864 \pm 0.013$  (Control) vs.  $0.803 \pm$   
 $0.007$  (Treated);  $p < 0.001$ ]. This reduction in RQ is

indicative of an increased utilization of fat during the animals' active (dark) cycle. Additionally, treated animals display significantly higher rates of energy expenditure than control animals.

5

#### Male KK/A<sup>y</sup> Mice

Male KK/A<sup>y</sup> mice are singly housed, maintained under standardized conditions (22°C, 12 h light:dark cycle), and provided free access to food and water throughout the duration of the study. At twenty-two weeks of age the mice are randomly assigned to vehicle control and treated groups based on plasma glucose levels. Mice are then dosed via oral gavage with either vehicle or a Compound of this invention (30 mg/kg) one hour after the initiation of the light cycle (7 A.M.) for 14 days. Plasma glucose, triglyceride, and insulin levels are assessed on day 14.

The results of p.o. dosing of compound for 14 days on plasma glucose, triglycerides, and insulin are evaluated to identify compounds of this invention which may be especially desired.

20

#### Method to Elucidate the LDL-cholesterol Total-cholesterol and Triglyceride Lowering Effect

Male Syrian hamsters (Harlan Sprague Dawley) weighing 80-120 g are placed on a high-fat cholesterol-rich diet for two to three weeks prior to use. Feed and water are provided ad libitum throughout the course of the experiment. Under these conditions, hamsters become hypercholesterolemic showing plasma cholesterol levels between 180-280 mg/dl. (Hamsters fed with normal chow have a total plasma cholesterol level between 100-150 mg/dl.) Hamsters with high plasma cholesterol (180 mg/dl and above) are randomized

30

into treatment groups based on their total cholesterol level using the GroupOptimizeV211.xls program.

A Compound of this invention is dissolved in an aqueous vehicle (containing CMC with Tween 80) such that  
5 each hamster receives once a day approx. 1 ml of the solution by gavage at doses 3 and 30 mg/kg body weight. The blank control is vehicle alone. Dosing is performed daily in the early morning for 14 days.

10 **Quantification of Plasma Lipids:**

On the last day of the test, hamsters are bled (400 ul) from the suborbital sinus while under isoflurane anesthesia 2 h after dosing. Blood samples are collected into heparinized microfuge tubes chilled in ice bath. Plasma samples are  
15 separated from the blood cells by brief centrifugation. Total cholesterol and triglycerides are determined by means of enzymatic assays carried out automatically in the Monarch equipment (Instrumentation Laboratory) following the manufacturer's procedure. Plasma lipoproteins (VLDL, LDL  
20 and HDL) are resolved by injecting 25 ul of the pooled plasma samples into an FPLC system eluted with phosphate buffered saline at 0.5 ml/min through a Superose 6 HR 10/30 column (Pharmacia) maintained room temp. Detection and characterization of the isolated plasma lipids are  
25 accomplished by postcolumn incubation of the effluent with a Cholesterol/HP reagent (for example, Roche Lab System; infused at 0.12 ml/min) in a knitted reaction coil maintained at 37°C. The intensity of the color formed is proportional to the cholesterol concentration and is  
30 measured photometrically at 505 nm.

The effect of administration of a Compound of this invention for 14 days is studied for the percent reduction



in LDL level with reference to the vehicle group. The LDL-lowering efficacy for certain compounds of this invention can be especially desired. Compounds of this invention that decrease LDL greater than or equal to 30% (thirty percent) compared to vehicle can be especially desired.

The total-cholesterol and triglyceride lowering effects of a Compound of this invention are also studied. The data for reduction in total cholesterol and triglyceride levels after treatment with a compound of this invention for 14 days is compared to the vehicle to suggest compounds that can be particularly desired.

#### **Method to Elucidate the Fibrinogen-Lowering Effect of PPAR Modulators**

##### **Zucker Fatty Rat Model:**

The life phase of the study on fibrinogen-lowering effect of compounds of this invention is part of the life phase procedures for the antidiabetic studies of the same compounds. On the last (14<sup>th</sup>) day of the treatment period, with the animals placed under surgical anesthesia, ~ 3ml of blood is collected, by cardiac puncture, into a syringe containing citrate buffer. The blood sample is chilled and centrifuged at 4°C to isolate the plasma that is stored at -70 °C prior to fibrinogen assay.

##### **Quantification of Rat Plasma Fibrinogen:**

Rat plasma fibrinogen levels are quantified by using a commercial assay system consists of a coagulation instrument following the manufacturer's protocol. In essence, 100 ul of plasma is sampled from each specimen and a 1/20 dilution is prepared with buffer. The diluted plasma is incubated at 37°C for 240 seconds. Fifty microliters of clotting reagent thrombin solution (provided by the instrument's manufacturer in a standard concentration) is then added. The instrument

monitors the clotting time, a function of fibrinogen concentration quantified with reference to standard samples.

**Results:**

- 5 Compounds of this invention may lowering fibrinogen level *in vivo*. Compounds that lower fibrinogen level greater than vehicle can be especially desired.

10 Method to Elucidate the Anti-body Weight Gain and Anti-appetite Effects of Compounds of this invention

**Fourteen-Day Study in Zucker Fatty Rat<sup>1</sup> or ZDF Rat<sup>2</sup> Models:**

- Male Zucker Fatty rats, non-diabetic (Charles River Laboratories, Wilmington, MA) or male ZDF rats (Genetic Models, Inc, Indianapolis, IN) of comparable age and weight  
15 are acclimated for 1 week prior to treatment. Rats are on normal chow and water is provided *ad libitum* throughout the course of the experiment.
- 20 Compounds of this invention are dissolved in an aqueous vehicle such that each rat receives once a day approximately 1 ml of the solution by gavage at doses 0.1, 0.3, 1 and 3 mg/kg body weight. Fenofibrate (Sigma Chemical, prepared as a suspension in the same vehicle) a known alpha-agonist  
25 given at doses of 300 mg/kg, as well as the vehicle are controls. Dosing is performed daily in the early morning for 14 days. Over the course of the experiment, body weight and food consumption are monitored.
- 30 Using this assay, compounds of this invention are found to result in significant weight reduction.


**EQUIVALENTS:**

While this invention has been particularly shown and described with references to preferred embodiments thereof, it will be understood by those skilled in the art that  
5 various changes in form and details may be made therein without departing from the scope of the invention encompassed by the appended claims.

## CLAIMS

What is claimed is:

1. A Compound of the structural Formula I,




and pharmaceutically acceptable salts, solvates and hydrates thereof, wherein:

- (a) R1 is selected from the group consisting of hydrogen, substituted or unsubstituted group selected from C<sub>1</sub>-C<sub>8</sub> alkyl, aryl-C<sub>0-4</sub>-alkyl, heteroaryl-C<sub>0-4</sub>-alkyl, C<sub>3</sub>-C<sub>6</sub> cycloalkyl-aryl-C<sub>0-2</sub>-alkyl, and -CH<sub>2</sub>-C(O)-R17-R18, wherein R17 is O or NH and R18 is optionally substituted benzyl;
- (b) R2 is H or a substituted or unsubstituted group selected from the group consisting of C<sub>1</sub>-C<sub>6</sub> alkyl, C<sub>1</sub>-C<sub>6</sub> alkenyl, aryl-C<sub>0-4</sub>-alkyl, heteroaryl-C<sub>0-4</sub>-alkyl, C<sub>1</sub>-C<sub>4</sub> alkyl sulfonamide, C<sub>1</sub>-C<sub>4</sub> alkyl amide, OR10 and C<sub>3</sub>-C<sub>6</sub> cycloalkyl;
- (c) W is O or S;
- (d) X is an optionally substituted C<sub>1</sub>-C<sub>5</sub> alkylene linker wherein one carbon atom of the linker may optionally be replaced with O, NH, S, and optionally two carbons together may form a double bond;

- (e) Y is selected from the group consisting of C, O, S, NH and a single bond; and
- (f) E is selected from the group consisting of C(R3)(R4)A, A, and a substituted or unsubstituted group selected from the group consisting of (CH<sub>2</sub>)<sub>n</sub> COOR<sub>19</sub>; and wherein
- (i) n is 0, 1, 2 or 3;
- (ii) A is an functional group selected from the group consisting of carboxyl, C<sub>1</sub>-C<sub>3</sub>alkylnitrile, carboxamide, substituted or unsubstituted sulfonamide, substituted or unsubstituted acylsulfonamide substituted or unsubstituted tetrazole, and substituted or unsubstituted isoxazole;
- (iii) R<sub>3</sub> is selected from the group consisting of H, C<sub>1</sub>-C<sub>5</sub> alkyl, and C<sub>1</sub>-C<sub>5</sub> alkoxy, and
- (iv) R<sub>4</sub> is selected from the group consisting of H, halo, and a substituted or unsubstituted group selected from C<sub>1</sub>-C<sub>5</sub> alkyl, C<sub>1</sub>-C<sub>5</sub> alkoxy, C<sub>3</sub>-C<sub>6</sub> cycloalkyl, aryl C<sub>0</sub>-C<sub>4</sub> alkyl, arylC<sub>0</sub>-C<sub>2</sub>alkoxy and phenyl; or R<sub>3</sub> and R<sub>4</sub> are combined to form a C<sub>3</sub>-C<sub>8</sub> cycloalkyl;
- (v) R<sub>19</sub> is selected from the group consisting of hydrogen, optionally substituted arylmethyl and optionally substituted C<sub>1</sub>-C<sub>4</sub>alkyl;
- (g) R<sub>8</sub> is selected from the group consisting of hydrogen, C<sub>1</sub>-C<sub>4</sub> alkyl, C<sub>1</sub>-C<sub>4</sub> alkenyl, and halo;
- (h) R<sub>9</sub> is selected from the group consisting of hydrogen, C<sub>1</sub>-C<sub>4</sub> alkyl, C<sub>1</sub>-C<sub>4</sub> alkenyl, halo, substituted or unsubstituted aryl, substituted or unsubstituted aryl-C<sub>1</sub>-C<sub>4</sub> alkyl, substituted or unsubstituted heteroaryl, C<sub>1</sub>-C<sub>6</sub> alkenyl, and OR<sub>10</sub>;


- (i) R10 is independently selected from the group consisting of hydrogen and C<sub>1</sub>-C<sub>4</sub> alkyl;
- (j) R21 is selected from the group consisting of hydrogen, =O, and a substituted or unsubstituted group selected from the group consisting of C<sub>1</sub>-C<sub>6</sub> alkyl, aryl, C<sub>1</sub>-C<sub>4</sub> alkylaryl, and heteroaryl;
- (k) R22 is selected from the group consisting of hydrogen, and a substituted or unsubstituted group selected from the group consisting of C<sub>1</sub>-C<sub>6</sub> alkyl, aryl, C<sub>1</sub>-C<sub>4</sub> alkylaryl, and heteroaryl; and
- (l) ---- represents an optional double bond.

2. A compound as claimed by Claim 1 that is represented by the following structural formula:




and salts, solvates and hydrates thereof.

3. A compound as claimed by Claim 1 that is represented by the following structural formula:



and salts, solvates and hydrates thereof.

4. A compound as claimed by Claim 1 that is represented by the following structural formula:



and salts, solvates and hydrates thereof.

5. A compound as claimed by any one of Claims 1, 2, 3 or 4 wherein W is O.

5 6. A compound as claimed by any one of Claims 1, 2, 3, 4 or 5 wherein E is A.


7. A compound as claimed by any one of Claims 1, 2, 3, 4, 5 or 6 wherein A is COOH.

8. A compound as claimed by any one of Claims 1, 2, 3, 4, 5, 6 or 7 wherein Y is O.

9. A compound as claimed by any one of Claims 1, 2, 3, 4, 5, 6, or 7 wherein Y is C.


10. A compound as claimed by any one of Claims 1, 2, 3, 4, 5, 6, 8, or 9 wherein E is C(R3)(R4)A.

15 11. A compound as claimed by any one of Claims 1, 2, 3, 4, 5, 6, 8, or 9 wherein E is a group of the formula:



wherein R14 is selected from the group consisting of H, CF<sub>3</sub>, substituted or unsubstituted phenyl, substituted or unsubstituted aryl-C<sub>0-4</sub>-alkyl, and C<sub>1-6</sub>-alkyl.

12. A compound as claimed by any one of Claims 1, 2, 3, 4, 5, 6, 8, or 9 wherein E is a group of the formula:



wherein R17 is selected from the group consisting of H, and C<sub>1-6</sub>-alkyl.

13. A compound as claimed by any one of Claims 1-12,  
5 wherein X is optionally substituted C<sub>2</sub>-C<sub>5</sub> alkylene.

14. A compound as claimed by any one of Claims 1-13  
wherein X is propylene.

15. A compound as claimed by any one of Claims 1  
through 14 wherein the E-Y group is in the para position in  
10 relation to the X linker.

16. A compound as claimed by any one of Claims 1  
through 15 wherein R2 is C<sub>1</sub>-C<sub>2</sub> alkyl.

17. A compound as claimed by any one of Claims 1  
through 16 wherein R1 is substituted benzyl.

15 18. A compound as claimed by any one of Claims 1  
through 17 wherein R1 is substituted benzyl wherein the  
benzyl substituent is one or two independently selected from  
the group consisting of CF<sub>3</sub>, C<sub>1</sub>-C<sub>4</sub> alkyl, and halo.


19. A compound as claimed by any one of Claims 1-18  
20 wherein R1 is substituted benzyl wherein there is one para  
substituted benzyl substituent on the benzyl ring.

20. A compound as claimed by any one of Claims 1  
through 19 wherein X is C<sub>1</sub>-C<sub>3</sub>alkyl-O-.

21. A compound as claimed by any one of Claims 1-5, 8,  
25 9 and 12-20 wherein R4 is aryloxy.

22. A compound as claimed by any one of Claims 1, 4-  
12, 15-20 that is represented by the following structural  
formula:






wherein

- (a) LI is O or CH<sub>2</sub>;
- (b) V is a bond or a unsubstituted or substituted C1-C3 alkylene group;
- (c) R5 is substituted or unsubstituted group selected from the group consisting of aryl, heteroaryl and cycloalkyl groups; and
- (d) salts, solvates and hydrates thereof.


23. A compound as claimed by any one of Claims 1-5, 8, 9, and 12-21 wherein R4 has S stereochemistry in relation to R3.

24. A compound as claimed by any one of Claims 1, 4-12, 15-21, and 23 represented by the following structural Formula:




and salts, solvates and hydrates thereof.

25. A compound as claimed by any one of Claims 1, 4-12, 15-21 and 24 represented by the following structural formula:




wherein Q is C, O or S; and salts, solvates and hydrates thereof.

26. A compound as claimed by any one of Claims 1, 4-  
 5 12, 15-21 and 23-25 represented by the following structural formula:



wherein Q is C, O or S; and salts, solvates and hydrates thereof.


10 27. A compound as claimed by any one of Claims 1, 4-  
 12, 15-21 and 23-26 that is represented by the following structural formula:



15 and pharmaceutically acceptable salts, solvates and hydrates thereof, wherein:

- (a) X is an optionally substituted C<sub>1</sub>-C<sub>5</sub> alkylene linker wherein one carbon atom of the linker may be replaced with O, NH or S;
- (b) R<sub>3</sub> is H, C<sub>1</sub>-C<sub>5</sub> alkyl, or C<sub>1</sub>-C<sub>5</sub> alkoxy;
- 5 (c) R<sub>4</sub> is H, halo, a substituted or unsubstituted group selected from C<sub>1</sub>-C<sub>5</sub> alkyl, C<sub>1</sub>-C<sub>5</sub> alkoxy, aryloxy, C<sub>3</sub>-C<sub>6</sub> cycloalkyl and phenyl, or R<sub>3</sub> and R<sub>4</sub> are combined to form a C<sub>3</sub>-C<sub>8</sub> cycloalkyl;
- (d) V is a bond or a unsubstituted or substituted C<sub>1</sub>-C<sub>3</sub> alkylene group; and
- 10 (e) R<sub>5</sub> is substituted or unsubstituted group selected from the group consisting of aryl, heteroaryl, and cycloalkyl groups.

- 15 28. A compound as claimed by any one of Claims 1, 4-12, 15-21 and 23-27 that is represented by the following structural formula:




- and pharmaceutically acceptable salts, solvates and hydrates thereof, wherein:
- 20

- (a) X is an optionally substituted C<sub>1</sub>-C<sub>5</sub> alkylene linker wherein one carbon atom of the linker may be replaced with O, NH or S;
- (b) R<sub>3</sub> is H, C<sub>1</sub>-C<sub>5</sub> alkyl, or C<sub>1</sub>-C<sub>5</sub> alkoxy;

- (c) V is a bond or a unsubstituted or substituted C1-C3 alkylene group; and
- (d) R5 is substituted or unsubstituted group selected from the group consisting of aryl, heteroaryl and cycloalkyl groups.


29. A compound as claimed by any one of Claims 1, 4-12, 15-21 and 22-28 that is represented by the following structural formula:



and salts, solvates, and hydrates thereof;  
wherein

- (a) R3 is H or C<sub>1</sub>-C<sub>5</sub> alkyl;
- (b) V is a bond or a unsubstituted or substituted C1-C3 alkylene group;
- (c) R5 is substituted or unsubstituted group selected from the group consisting of aryl, heteroaryl and cycloalkyl groups.


30. A compound as claimed by any one of Claims 1, 4-12, 15-21, and 23-29 that is represented by the following structural formula:



wherein

- (a) R3 is H or C<sub>1</sub>-C<sub>5</sub> alkyl;
- (b) V is a bond or a unsubstituted or substituted C<sub>1</sub>-C<sub>3</sub> alkylene group;
- (c) R5 is substituted or unsubstituted group selected from the group consisting of aryl, heteroaryl and cycloalkyl groups; and
- (d) salts, solvates and hydrates thereof.

31. A Compound as claimed by Claim 1 that is represented by the following structural formula:



and salts, solvates, and hydrates thereof.

32. A compound as claimed by Claim 4 that is 2-methyl-3-(4-{3-[3-methyl-2-oxo-1-(4-trifluoromethyl-benzyl)-imidazolidin-4-yl]-propyl}-phenyl)-2-phenoxy-propionic acid or 2-Methyl-3-(4-{2-[3-methyl-2-oxo-1-(4-trifluoromethyl-benzyl)-imidazolidin-4-yl]-ethoxy}-phenyl)-2-phenoxy-propionic acid.

33. A pharmaceutical composition, comprising a pharmaceutically acceptable carrier and at least one

compound as claimed by any one of Claims 1-32, or a pharmaceutically acceptable salt, solvate or hydrate thereof.

34. A method of modulating a peroxisome proliferator  
5 activated receptor, comprising the step of contacting the receptor with at least one compound as claimed by any one of Claims 1-32, or a pharmaceutically acceptable salt, solvate or hydrate thereof.

35. A method of treating diabetes mellitus in a  
10 mammal, comprising the step of administering to the mammal in need thereof, a therapeutically effective amount of at least one compound of Claims 1-32, or a pharmaceutically acceptable salt, solvate or hydrate thereof.

36. A method of preventing diabetes mellitus in a  
15 mammal, comprising the step of administering to the mammal in need thereof, a pharmaceutically effective amount of at least one compound of Claims 1-32, or a pharmaceutically acceptable salt, solvate or hydrate thereof.

37. A method of treating Syndrome X in a mammal,  
20 comprising the step of administering to the mammal in need thereof, a therapeutically effective amount of at least one compound of Claims 1-32, or a pharmaceutically acceptable salt, solvate or hydrate thereof.

38. A method of treating cardiovascular disease in a  
25 mammal, comprising the step of administering to the mammal in need thereof, a therapeutically effective amount of at least one compound of Claims 1-32, or a pharmaceutically acceptable salt, solvate or hydrate thereof.

39. Use of a compound for the manufacture of a  
30 medicament for the treatment of a condition modulated by a

peroxisome proliferator activated receptor, wherein the compound, or pharmaceutically acceptable salt, solvate or hydrate thereof, is a compound of Claims 1-32.

40. All methods disclosed herein of preparing the  
5 compounds represented by Structural Formula I.

41. A compound as disclosed by any one of the examples herein.

42. A compound of Claim 1 selected from the group consisting of:

- 10 2-(4-{2-[1-(4-tert-Butyl-benzyl)-3-methyl-2-oxo-  
imidazolidin-4-yl]-ethoxy}-2-methyl-phenoxy)-2-methyl-  
propionic acid;  
2-(4-{2-[1-(4-Methoxy-benzyl)-3-methyl-2-oxo-imidazolidin-4-  
yl]-ethoxy}-2-methyl-phenoxy)-2-methyl-propionic acid ethyl  
15 ester 2-(4-{2-[1-(4-tert-Butyl-benzyl)-3-methyl-2-oxo-  
imidazolidin-4-yl]-ethoxy}-3-propyl-phenoxy)-2-methyl-  
propionic acid;  
2-(4-{3-[1-(4-tert-Butyl-benzyl)-3-methyl-2-oxo-  
imidazolidin-4-yl]-propyl}-2-methyl-phenoxy)-2-methyl-  
20 propionic acid;  
1-(4-tert-Butyl-benzyl)-4-{3-(4-methoxy-3-methyl-phenyl)-  
2-(2-Butyl-4-{3-[3-methyl-2-oxo-1-(4-trifluoromethyl-  
benzyl)-imidazolidin-4-yl]-propyl}-phenoxy)-2-methyl-  
propionic acid;  
25 2-Methyl-3-(4-{2-[3-methyl-2-oxo-1-(3-phenoxy-benzyl)-  
imidazolidin-4-yl]-ethoxy}-phenyl)-2-phenoxy-propionic acid;  
2-Methyl-3-(4-{2-[3-methyl-2-oxo-1-(3-trifluoromethoxy-  
benzyl)-imidazolidin-4-yl]-ethoxy}-phenyl)-2-phenoxy-  
propionic acid;  
30 2-Methyl-3-(4-{2-[3-methyl-2-oxo-1-(3-trifluoromethoxy-  
benzyl)-imidazolidin-4-yl]-ethoxy}-phenyl)-2-phenoxy-  
propionic acid;

- 2-(2-Fluoro-phenoxy)-3-(4-{2-[1-(4-methoxy-benzyl)-3-methyl-2-oxo-imidazolidin-4-yl]-ethoxy}-phenyl)-2-methyl-propionic acid;
- 2-(2-Fluoro-phenoxy)-3-(4-hydroxy-phenyl)-2-methyl-propionic acid methyl ester;
- 2-(2-Fluoro-phenoxy)-3-(4-{2-[1-(4-methoxy-benzyl)-3-methyl-2-oxo-imidazolidin-4-yl]-ethoxy}-phenyl)-2-methyl-propionic acid;
- 3-{4-[2-(1-Biphenyl-4-ylmethyl-3-methyl-2-oxo-imidazolidin-4-yl)-ethoxy]-phenyl}-2-(2-fluoro-phenoxy)-2-methyl-propionic acid;
- 3-(4-{2-[1-(4-tert-Butyl-benzyl)-3-methyl-2-oxo-imidazolidin-4-yl]-ethoxy}-phenyl)-2-(2-fluoro-phenoxy)-2-methyl-propionic acid;
- 2-(2-Fluoro-phenoxy)-2-methyl-3-{4-[2-(3-methyl-1-naphthalen-2-ylmethyl-2-oxo-imidazolidin-4-yl)-ethoxy]-phenyl}-propionic acid;
- 3-{4-[2-(1-Benzyl-3-methyl-2-oxo-imidazolidin-4-yl)-ethoxy]-phenyl}-2-(2-fluoro-phenoxy)-2-methyl-propionic acid;
- 3-(4-{2-[1-(4-Chloro-benzyl)-3-methyl-2-oxo-imidazolidin-4-yl]-ethoxy}-phenyl)-2-(2-fluoro-phenoxy)-2-methyl-propionic acid;
- 2-(2-Fluoro-phenoxy)-2-methyl-3-(4-{2-[3-methyl-2-oxo-1-(4-trifluoromethyl-benzyl)-imidazolidin-4-yl]-ethoxy}-phenyl)-propionic acid;
- 2-(2-Fluoro-phenoxy)-2-methyl-3-(4-{2-[3-methyl-2-oxo-1-(3-trifluoromethyl-benzyl)-imidazolidin-4-yl]-ethoxy}-phenyl)-propionic acid;
- 2-(2-Fluoro-phenoxy)-3-(4-{2-[1-(3-methoxy-benzyl)-3-methyl-2-oxo-imidazolidin-4-yl]-ethoxy}-phenyl)-2-methyl-propionic acid;



- 3-{4-[2-(1-Benzyl-3-methyl-2-oxo-imidazolidin-4-yl)-ethoxy]-phenyl}-2-(2-fluoro-phenoxy)-2-methyl-propionic acid
- 3-(4-{2-[1-(4-Chloro-benzyl)-3-methyl-2-oxo-imidazolidin-4-yl]-ethoxy}-phenyl)-2-(2-fluoro-phenoxy)-2-methyl-propionic acid;
- 5 2-(2-Fluoro-phenoxy)-2-methyl-3-(4-{2-[3-methyl-2-oxo-1-(4-trifluoromethyl-benzyl)-imidazolidin-4-yl]-ethoxy}-phenyl)-propionic acid;
- 2-(2-Fluoro-phenoxy)-2-methyl-3-(4-{2-[3-methyl-2-oxo-1-(3-trifluoromethyl-benzyl)-imidazolidin-4-yl]-ethoxy}-phenyl)-propionic acid;
- 10 2-(2-Fluoro-phenoxy)-3-(4-{2-[1-(3-methoxy-benzyl)-3-methyl-2-oxo-imidazolidin-4-yl]-ethoxy}-phenyl)-2-methyl-propionic acid ethyl ester;
- 15 2-(2-Fluoro-phenoxy)-3-(4-{2-[1-(3-methoxy-benzyl)-3-methyl-2-oxo-imidazolidin-4-yl]-ethoxy}-phenyl)-2-methyl-propionic acid ethyl ester;
- 3-{4-[2-(1-Benzyl-3-methyl-2-oxo-imidazolidin-4-yl)-ethoxy]-phenyl}-2-butoxy-2-methyl-propionic acid;
- 20 2-Butoxy-3-(4-{2-[1-(3-methoxy-benzyl)-3-methyl-2-oxo-imidazolidin-4-yl]-ethoxy}-phenyl)-2-methyl-propionic acid; 2-Butoxy-3-(4-{2-[1-(4-chloro-benzyl)-3-methyl-2-oxo-imidazolidin-4-yl]-ethoxy}-phenyl)-2-methyl-propionic acid;
- 2-Butoxy-2-methyl-3-(4-{2-[3-methyl-2-oxo-1-(3-trifluoromethyl-benzyl)-imidazolidin-4-yl]-ethoxy}-phenyl)-propionic acid;
- 25 2-Butoxy-3-(4-{2-[1-(4-chloro-benzyl)-3-methyl-2-oxo-imidazolidin-4-yl]-ethoxy}-phenyl)-2-methyl-propionic acid;
- 2-Butoxy-3-(4-{2-[1-(4-methoxy-benzyl)-3-methyl-2-oxo-imidazolidin-4-yl]-ethoxy}-phenyl)-2-methyl-propionic acid;
- 30 2-Ethoxy-3-(4-{2-[1-(4-methoxy-benzyl)-3-methyl-2-oxo-imidazolidin-4-yl]-ethoxy}-phenyl)-2-methyl-propionic acid;

- 2-Ethoxy-2-methyl-3-(4-{2-[3-methyl-2-oxo-1-(4-trifluoromethyl-benzyl)-imidazolidin-4-yl]-ethoxy}-phenyl)-propionic acid;
- 2-Methyl-3-(4-{2-[3-methyl-2-oxo-1-(4-trifluoromethyl-benzyl)-imidazolidin-4-yl]-ethoxy}-phenyl)-2-propoxy-propionic acid
- 5 2-Butoxy-2-methyl-3-(4-{2-[3-methyl-2-oxo-1-(4-trifluoromethyl-benzyl)-imidazolidin-4-yl]-ethoxy}-phenyl)-propionic acid;
- 2-Ethoxy-2-methyl-3-(4-{2-[3-methyl-2-oxo-1-(4-trifluoromethoxy-benzyl)-imidazolidin-4-yl]-ethoxy}-phenyl)-propionic acid;
- 10 2-Ethoxy-3-(4-{2-[1-(3-methoxy-benzyl)-3-methyl-2-oxo-imidazolidin-4-yl]-ethoxy}-phenyl)-2-methyl-propionic acid;
- 2-Ethoxy-2-methyl-3-(4-{2-[3-methyl-2-oxo-1-(3-trifluoromethoxy-benzyl)-imidazolidin-4-yl]-ethoxy}-phenyl)-propionic acid;
- 15 2-Ethoxy-2-methyl-3-(4-{2-[3-methyl-2-oxo-1-(3-trifluoromethyl-benzyl)-imidazolidin-4-yl]-ethoxy}-phenyl)-propionic acid;
- 3-{4-[2-(1-Biphenyl-4-ylmethyl-3-methyl-2-oxo-imidazolidin-4-yl)-ethoxy]-phenyl}-2-ethoxy-2-methyl-propionic acid;
- 20 3-(4-{2-[1-(3,4-Difluoro-benzyl)-3-methyl-2-oxo-imidazolidin-4-yl]-ethoxy}-phenyl)-2-methyl-2-phenoxy-propionic acid;
- 25 2-Ethoxy-2-methyl-3-(4-{2-[3-methyl-2-oxo-1-(4-trifluoromethyl-benzyl)-imidazolidin-4-yl]-ethoxy}-phenyl)-propionic acid;
- 2-Ethoxy-2-methyl-3-(4-{2-[3-methyl-2-oxo-1-(4-trifluoromethyl-benzyl)-imidazolidin-4-yl]-ethoxy}-phenyl)-propionic acid;
- 30 2-Methyl-3-(4-{2-[3-methyl-2-oxo-1-(4-trifluoromethyl-benzyl)-imidazolidin-4-yl]-ethoxy}-phenyl)-2-phenoxy-

- propionic acid; 2-Methyl-3-(4-{2-[3-methyl-2-oxo-1-(4-trifluoromethyl-benzyl)-imidazolidin-4-yl]-ethoxy}-phenyl)-2-phenoxy-propionic acid;
- 3-(4-{2-[3-Ethyl-2-oxo-1-(4-trifluoromethyl-benzyl)-imidazolidin-4-yl]-ethoxy}-phenyl)-2-methyl-2-phenoxy-propionic acid;
- 3-(4-{2-[3-(2-Methoxy-ethyl)-2-oxo-1-(4-trifluoromethyl-benzyl)-imidazolidin-4-yl]-ethoxy}-phenyl)-2-methyl-2-phenoxy-propionic acid; 2-Butoxy-2-methyl-3-(4-{2-[3-methyl-2-oxo-1-(3-trifluoromethyl-benzyl)-imidazolidin-4-yl]-ethoxy}-phenyl)-propionic acid; 3-{4-[2-(1-Biphenyl-4-ylmethyl-3-methyl-2-oxo-imidazolidin-4-yl)-ethoxy]-phenyl}-2-butoxy-2-methyl-propionic acid;
- 3-{4-[2-(1-Benzyl-3-methyl-2-oxo-imidazolidin-4-yl)-ethoxy]-phenyl}-2-butoxy-2-methyl-propionic acid;
- 2-Methoxy-2-methyl-3-(4-{2-[3-methyl-2-oxo-1-(4-trifluoromethyl-benzyl)-imidazolidin-4-yl]-ethoxy}-phenyl)-propionic acid; 2-Methoxy-2-methyl-3-(4-{2-[3-methyl-2-oxo-1-(4-trifluoromethyl-benzyl)-imidazolidin-4-yl]-ethoxy}-phenyl)-propionic acid; 2-Methyl-3-(4-{2-[3-methyl-2-oxo-1-(4-trifluoromethyl-benzyl)-imidazolidin-4-yl]-ethoxy}-phenyl)-2-phenoxy-propionic acid;
- 2,2-Dimethyl-3-(4-{2-[3-methyl-2-oxo-1-(4-trifluoromethyl-benzyl)-imidazolidin-4-yl]-ethoxy}-phenyl)-propionic acid;
- 2-methyl-3-(4-{2-[1-(4-methyl-benzyl)-2-oxo-3-propyl-imidazolidin-4-yl]-ethoxy}-phenyl)-2-phenoxy-propionic acid;
- 2-methyl-3-{4-[2-(3-methyl-1-naphthalen-2-ylmethyl-2-oxo-imidazolidin-4-yl)-ethoxy]-phenyl}-2-phenoxy-propionic acid;
- 2-methyl-3-{4-[2-(1-naphthalen-2-ylmethyl-2-oxo-imidazolidin-4-yl)-ethoxy]-phenyl}-2-phenoxy-propionic acid;
- 2-methyl-3-{4-[2-(1-naphthalen-2-ylmethyl-2-oxo-imidazolidin-4-yl)-ethoxy]-phenyl}-2-phenoxy-propionic acid;

- 3-(4-{2-[1-(3-methoxy-benzyl)-3-methyl-2-oxo-imidazolidin-4-yl]-ethoxy}-phenyl)-2-methyl-2-phenoxy-propionic acid;  
2-methyl-3-(4-{2-[3-methyl-2-oxo-1-(2-trifluoromethyl-benzyl)-imidazolidin-4-yl]-ethoxy}-phenyl)-2-phenoxy-  
5 propionic acid;  
3-(4-{2-[1-(2-methoxy-benzyl)-3-methyl-2-oxo-imidazolidin-4-yl]-ethoxy}-phenyl)-2-methyl-2-phenoxy-propionic acid;  
2-Methyl-3-(4-{2-[3-methyl-2-oxo-1-(4-trifluoromethyl-benzyl)-imidazolidin-4-yl]-ethoxy}-phenyl)-2-phenoxy-  
10 propionic acid;  
3-(4-{2-[1-(4-tert-Butyl-benzyl)-3-methyl-2-oxo-imidazolidin-4-yl]-ethoxy}-phenyl)-2-(4-fluoro-phenoxy)-2-methyl-propionic acid;  
2-(4-Fluoro-phenoxy)-2-methyl-3-{4-[2-(3-methyl-1-naphthalen-2-ylmethyl-2-oxo-imidazolidin-4-yl)-ethoxy]-phenyl}-propionic acid;  
15 3-(4-{2-[1-(3,4-Dimethyl-benzyl)-3-methyl-2-oxo-imidazolidin-4-yl]-ethoxy}-phenyl)-2-(4-fluoro-phenoxy)-2-methyl-propionic acid;  
3-(4-{2-[1-(4-tert-Butyl-benzyl)-3-methyl-2-oxo-imidazolidin-4-yl]-ethoxy}-phenyl)-2-methyl-2-(4-trifluoromethyl-phenoxy)-propionic acid;  
20 2-Methyl-3-{4-[2-(3-methyl-1-naphthalen-2-ylmethyl-2-oxo-imidazolidin-4-yl)-ethoxy]-phenyl}-2-(4-trifluoromethyl-phenoxy)-propionic acid;  
25 3-(4-{2-[1-(3,4-Dimethyl-benzyl)-3-methyl-2-oxo-imidazolidin-4-yl]-ethoxy}-phenyl)-2-methyl-2-(4-trifluoromethyl-phenoxy)-propionic acid;  
2-Methyl-3-(4-{2-[3-methyl-2-oxo-1-(3-phenoxy-benzyl)-imidazolidin-4-yl]-ethoxy}-phenyl)-2-phenoxy-propionic acid;  
30 2-Methyl-3-(4-{2-[3-methyl-2-oxo-1-(3-trifluoromethoxy-benzyl)-imidazolidin-4-yl]-ethoxy}-phenyl)-2-phenoxy-


- propionic acid; 2-Methyl-3-(4-{2-[3-methyl-2-oxo-1-(3-trifluoromethoxy-benzyl)-imidazolidin-4-yl]-ethoxy}-phenyl)-2-phenoxy-propionic acid;
- 2-(2-Fluoro-phenoxy)-2-methyl-3-(4-{2-[3-methyl-2-oxo-1-(4-trifluoromethoxy-benzyl)-imidazolidin-4-yl]-ethoxy}-phenyl)-propionic acid;
- 2-(2-Fluoro-phenoxy)-2-methyl-3-(4-{2-[3-methyl-1-(4-methyl-benzyl)-2-oxo-imidazolidin-4-yl]-ethoxy}-phenyl)-propionic acid;
- 3-(4-{2-[1-(3,4-Dichloro-benzyl)-3-methyl-2-oxo-imidazolidin-4-yl]-ethoxy}-phenyl)-2-(2-fluoro-phenoxy)-2-methyl-propionic acid;
- 3-(4-{2-[1-(3,4-Difluoro-benzyl)-3-methyl-2-oxo-imidazolidin-4-yl]-ethoxy}-phenyl)-2-(2-fluoro-phenoxy)-2-methyl-propionic acid;
- 3-(4-{2-[1-(3,5-Difluoro-benzyl)-3-methyl-2-oxo-imidazolidin-4-yl]-ethoxy}-phenyl)-2-(2-fluoro-phenoxy)-2-methyl-propionic acid;
- 2-(2-Fluoro-phenoxy)-2-methyl-3-{4-[2-(3-methyl-1-naphthalen-2-ylmethyl-2-oxo-imidazolidin-4-yl)-ethoxy]-phenyl}-propionic acid;
- 3-{4-[2-(1-Biphenyl-4-ylmethyl-3-methyl-2-oxo-imidazolidin-4-yl)-ethoxy]-phenyl}-2-methyl-2-phenoxy-propionic acid;
- 2-Methyl-3-(4-{2-[3-methyl-2-oxo-1-(4-trifluoromethoxy-benzyl)-imidazolidin-4-yl]-ethoxy}-phenyl)-2-phenoxy-propionic acid;
- 3-(4-{2-[1-(4-Fluoro-3-trifluoromethyl-benzyl)-3-methyl-2-oxo-imidazolidin-4-yl]-ethoxy}-phenyl)-2-methyl-2-phenoxy-propionic acid;
- 3-(4-{2-[1-(3-Fluoro-4-trifluoromethyl-benzyl)-3-methyl-2-oxo-imidazolidin-4-yl]-ethoxy}-phenyl)-2-methyl-2-phenoxy-propionic acid;

- 2-Methyl-3-(4-{2-[3-methyl-2-oxo-1-(3-trifluoromethyl-benzyl)-imidazolidin-4-yl]-ethoxy}-phenyl)-2-phenoxy-propionic acid;
- 3-(4-{2-[1-(3,4-Dichloro-benzyl)-3-methyl-2-oxo-
- 5 imidazolidin-4-yl]-ethoxy}-phenyl)-2-methyl-2-phenoxy-propionic acid;
- 3-(4-{2-[1-(3,5-Bis-trifluoromethyl-benzyl)-3-methyl-2-oxo-imidazolidin-4-yl]-ethoxy}-phenyl)-2-methyl-2-phenoxy-propionic acid;
- 10 3-(4-{2-[1-(3,5-Bis-trifluoromethyl-benzyl)-3-methyl-2-oxo-imidazolidin-4-yl]-ethoxy}-phenyl)-2-methyl-2-phenoxy-propionic acid;
- 3-(4-{2-[1-(4-Benzoyl-benzyl)-3-methyl-2-oxo-imidazolidin-4-yl]-ethoxy}-phenyl)-2-methyl-2-phenoxy-propionic acid;
- 15 3-(4-{2-[1-(4-Isopropyl-benzyl)-3-methyl-2-oxo-imidazolidin-4-yl]-ethoxy}-phenyl)-2-methyl-2-phenoxy-propionic acid;
- 2-Methyl-3-(4-{2-[3-methyl-1-(6-methyl-naphthalen-2-ylmethyl)-2-oxo-imidazolidin-4-yl]-ethoxy}-phenyl)-2-phenoxy-propionic acid;
- 20 2-Methyl-3-(3-{2-[3-methyl-2-oxo-1-(4-trifluoromethyl-benzyl)-imidazolidin-4-yl]-ethoxy}-phenyl)-2-phenoxy-propionic acid (diastereomer 4);
- 2-methyl-3-(4-{3-[3-methyl-2-oxo-1-(4-trifluoromethyl-benzyl)-imidazolidin-4-yl]-propyl}-phenyl)-2-phenoxy-
- 25 propionic acid;
- 2-(2-Butyl-4-{3-[1-(4-tert-butyl-benzyl)-3-methyl-2-oxo-imidazolidin-4-yl]-propyl}-phenoxy)-2-methyl-propionic acid;
- 2-(4-{3-[1-(4-tert-Butyl-benzyl)-3-methyl-2-oxo-imidazolidin-4-yl]-propyl}-2-isobutyl-phenoxy)-2-methyl-
- 30 propionic acid;

- 2-(5-{3-[1-(4-tert-Butyl-benzyl)-3-methyl-2-oxo-imidazolidin-4-yl]-propyl}-biphenyl-2-yloxy)-2-methyl-propionic acid;
- 2-(4-{3-[1-(4-tert-Butyl-benzyl)-3-methyl-2-oxo-imidazolidin-4-yl]-propyl}-2-vinyl-phenoxy)-2-methyl-propionic acid;
- 5 2-(4-{3-[1-(4-tert-Butyl-benzyl)-3-methyl-2-oxo-imidazolidin-4-yl]-propyl}-2-ethyl-phenoxy)-2-methyl-propionic acid;
- 10 2-(2-Allyl-4-{3-[1-(4-tert-butyl-benzyl)-3-methyl-2-oxo-imidazolidin-4-yl]-propyl}-phenoxy)-2-methyl-propionic acid;
- 2-(4-{3-[1-(4-tert-Butyl-benzyl)-3-methyl-2-oxo-imidazolidin-4-yl]-propyl}-2-propyl-phenoxy)-2-methyl-propionic acid;
- 15 2-Methyl-2-{2-methyl-4-[3-(3-methyl-2-oxo-imidazolidin-4-yl)-propyl]-phenoxy}-propionic acid;
- 2-{4-[3-(1-Biphenyl-4-ylmethyl-3-methyl-2-oxo-imidazolidin-4-yl)-propyl]-2-methyl-phenoxy}-2-methyl-propionic acid;
- 2-(4-{3-[1-(4-Bromo-benzyl)-3-methyl-2-oxo-imidazolidin-4-yl]-propyl}-2-methyl-phenoxy)-2-methyl-propionic acid;
- 20 2-(4-{3-[1-(3-Methoxy-benzyl)-3-methyl-2-oxo-imidazolidin-4-yl]-propyl}-2-methyl-phenoxy)-2-methyl-propionic acid;
- 2-Methyl-2-{2-methyl-4-[3-(3-methyl-2-oxo-1-quinolin-2-ylmethyl-imidazolidin-4-yl)-propyl]-phenoxy}-propionic acid;
- 25 2-Methyl-2-{2-methyl-4-[3-(3-methyl-1-naphthalen-2-ylmethyl-2-oxo-imidazolidin-4-yl)-propyl]-phenoxy}-propionic acid;
- 2-{4-[3-(1-Biphenyl-4-ylmethyl-3-methyl-2-oxo-imidazolidin-4-yl)-propyl]-2-butyl-phenoxy}-2-methyl-propionic acid;
- 2-(2-Butyl-4-{3-[1-(4-tert-butyl-benzyl)-3-methyl-2-oxo-imidazolidin-4-yl]-propyl}-phenoxy)-2-methyl-propionic acid;
- 30 2-(4-{3-[1-(4-tert-Butyl-benzyl)-3-methyl-2-oxo-

imidazolidin-4-yl]-propyl}-2-isobutyl-phenoxy)-2-methyl-propionic acid;  
2-(5-{3-[1-(4-tert-Butyl-benzyl)-3-methyl-2-oxo-imidazolidin-4-yl]-propyl}-biphenyl-2-yloxy)-2-methyl-propionic acid;  
5 2-(4-{3-[1-(4-tert-Butyl-benzyl)-3-methyl-2-oxo-imidazolidin-4-yl]-propyl}-2-vinyl-phenoxy)-2-methyl-propionic acid;  
2-(4-{3-[1-(4-tert-Butyl-benzyl)-3-methyl-2-oxo-imidazolidin-4-yl]-propyl}-2-ethyl-phenoxy)-2-methyl-propionic acid;  
10 2-(2-Allyl-4-{3-[1-(4-tert-butyl-benzyl)-3-methyl-2-oxo-imidazolidin-4-yl]-propyl}-phenoxy)-2-methyl-propionic acid;  
2-(4-{3-[1-(4-tert-Butyl-benzyl)-3-methyl-2-oxo-imidazolidin-4-yl]-propyl}-2-propyl-phenoxy)-2-methyl-propionic acid;  
15 2-Methyl-2-{2-methyl-4-[3-(3-methyl-2-oxo-imidazolidin-4-yl)-propyl]-phenoxy}-propionic acid;  
2-{4-[3-(1-Biphenyl-4-ylmethyl-3-methyl-2-oxo-imidazolidin-4-yl)-propyl]-2-methyl-phenoxy}-2-methyl-propionic acid;  
20 2-(4-{3-[1-(4-Bromo-benzyl)-3-methyl-2-oxo-imidazolidin-4-yl]-propyl}-2-methyl-phenoxy)-2-methyl-propionic acid;  
2-(4-{3-[1-(3-Methoxy-benzyl)-3-methyl-2-oxo-imidazolidin-4-yl]-propyl}-2-methyl-phenoxy)-2-methyl-propionic acid;  
25 2-Methyl-2-{2-methyl-4-[3-(3-methyl-2-oxo-1-quinolin-2-ylmethyl-imidazolidin-4-yl)-propyl]-phenoxy}-propionic acid;  
2-Methyl-2-{2-methyl-4-[3-(3-methyl-1-naphthalen-2-ylmethyl-2-oxo-imidazolidin-4-yl)-propyl]-phenoxy}-propionic acid;  
2-{4-[3-(1-Biphenyl-4-ylmethyl-3-methyl-2-oxo-imidazolidin-4-yl)-propyl]-2-butyl-phenoxy}-2-methyl-propionic acid;  
30 and a compound of the formula:





43. A compound of Claim 1, selected from the group consisting of:

2-(4-{3-[1-(4-tert-Butyl-benzyl)-2-oxo-2,3-dihydro-1H-imidazol-4-yl]-propyl}-phenoxy)-2-methyl-propionic acid,  
 2-(4-{3-[1-(3,4-Dimethyl-benzyl)-2-oxo-2,3-dihydro-1H-imidazol-4-yl]-propyl}-phenoxy)-2-methyl-propionic acid,  
 2-(4-{3-[1-(3-Methoxy-benzyl)-2-oxo-2,3-dihydro-1H-imidazol-4-yl]-propyl}-phenoxy)-2-methyl-propionic acid,  
 2-Methyl-2-{4-[3-(2-oxo-1-quinolin-2-ylmethyl-2,3-dihydro-1H-imidazol-4-yl)-propyl]-phenoxy}-propionic acid,  
 2-Methyl-2-(4-{3-[1-(4-methyl-benzyl)-2-oxo-2,3-dihydro-1H-imidazol-4-yl]-propyl}-phenoxy)-propionic acid,  
 2-(4-{3-[1-(3,5-Difluoro-benzyl)-2-oxo-2,3-dihydro-1H-imidazol-4-yl]-propyl}-phenoxy)-2-methyl-propionic acid,  
 2-{4-[3-(1-Benzyl-2-oxo-2,3-dihydro-1H-imidazol-4-yl)-propyl]-phenoxy}-2-methyl-propionic acid,  
 2-Methyl-2-{4-[3-(1-naphthalen-2-ylmethyl-2-oxo-2,3-dihydro-1H-imidazol-4-yl)-propyl]-phenoxy}-propionic acid,  
 2-Methyl-2-{4-[3-(2-oxo-1-phenethyl-2,3-dihydro-1H-imidazol-4-yl)-propyl]-phenoxy}-propionic acid,  
 2-(4-{3-[1-(4-Isopropyl-benzyl)-2-oxo-2,3-dihydro-1H-imidazol-4-yl]-propyl}-phenoxy)-2-methyl-propionic acid,  
 2-(4-{3-[1-(4-Ethyl-benzyl)-2-oxo-2,3-dihydro-1H-imidazol-4-yl]-propyl}-phenoxy)-2-methyl-propionic acid,  
 2-{4-[3-(1,3-Dibenzyl-2-oxo-2,3-dihydro-1H-imidazol-4-yl)-propyl]-phenoxy}-2-methyl-propionic acid,  
 2-(4-{3-[1-(3,4-Dimethyl-benzyl)-2-oxo-3-propyl-2,3-dihydro-1H-imidazol-4-yl]-propyl}-phenoxy)-2-methyl-

propionic acid,

2-(4-{3-[1-(4-tert-Butyl-benzyl)-3-methyl-2-oxo-2,3-dihydro-1H-imidazol-4-yl]-propyl}-phenoxy)-2-methyl-propionic acid,

2-(4-{3-[1-(4-tert-Butyl-benzyl)-3-ethyl-2-oxo-2,3-dihydro-1H-imidazol-4-yl]-propyl}-phenoxy)-2-methyl-propionic acid,

2-(4-{3-[1-(4-tert-Butyl-benzyl)-2-oxo-3-propyl-2,3-dihydro-1H-imidazol-4-yl]-propyl}-phenoxy)-2-methyl-propionic acid,

2-(4-{3-[1-(3-Methoxy-benzyl)-2-oxo-3-propyl-2,3-dihydro-1H-imidazol-4-yl]-propyl}-phenoxy)-2-methyl-propionic acid,

2-Methyl-2-(4-{3-[3-methyl-1-(4-methyl-benzyl)-2-oxo-2,3-dihydro-1H-imidazol-4-yl]-propyl}-phenoxy)-propionic acid,

2-Methyl-2-(4-[3-(2-oxo-3-propyl-1-quinolin-2-yl)methyl-2,3-dihydro-1H-imidazol-4-yl]-propyl)-phenoxy)-propionic acid,

2-Methyl-2-(4-{3-[1-(4-methyl-benzyl)-2-oxo-3-propyl-2,3-dihydro-1H-imidazol-4-yl]-propyl}-phenoxy)-propionic acid,

2-(4-{3-[3-Hexyl-1-(4-methyl-benzyl)-2-oxo-2,3-dihydro-1H-imidazol-4-yl]-propyl}-phenoxy)-2-methyl-propionic acid,

2-Methyl-2-(4-[3-(1-naphthalen-2-yl)methyl-2-oxo-3-propyl-2,3-dihydro-1H-imidazol-4-yl]-propyl)-phenoxy)-propionic acid,

2-(4-{3-[1-(3,5-Difluoro-benzyl)-2-oxo-3-propyl-2,3-dihydro-1H-imidazol-4-yl]-propyl}-phenoxy)-2-methyl-propionic acid,

2-(4-{3-[3-Cyclopropylmethyl-1-(3,5-difluoro-benzyl)-2-oxo-2,3-dihydro-1H-imidazol-4-yl]-propyl}-phenoxy)-2-methyl-propionic acid,

2-{4-[3-(1-Benzyl-2-oxo-3-propyl-2,3-dihydro-1H-imidazol-4-yl)-propyl]-phenoxy}-2-methyl-propionic acid,

2-(4-{3-[1-(3,4-Dimethyl-benzyl)-3-(2-methoxy-ethyl)-2-oxo-

2,3-dihydro-1H-imidazol-4-yl]-propyl}-phenoxy)-2-methyl-propionic acid,

2-(4-{3-[1-(3,4-Dimethyl-benzyl)-3-methyl-2-oxo-2,3-dihydro-1H-imidazol-4-yl]-propyl}-phenoxy)-2-methyl-propionic acid,

2-(4-{3-[1-(3-Chloro-benzyl)-2-oxo-3-propyl-2,3-dihydro-1H-imidazol-4-yl]-propyl}-phenoxy)-2-methyl-propionic acid,

2-(4-{3-[1-(3,4-Difluoro-benzyl)-2-oxo-3-propyl-2,3-dihydro-1H-imidazol-4-yl]-propyl}-phenoxy)-2-methyl-propionic acid,

2-(4-{3-[1-(3,4-Dichloro-benzyl)-2-oxo-3-propyl-2,3-dihydro-1H-imidazol-4-yl]-propyl}-phenoxy)-2-methyl-propionic acid,

2-(4-{3-[1-(4-Fluoro-benzyl)-2-oxo-3-propyl-2,3-dihydro-1H-imidazol-4-yl]-propyl}-phenoxy)-2-methyl-propionic acid,

2-(4-{3-[1-(3,5-Bis-trifluoromethyl-benzyl)-2-oxo-3-propyl-2,3-dihydro-1H-imidazol-4-yl]-propyl}-phenoxy)-2-methyl-propionic acid,

2-Methyl-2-{4-[3-(2-oxo-1-phenethyl-3-propyl-2,3-dihydro-1H-imidazol-4-yl)-propyl]-phenoxy}-propionic acid,

2-(4-{3-[1-(4-Isopropyl-benzyl)-2-oxo-3-propyl-2,3-dihydro-1H-imidazol-4-yl]-propyl}-phenoxy)-2-methyl-propionic acid,

2-(4-{3-[1-(4-Ethyl-benzyl)-2-oxo-3-propyl-2,3-dihydro-1H-imidazol-4-yl]-propyl}-phenoxy)-2-methyl-propionic acid,

2-(4-{3-[1-(3,4-Dimethyl-benzyl)-2-oxo-3-pyridin-4-ylmethyl-2,3-dihydro-1H-imidazol-4-yl]-propyl}-phenoxy)-2-methyl-propionic acid,

2-(4-{3-[3-(3,4-Dimethyl-benzyl)-2-oxo-1-propyl-2,3-dihydro-1H-imidazol-4-yl]-propyl}-phenoxy)-2-methyl-propionic acid,

2-(4-{3-[1-(3,4-Dimethyl-benzyl)-2,5-dioxo-imidazolidin-4-

yl]-propyl}-phenoxy)-2-methyl-propionic acid,  
2-Methyl-2-{4-[3-(1-naphthalen-2-ylmethyl-2,5-dioxo-imidazolidin-4-yl)-propyl]-phenoxy}-propionic acid,  
2-(4-{3-[1-(3,4-Dimethyl-benzyl)-2,5-dioxo-3-propyl-imidazolidin-4-yl]-propyl}-phenoxy)-2-methyl-propionic acid,  
2-(4-{3-[3-(4-Methoxy-benzyl)-1-naphthalen-2-ylmethyl-2,5-dioxo-imidazolidin-4-yl]-propyl}-phenoxy)-2-methyl-propionic acid,  
2-(4-{3-[1-(3,5-Difluoro-benzyl)-4-methyl-2,5-dioxo-3-propyl-imidazolidin-4-yl]-propyl}-phenoxy)-2-methyl-propionic acid,  
2-(4-{3-[1-(3,4-Dimethyl-benzyl)-4-methyl-2,5-dioxo-3-propyl-imidazolidin-4-yl]-propyl}-phenoxy)-2-methyl-propionic acid,  
2-(4-{3-[1-(3,4-Dimethyl-benzyl)-4-methyl-2,5-dioxo-imidazolidin-4-yl]-propyl}-phenoxy)-2-methyl-propionic acid,  
2-(4-{3-[1-(4-Ethyl-benzyl)-4-methyl-2,5-dioxo-imidazolidin-4-yl]-propyl}-phenoxy)-2-methyl-propionic acid,  
2-Methyl-2-{4-[3-(1-naphthalen-2-ylmethyl-2-oxo-imidazolidin-4-yl)-propyl]-phenoxy}-propionic acid, and  
2-{4-[2-(1,3-Dibenzyl-2,5-dioxo-imidazolidin-4-yloxy)-ethyl]-phenoxy}-2-methyl-propionic acid.

(19) World Intellectual Property Organization  
International Bureau(43) International Publication Date  
12 June 2003 (12.06.2003)

PCT

(10) International Publication Number  
WO 03/048130 A3(51) International Patent Classification<sup>7</sup>: C07D 233/32,  
A61K 31/4166, A61P 43/00, C07D 233/76, 233/70,  
401/06, 403/12, 405/06, 405/12

(21) International Application Number: PCT/US02/36128

(22) International Filing Date:  
26 November 2002 (26.11.2002)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:  
60/334,453 30 November 2001 (30.11.2001) US(71) Applicant (for all designated States except US): ELI  
LILLY AND COMPANY [US/US]; Lilly Corporate  
Center, Indianapolis, IN 46285 (US).

(72) Inventors; and

(75) Inventors/Applicants (for US only): GIBSON, Tracey,  
Ann [US/US]; 4416 Par Drive, Indianapolis, IN 46268  
(US). JOHNSTON, Richard, Duane [US/US]; 3712 For-  
est Lane, Greenfield, IN 46140 (US). MANTLO, Nathan,  
Bryan [US/US]; 7325 East 800 North, Brownsburg, IN  
46112 (US). THOMPSON, Richard, Craig [US/US]; 763  
North 900 West, Frankfort, IN 46041 (US). WANG, Xi-  
aodong [CN/US]; 14066 Honey Tree Drive, Carmel, IN  
46032 (US). WINNEROSKI, Leonard, Larry, Junior  
[US/US]; 587 Thoroughbred Lane, Greenwood, IN 46142  
(US). XU, Yanping [CN/US]; 13832 Barnett Place, Fish-  
ers, IN 46038 (US).(74) Agents: VORNDRAN-JONES, MaCharri et al.; Eli  
Lilly and Company, P. O. Box 6288, Indianapolis, IN  
46206-6288 (US).(81) Designated States (national): AE, AG, AL, AM, AT (uti-  
lity model), AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA,  
CH, CN, CO, CR, CU, CZ (utility model), CZ, DE (uti-  
lity model), DE, DK (utility model), DK, DM, DZ, EC, EE(utility model), EE, ES, FI (utility model), FI, GB, GD, GE,  
GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ,  
LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN,  
MW, MX, MZ, NO, NZ, OM, PH, PL, PT, RO, RU, SC,  
SD, SE, SG, SI, SK (utility model), SK, SL, TJ, TM, TN,  
TR, TT, TZ, UA, UG, US, UZ, VC, VN, YU, ZA, ZM, ZW.(84) Designated States (regional): ARIPO patent (GH, GM,  
KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW),  
Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM),  
European patent (AT, BE, BG, CH, CY, CZ, DE, DK, EE,  
ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, SK,  
TR), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GQ,  
GW, ML, MR, NE, SN, TD, TG).

## Declaration under Rule 4.17:


— as to applicant's entitlement to apply for and be granted  
a patent (Rule 4.17(ii)) for the following designations AE,  
AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA,  
CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, ES,  
FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE,  
KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD,  
MG, MK, MN, MW, MX, MZ, NO, NZ, OM, PH, PL, PT,  
RO, RU, SC, SD, SE, SG, SI, SK, SL, TJ, TM, TN, TR, TT,  
TZ, UA, UG, UZ, VC, VN, YU, ZA, ZM, ZW, ARIPO patent  
(GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW),  
Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM),  
European patent (AT, BE, BG, CH, CY, CZ, DE, DK, EE,  
ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, SK, TR),  
OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW,  
ML, MR, NE, SN, TD, TG)

## Published:

— with international search report

(88) Date of publication of the international search report:  
20 November 2003For two-letter codes and other abbreviations, refer to the "Guid-  
ance Notes on Codes and Abbreviations" appearing at the begin-  
ning of each regular issue of the PCT Gazette.

(54) Title: PEROXISOME PROLIFERATOR ACTIVATED RECEPTOR AGONISTS

(57) Abstract: The present invention is directed  
to compounds represented by the following struc-  
tural Formula (I), (c) W is O or S; (d) X is an op-  
tionally substituted C<sub>1</sub>-C<sub>5</sub> alkylene linker wherein  
one carbon atom of the linker may optionally be  
replaced with O, NH, S, and optionally two car-  
bons together may form a double bond; (e) Y is  
selected from the group consisting of C, O, S, NH  
and a single bond; and (f) E is selected from the  
group consisting of C(R<sub>3</sub>)(R<sub>4</sub>)A, A, and a sub-  
stituted or unsubstituted group selected from the  
group consisting of (CH<sub>2</sub>)<sub>n</sub> COOR<sub>19</sub>.

## INTERNATIONAL SEARCH REPORT

International Application No

PCT/US 02/36128

<b>A. CLASSIFICATION OF SUBJECT MATTER</b>		
IPC 7	C07D233/32 A61K31/4166 A61P43/00 C07D233/76 C07D233/70	C07D401/06 C07D403/12 C07D405/06 C07D405/12
According to International Patent Classification (IPC) or to both national classification and IPC		
<b>B. FIELDS SEARCHED</b>		
Minimum documentation searched (classification system followed by classification symbols)		
IPC 7 C07D A61K A61P		
Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched		
Electronic data base consulted during the international search (name of data base and, where practical, search terms used)		
CHEM ABS Data		
<b>C. DOCUMENTS CONSIDERED TO BE RELEVANT</b>		
Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	WO 98 39303 A (BOEHRINGER INGELHEIM PHARMACEUTICALS, INC.) 11 September 1998 (1998-09-11) * page 90, example 46; page 146, example 140; page 159, example 174 *	1
X	J. KNABE ET AL.: "Razemate und Enantiomere basisch substituierter 5-Phenylhydantoine, Synthese und antiarrhythmische Wirkung" PHARMAZIE., vol. 52, - 1997 pages 912-919, XP002240097 VEB VERLAG VOLK UND GESUNDHEIT. BERLIN., DD ISSN: 0031-7144 * compounds 4a and 4b *	1
--- -/--		
<input checked="" type="checkbox"/> Further documents are listed in the continuation of box C. <input checked="" type="checkbox"/> Patent family members are listed in annex.		
<b>* Special categories of cited documents:</b>		
<b>*A*</b> document defining the general state of the art which is not considered to be of particular relevance		
<b>*E*</b> earlier document but published on or after the international filing date		
<b>*L*</b> document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)		
<b>*O*</b> document referring to an oral disclosure, use, exhibition or other means		
<b>*P*</b> document published prior to the international filing date but later than the priority date claimed		
<b>*T*</b> later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention		
<b>*X*</b> document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone		
<b>*Y*</b> document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.		
<b>*Δ*</b> document member of the same patent family		
Date of the actual completion of the international search		Date of mailing of the international search report
13 May 2003		10/06/2003
Name and mailing address of the ISA European Patent Office, P.B. 5818 Patentlaan 2 NL - 2280 HV Rijswijk Tel. (+31-70) 340-2040, Tx. 31 651 epo nl, Fax: (+31-70) 340-3016		Authorized officer  Van Bijlen, H

## INTERNATIONAL SEARCH REPORT

International Application No

PCT/US 02/36128

## C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	ROBERT C. GLEN ET AL.: "Computer-aided design and synthesis of 5-substituted tryptamines and their pharmacology at the 5-HT1D receptor: discovery of compounds with potential anti-migraine properties" JOURNAL OF MEDICINAL CHEMISTRY., vol. 38, no. 18, - 1995 pages 3566-3580, XP002240098 AMERICAN CHEMICAL SOCIETY. WASHINGTON., US ISSN: 0022-2623 * page 3569, table 2, compound 38 *	1
X	DONALD L. ROSS ET AL.: "N-(gamma-L-Glutamyl)aminobenzoic acids" JOURNAL OF MEDICINAL CHEMISTRY., vol. 6, - 1963 pages 208-210, XP002240099 AMERICAN CHEMICAL SOCIETY. WASHINGTON., US ISSN: 0022-2623 * page 210, Table I, the last three compounds *	1
X	CHEMICAL ABSTRACTS, vol. 102, no. 12, 25 March 1985 (1985-03-25) Columbus, Ohio, US; abstract no. 103450a, "Silver halide color photographic material" XP002240103 abstract -& DATABASE CAPLUS 'Online! CHEMICAL ABSTRACTS SERVICE, COLUMBUS, OHIO, US; Database accession no. 1985:103450 XP002240106 compound with RN 95035-70-4 abstract & JP 59 159163 A (FUJI PHOTO FILM CO., LTD.)	1
X	EP 0 005 647 A (LILLY INDUSTRIES LTD.) 28 November 1979 (1979-11-28) * example 24; page 10, last line *	1
X	DE 27 24 948 A (THE WELLCOME FOUNDATION LTD.) 15 December 1977 (1977-12-15) * page 56: example 23c and 24c *	1

-/--

# INTERNATIONAL SEARCH REPORT

International Application No.

PCT/US 02/36128

## C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	<p>STANLEY C. BELL ET AL.: "Rearrangement reactions of phenyl chloroformate derivatives of 2-hydroxyaminoacetanilides to hydantoins, ureas and hydantoic acid derivatives"</p> <p>JOURNAL OF HETEROCYCLIC CHEMISTRY., vol. 13, no. 1, - 1976 pages 51-55, XP002240100</p> <p>HETEROCORPORATION. PROVO., US</p> <p>ISSN: 0022-152X</p> <p>* compound 30 *</p>	1
X	<p>JOSEPH WEINSTOCK ET AL.: "Synthesis of a 10,10a-dihydro-1H-imidazo-(3,4-b)(1,2)benzothiazine 5,5-dioxide"</p> <p>JOURNAL OF ORGANIC CHEMISTRY., vol. 33, no. 8, - 1968 pages 3342-3343, XP002240101</p> <p>AMERICAN CHEMICAL SOCIETY. EASTON., US</p> <p>ISSN: 0022-3263</p> <p>* page 3342, formula 2 *</p>	1
X	<p>WALFREDD S. SAAR ET AL.: "Tyrosine hydrolase inhibitors. Synthesis and activity of substituted aromatic amino acids"</p> <p>JOURNAL OF MEDICINAL CHEMISTRY., vol. 10, no. 6, - 1967 pages 1008-1014, XP002240102</p> <p>AMERICAN CHEMICAL SOCIETY. WASHINGTON., US</p> <p>ISSN: 0022-2623</p> <p>* page 1009, intermediate between III and V ; page 1012:</p> <p>5-methyl-5-(4-sulfamoylbenzyl)hydantoin *</p>	1
X	<p>US 3 452 041 A (STANLEY C. BELL ET AL.)</p> <p>24 June 1969 (1969-06-24)</p> <p>example 13</p>	1

-/--



## INTERNATIONAL SEARCH REPORT

International Application No.

PCT/US 02/36128

## C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	CHEMICAL ABSTRACTS, vol. 102, no. 17, 29 April 1985 (1985-04-29) Columbus, Ohio, US; abstract no. 145646z, STEIMAN, DAVID M. ET AL.: "Synthesis of side chain-protected amino acid phenylthiohydantoins and their use in quantitative solid-phase Edman degradation." XP002240104 & ANAL. BIOCHEM., vol. 145, no. 1, - 1985 pages 91-95, -& DATABASE CAPLUS 'Online! CHEMICAL ABSTRACTS SERVICE, COLUMBUS, OHIO, US; Database accession no. 1985:145646 XP002240107 compound with RN 95759-20-9 ---	1
X	CHEMICAL ABSTRACTS, vol. 99, no. 3, 18 July 1983 (1983-07-18) Columbus, Ohio, US; abstract no. 22893q, KENT, STEPHEN B. H. ET AL.: "A study of the Edman degradation in the assessment of the purity of synthetic peptides" XP002240105 abstract -& DATABASE CAPLUS 'Online! CHEMICAL ABSTRACTS SERVICE, COLUMBUS, OHIO, US; Database accession no. 1983:422893 XP002240108 compound with RN 86124-60-9 & METHODS PROTEIN SEQUENCE ANAL.(PROC. INT. CONF.), 4TH 1981, - 1982 pages 205-213, --- -/--	1

## INTERNATIONAL SEARCH REPORT

International Application No

PCT/US 02/36128

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT		
Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	CHEMICAL ABSTRACTS, vol. 134, 2001 Columbus, Ohio, US; abstract no. 246873n, JANDU, KARAMJIT SINGH ET AL.: "Discovery of 4-(3-trans-3-dimethylaminocyclobutyl)-1H-indol-5-ylmethyl)-(4S)-oxazolidin-2-one(499 1W93), a 5HT1H/1D receptor partial agonist and a potent inhibitor of electrically induced plasma extravasation." XP002240898 abstract -& DATABASE CAPLUS 'Online! CHEMICAL ABSTRACTS SERVICE, COLUMBUS, OHIO, US; Database accession no. 2001:55835 XP002240899 compound with RN 331441-72-6, -71-5, -70-4, -69-1 and -68-0 & J. MED: CHEM., vol. 44, no. 5, - 2001 pages 681-693, ---	1
A	WO 97 31907 A (GLAXO GROUP LTD.) 4 September 1997 (1997-09-04) claims 1,27 ---	1,39
A	WO 01 16120 A (ELI LILLY AND COMPANY) 8 March 2001 (2001-03-08) claims 1,78 -----	1,39

# INTERNATIONAL SEARCH REPORT

International application No.  
PCT/US 02/36128

## Box I Observations where certain claims were found unsearchable (Continuation of item 1 of first sheet)

This International Search Report has not been established in respect of certain claims under Article 17(2)(a) for the following reasons:

1. ☒ Claims Nos.:  
because they relate to subject matter not required to be searched by this Authority, namely:  
  
Although claims 35-38 are directed to a method of treatment of the human/animal body, the search has been carried out and based on the alleged effects of the compound/composition.
2. ☐ Claims Nos.:  
because they relate to parts of the International Application that do not comply with the prescribed requirements to such an extent that no meaningful International Search can be carried out, specifically:
3. ☐ Claims Nos.:  
because they are dependent claims and are not drafted in accordance with the second and third sentences of Rule 6.4(a).

## Box II Observations where unity of invention is lacking (Continuation of item 2 of first sheet)

This International Searching Authority found multiple inventions in this international application, as follows:

1. ☐ As all required additional search fees were timely paid by the applicant, this International Search Report covers all searchable claims.
2. ☐ As all searchable claims could be searched without effort justifying an additional fee, this Authority did not invite payment of any additional fee.
3. ☐ As only some of the required additional search fees were timely paid by the applicant, this International Search Report covers only those claims for which fees were paid, specifically claims Nos.:
4. ☐ No required additional search fees were timely paid by the applicant. Consequently, this International Search Report is restricted to the invention first mentioned in the claims; it is covered by claims Nos.:

### Remark on Protest

- ☐ The additional search fees were accompanied by the applicant's protest.
- ☐ No protest accompanied the payment of additional search fees.

## INTERNATIONAL SEARCH REPORT

Information on patent family members

International Application No

PCT/US 02/36128

Patent document cited in search report		Publication date	Patent family member(s)	Publication date
WO 9839303	A	11-09-1998	AT 233738 T	15-03-2003
			AU 6541898 A	22-09-1998
			BG 103711 A	28-09-2001
			BR 9811260 A	08-08-2000
			CN 1249748 T	05-04-2000
			DE 69811867 D1	10-04-2003
			EE 9900481 A	15-06-2000
			EP 0966447 A1	29-12-1999
			HU 0002347 A2	28-10-2000
			JP 2001513821 T	04-09-2001
			NO 994256 A	02-11-1999
			PL 336580 A1	03-07-2000
			SK 117499 A3	16-05-2000
			TR 9902124 T2	21-06-2000
			WO 9839303 A1	11-09-1998
			US 6355664 B1	12-03-2002
EP 5647	A	28-11-1979	GB 1601310 A	28-10-1981
			AR 225143 A1	26-02-1982
			AR 224186 A1	30-10-1981
			AT 366034 B	10-03-1982
			AT 373379 A	15-07-1981
			AT 366035 B	10-03-1982
			AT 551580 A	15-07-1981
			AU 530968 B2	04-08-1983
			AU 4728979 A	29-11-1979
			BE 876435 A1	21-11-1979
			BG 32115 A4	14-05-1982
			BG 31799 A3	15-03-1982
			CA 1120481 A1	23-03-1982
			CH 643546 A5	15-06-1984
			CH 639650 A5	30-11-1983
			CS 208654 B2	15-09-1981
			CS 208653 B2	15-09-1981
			DD 143769 A5	10-09-1980
			DE 2963061 D1	29-07-1982
			DK 207579 A	24-11-1979
			EP 0005647 A1	28-11-1979
			ES 480850 A1	01-02-1980
			FI 791614 A	24-11-1979
			FR 2426681 A1	21-12-1979
			GR 68363 A1	23-12-1981
			HU 184648 B	28-09-1984
			IL 57343 A	30-09-1982
			JP 54160373 A	19-12-1979
			KR 8300635 A	23-03-1983
			LU 81300 A1	16-12-1980
			MX 5604 E	07-11-1983
			NZ 190512 A	19-10-1984
			PL 215773 A1	14-07-1980
			PL 222701 A1	15-12-1980
			PT 69648 A	01-06-1979
			RO 76807 A1	26-04-1981
			RO 79703 A1	17-08-1982
			SU 873878 A3	15-10-1981
			SU 988190 A3	07-01-1983
			US 4241073 A	23-12-1980
			US 4293563 A	06-10-1981

# INTERNATIONAL SEARCH REPORT

Information on patent family members

International Application No

PCT/US 02/36128

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
EP 5647	A	YU 120179 A1 ZA 7902501 A	28-02-1983 28-05-1980
DE 2724948	A 15-12-1977	AT 373882 B AT 92080 A AT 374460 B AT 745579 A AU 2576877 A CA 1176261 A1 CH 647511 A5 CH 646426 A5 CH 645631 A5 DE 2724948 A1 DK 244577 A ES 459412 A1 ES 470428 A1 ES 478156 A1 ES 482010 A1 ES 8103057 A1 FI 771763 A , B, FR 2362839 A1 HU 183046 B IL 64162 A IT 1143583 B JP 52151168 A JP 62022986 B NL 7706064 A NZ 184281 A SE 446336 B SE 7706428 A SE 8203161 A US 4486443 A US 4204068 A US 4262128 A AU 515174 B2 BE 855337 A1 IL 52229 A ZA 7703324 A	27-02-1984 15-07-1983 25-04-1984 15-09-1983 07-12-1978 16-10-1984 31-01-1985 30-11-1984 15-10-1984 15-12-1977 04-12-1977 01-11-1978 01-09-1979 01-11-1979 16-08-1980 16-05-1981 04-12-1977 24-03-1978 28-04-1984 29-02-1984 22-10-1986 15-12-1977 20-05-1987 06-12-1977 28-04-1980 01-09-1986 04-12-1977 19-05-1982 04-12-1984 20-05-1980 14-04-1981 19-03-1981 02-12-1977 29-02-1984 31-01-1979
US 3452041	A 24-06-1969	US 3761476 A	25-09-1973
WO 9731907	A 04-09-1997	AP 780 A AT 205485 T AU 717699 B2 AU 2093597 A BG 102792 A BR 9707786 A CA 2247443 A1 CN 1218460 A , B CZ 9802750 A3 DE 69706658 D1 DE 69706658 T2 DK 888317 T3 EA 1403 B1 EE 9800288 A WO 9731907 A1 EP 0888317 A1 ES 2163125 T3	22-11-1999 15-09-2001 30-03-2000 16-09-1997 31-08-1999 27-07-1999 04-09-1997 02-06-1999 13-01-1999 18-10-2001 20-06-2002 21-01-2002 26-02-2001 15-02-1999 04-09-1997 07-01-1999 16-01-2002

# INTERNATIONAL SEARCH REPORT

Information on patent family members

International Application No

PCT/US 02/36128

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
WO 9731907 A		HK 1015369 A1	15-02-2002
		HR 970110 A1	30-04-1998
		HU 0004845 A2	28-05-2001
		IL 125796 A	14-06-2001
		JP 3255930 B2	12-02-2002
		JP 2000507216 T	13-06-2000
		NO 983940 A	27-10-1998
		NZ 331381 A	23-06-2000
		PL 328871 A1	01-03-1999
		PT 888317 T	28-03-2002
		SI 888317 T1	30-04-2002
		SK 116398 A3	13-04-1999
		TR 9801707 T2	21-12-1998
		US 6294580 B1	25-09-2001
		ZA 9701645 A	10-12-1997
WO 0116120 A	08-03-2001	AU 7073400 A	26-03-2001
		CA 2382966 A1	08-03-2001
		EP 1206457 A1	22-05-2002
		JP 2003508389 T	04-03-2003
		WO 0116120 A1	08-03-2001
		US 2003045558 A1	06-03-2003
		US 6417212 B1	09-07-2002